

SPINES, SKELETONS AND THE STRONG LAW OF LARGE NUMBERS FOR SUPERDIFFUSIONS

BY MAREN ECKHOFF, ANDREAS E. KYPRIANOU AND MATTHIAS WINKEL

University of Bath, University of Bath and University of Oxford

Consider a supercritical superdiffusion $(X_t)_{t \geq 0}$ on a domain $D \subseteq \mathbb{R}^d$ with branching mechanism

$$(x, z) \mapsto -\beta(x)z + \alpha(x)z^2 + \int_{(0, \infty)} (e^{-zy} - 1 + zy)\Pi(x, dy).$$

The skeleton decomposition provides a pathwise description of the process in terms of immigration along a branching particle diffusion. We use this decomposition to derive the strong law of large numbers (SLLN) for a wide class of superdiffusions from the corresponding result for branching particle diffusions. That is, we show that for suitable test functions f and starting measures μ ,

$$\frac{\langle f, X_t \rangle}{P_\mu[\langle f, X_t \rangle]} \rightarrow W_\infty \quad P_\mu\text{-almost surely as } t \rightarrow \infty,$$

where W_∞ is a finite, non-deterministic random variable characterized as a martingale limit. Our method is based on skeleton and spine techniques and offers structural insights into the driving force behind the SLLN for superdiffusions. The result covers many of the key examples of interest and, in particular, proves a conjecture by Fleischmann and Swart [*Stochastic Process. Appl.* **106** (2003) 141–165] for the super-Wright–Fisher diffusion.

1. Introduction. The asymptotic behavior of the total mass assigned to a compact set by a superprocess was first characterized by Pinsky [51] at the level of the first moment. Motivated by this study, Engländer and Turaev [21] proved weak convergence of the ratio of the total mass in a compact set and its expectation. Others have further improved the mode of convergence; specifically, several authors conjectured an almost sure convergence result for a wide class of superprocesses [15, 22, 30, 46]. However, up to now it has not been possible to deal with many of the classical examples of interest. In the existing literature, for almost sure convergence, either motion and branching mechanism have to obey restrictive conditions [7], or the domain is assumed to be of finite Lebesgue measure [46]. In this article, we make a significant step toward closing the gap and establish the strong law of

Received September 2013; revised May 2014.

MSC2010 subject classifications. Primary 60J68; secondary 60J80, 60F15.

Key words and phrases. Superdiffusion, measure-valued diffusion, skeleton decomposition, spine decomposition, strong law of large numbers, additive and multiplicative martingales, almost sure limit theorem.

large numbers (SLLN) for a large class of superdiffusions on arbitrary domains. In particular, we prove a conjecture by Fleischmann and Swart for the super-Wright–Fisher diffusion.

Methodologically, previous articles concerned with almost sure limit behavior of superprocesses relied on Fourier analysis, functional analytic arguments or used the martingale formulation for superprocesses combined with stochastic analysis. We take a different approach. The core of our proof is the skeleton decomposition that represents the superprocess as an immigration process along a branching particle process, called the skeleton, where immigration occurs in a Poissonian way along the space–time trajectories and at the branch points of the skeleton. The skeleton may be interpreted as immortal particles that determine the long-term behavior of the process. We exploit this fact and carry the SLLN from the skeleton over to the superprocess. Apart from the result itself, this approach provides insights into the driving force behind the law of large numbers for superprocesses.

A more detailed literature review and discussion of the ideas of proof is deferred to Sections 1.4 and 1.5. Before we introduce the model in Section 1.1, our assumptions are stated in Section 1.2, and the main results are collected in Section 1.3.

1.1. *Model and notation.* Let $d \in \mathbb{N}$, and let $D \subseteq \mathbb{R}^d$ be a non-empty domain. For $k \in \mathbb{N}_0$, $\eta > 0$, we write $C^{k,\eta}(D)$ for the space of real-valued functions on D , whose k th order partial derivatives are locally η -Hölder continuous, $C^\eta(D) := C^{0,\eta}(D)$. We denote by $\mathcal{B}(D)$ the Borel σ -algebra on D . The notation $B \subset\subset D$ means that $B \in \mathcal{B}(D)$ is bounded, and there is an open set B_1 such that $B \subseteq B_1 \subseteq \overline{B_1} \subseteq D$. The Lebesgue measure on $\mathcal{B}(D)$ is denoted by ℓ ; the set of finite (and compactly supported) measures on $\mathcal{B}(D)$ is denoted by $\mathcal{M}_f(D)$ [and $\mathcal{M}_c(D)$, resp.]. When μ is a measure on $\mathcal{B}(D)$ and $f : D \rightarrow \mathbb{R}$ measurable, let $\langle f, \mu \rangle := \int_D f(x)\mu(dx)$, whenever the right-hand side makes sense. If μ has a density ρ with respect to ℓ , we write $\langle f, \rho \rangle = \langle f, \mu \rangle$. For any metric space E , we denote by $p(E)$ and $b(E)$ the sets of Borel measurable and, respectively, nonnegative and bounded functions on E , and let $bp(E) = b(E) \cap p(E)$.

Let $(\xi = (\xi_t)_{t \geq 0}; (\mathbb{P}_x)_{x \in D})$ be a diffusion process on D with generator

$$L(x) = \frac{1}{2} \nabla \cdot a(x) \nabla + b(x) \cdot \nabla \quad \text{on } D.$$

The diffusion matrix $a : D \rightarrow \mathbb{R}^{d \times d}$ takes values in the set of symmetric, positive definite matrices. Moreover, all components of a and $b : D \rightarrow \mathbb{R}^d$ belong to $C^{1,\eta}(D)$ for some $\eta \in (0, 1]$ (the parameter η remains fixed throughout the article). In other words, ξ denotes the unique solution to the generalized martingale problem associated with L on $D \cup \{\dagger\}$, the one-point compactification of D with cemetery state \dagger ; cf. Chapter I in [50]. We write $\tau_D = \inf\{t \geq 0 : \xi_t \notin D\}$.

Let $\beta \in C^\eta(D)$ be bounded and

$$(1.1) \quad \psi_0(x, z) := \alpha(x)z^2 + \int_{(0,\infty)} (e^{-zy} - 1 + zy)\Pi(x, dy),$$

where $\alpha \in bp(D)$, and Π is a kernel from D to $(0, \infty)$ such that $x \mapsto \int_{(0, \infty)} (y \wedge y^2) \Pi(x, dy)$ belongs to $bp(D)$. The function $\psi_\beta(x, z) := -\beta(x)z + \psi_0(x, z)$ is called the *branching mechanism*. If $\Pi \equiv 0$, we say that the branching mechanism is quadratic. In Section 4.2, we explain that our results carry over to a class of quadratic branching mechanisms with unbounded α and β .

The main process of interest in this article is the $(L, \psi_\beta; D)$ -superdiffusion, which we denote by $X = (X_t)_{t \geq 0}$. Its distribution is denoted by P_μ if the process is started in $\mu \in \mathcal{M}_f(D)$. That is, X is an $\mathcal{M}_f(D)$ -valued time-homogeneous Markov process such that for all $\mu \in \mathcal{M}_f(D)$, $f \in bp(D)$ and $t \geq 0$,

$$(1.2) \quad P_\mu[e^{-\langle f, X_t \rangle}] = e^{-\langle u_f(\cdot, t), \mu \rangle},$$

where u_f is the unique nonnegative solution to the *mild equation*

$$(1.3) \quad u(x, t) = S_t f(x) - \int_0^t S_s [\psi_0(\cdot, u(\cdot, t - s))](x) ds$$

for all $(x, t) \in D \times [0, \infty)$. Here $S_t g(x) := \mathbb{P}_x[e^{\int_0^t \beta(\xi_s) ds} g(\xi_t) \mathbb{1}_{\{t < \tau_D\}}]$ for all $g \in p(D)$; that is, $(S_t)_{t \geq 0}$ denotes the semigroup of the differential operator $L + \beta$. Every function g on D is automatically extended to $D \cup \{\dagger\}$ by $g(\dagger) := 0$. Hence

$$S_t g(x) = \mathbb{P}_x[e^{\int_0^t \beta(\xi_s) ds} g(\xi_t)].$$

We refer to ξ as the *underlying motion* or just the *motion* in the space D . Informally, the $\mathcal{M}_f(D)$ -valued process $X = (X_t)_{t \geq 0}$ describes a cloud of infinitesimal particles independently evolving according to the motion ξ and branching in a spatially dependent way according to the branching mechanism ψ_β . The existence of the superprocess X is guaranteed by [11, 28], and it satisfies the branching property; see (1.1) in [28] for a definition. By Theorem 3.1 in [10] or Theorem 2.11 in [28], there is a version of X such that $t \mapsto \langle f, X_t \rangle$ is almost surely right-continuous for all continuous $f \in bp(D)$. We will always work with this version. In most texts, the mild equation (1.3) is written in a slightly different form: instead of $(S_t)_{t \geq 0}$, the semigroup of L is used, and ψ_0 is replaced by ψ_β . Using Feynman–Kac arguments [see Lemma A.1(i) in the Appendix] and Gronwall’s lemma, one easily checks that (for bounded β) the two equations are equivalent.

The main goal of this article is to determine the large-time behavior of

$$(1.4) \quad \frac{\langle f, X_t \rangle}{P_\mu[\langle f, X_t \rangle]}$$

for suitable test functions f and starting measures μ . We say that X satisfies the *strong law of large numbers* (SLLN) if, for all test functions $f \in C_c^+(D)$, $f \neq \mathbf{0}$, the ratio in (1.4) converges to a finite, non-deterministic random variable which is independent of f . Here $C_c^+(D)$ denotes the space of nonnegative, continuous functions of compact support, and $\mathbf{0}$ is the constant function with value 0.

1.2. *Statement of assumptions.* A probabilistic view on supercritical superprocesses is offered by the *skeleton decomposition*. This, by now classical (cf. [3, 5, 9, 19, 27, 43]) decomposition has been studied under a variety of names. It provides a pathwise representation of the superprocess as an immigration process along a supercritical branching particle process, that we call the *skeleton*. The skeleton captures the global behavior of the superprocess, and its discrete nature makes it much more tractable than the superprocess itself. We exploit these facts to establish the SLLN for superdiffusions. Specifically, our fundamental aim is to show that the SLLN for superdiffusions follows as soon as an appropriate SLLN holds for its skeleton. Given the existing knowledge for branching particle processes, this will lead us to a large class of superprocesses for which the SLLN can be stated.

Classically, the skeleton was constructed using the event $\mathcal{E}_{\text{fin}} = \{\exists t \geq 0 : X_t(D) = 0\}$ of extinction after finite time to guide the branching particle process into regions where extinction of the superprocess is unlikely. The key property of \mathcal{E}_{fin} exploited in the skeleton decomposition is that the function $x \mapsto w(x) = -\log P_{\delta_x}(\mathcal{E}_{\text{fin}})$ gives rise to the multiplicative martingale $((e^{-\langle w, X_t \rangle})_{t \geq 0}; P_\mu)$. In the more general setup of the present article, we assume only the existence of such a *martingale function* w .

ASSUMPTION 1 (Skeleton assumption). There exists a function $w \in p(D)$ with $w(x) > 0$ for all $x \in D$,

$$(1.5) \quad \sup_{x \in B} w(x) < \infty \quad \text{for all } B \subset\subset D,$$

$$(1.6) \quad P_\mu[e^{-\langle w, X_t \rangle}] = e^{-\langle w, \mu \rangle} \quad \text{for all } \mu \in \mathcal{M}_c(D), t \geq 0.$$

The martingale function w allows us to define the skeleton as a branching particle diffusion Z , where the spatial movement of each particle is equal in distribution to $(\xi = (\xi_t)_{t \geq 0}; (\mathbb{P}_x^w)_{x \in D})$ with

$$(1.7) \quad \frac{d\mathbb{P}_x^w}{d\mathbb{P}_x} \Big|_{\sigma(\xi_s : s \in [0, t])} = \frac{w(\xi_t)}{w(x)} \exp\left(-\int_0^t \frac{\psi_\beta(\xi_s, w(\xi_s))}{w(\xi_s)} ds\right)$$

on $\{t < \tau_D\}$ for all $t \geq 0$.

We will see in Lemma 2.2 that \mathbb{P}_x^w is well-defined. Each particle dies at spatially dependent rate $q \in p(D)$ and is replaced by a random number of offspring with distribution $(p_k(x))_{k \geq 2}$, where x is the location of its death. The branching rate q and the offspring distribution $(p_k)_{k \geq 2}$ are uniquely identified by

$$(1.8) \quad G(x, s) := q(x) \sum_{k=2}^\infty p_k(x)(s^k - s)$$

$$= \frac{1}{w(x)} (\psi_0(x, w(x)(1-s)) - (1-s)\psi_0(x, w(x)))$$

for all $s \in [0, 1]$ and $x \in D$. The fact that q and $(p_k)_{k \geq 2}$ are well-defined by (1.8) is contained in Theorem 2.3 below. In Section 2.1.1, we define Z on a rich probability space with probability measures \mathbf{P}_μ , $\mu \in \mathcal{M}_f(D)$, where the initial configuration of Z under \mathbf{P}_μ is given by a Poisson random measure with intensity $w(x)\mu(dx)$.

As noted earlier, we are interested in the situation where the skeleton itself satisfies a SLLN. There is a substantial body of literature available that analyzes the long-term behavior of branching particle diffusions. To delimit the regime we want to study, we make two regularity assumptions. A detailed discussion of all assumptions can be found in Section 2.1.

The first condition ensures that the semigroup $(S_t)_{t \geq 0}$ of $L + \beta$ grows precisely exponentially on compactly supported, continuous functions.

ASSUMPTION 2 (Criticality assumption). The second order differential operator $L + \beta$ has positive *generalized principal eigenvalue*

$$(1.9) \quad \begin{aligned} \lambda_c &:= \lambda_c(L + \beta) \\ &:= \inf\{\lambda \in \mathbb{R} : \exists u \in C^{2,\eta}(D), u > 0, (L + \beta - \lambda)u = 0\} > 0. \end{aligned}$$

Moreover, we assume that the operator $L + \beta - \lambda_c$ is *critical*; that is, it does not possess a Green’s function, but there exists $\phi \in C^{2,\eta}(D)$, $\phi > 0$, such that $(L + \beta - \lambda_c)\phi = 0$. In this case, ϕ is unique up to constant multiples and is called the *ground state*. With $L + \beta - \lambda_c$, its formal adjoint is also critical (cf. Pinsky [50]), and the corresponding ground state is denoted by $\tilde{\phi}$. We further assume that $L + \beta - \lambda_c$ is *product L^1 -critical*, that is, $\langle \phi, \tilde{\phi} \rangle < \infty$, and we normalize to obtain $\langle \phi, \tilde{\phi} \rangle = 1$.

Corollary 2.7 below shows that under Assumptions 1 and 2 the process

$$W_t^{\phi/w}(Z) = e^{-\lambda_c t} \langle \phi/w, Z_t \rangle, \quad t \geq 0,$$

is a nonnegative \mathbf{P}_μ -martingale for all $\mu \in \mathcal{M}_f^\phi(D) := \{\mu \in M_f(D) : \langle \phi, \mu \rangle < \infty\}$, and $(W_t^{\phi/w}(Z))_{t \geq 0}$ has an almost sure limit. To have the notation everywhere, we define $W_\infty^{\phi/w}(Z) := \liminf_{t \rightarrow \infty} W_t^{\phi/w}(Z)$.

Our second regularity assumption consists essentially of moment conditions.

ASSUMPTION 3 (Moment assumption). There exists $p \in (1, 2]$ such that

$$(1.10) \quad \sup_{x \in D} \phi(x)\alpha(x) < \infty,$$

$$(1.11) \quad \sup_{x \in D} \phi(x) \int_{(0,1]} y^2 \Pi(x, dy) < \infty,$$

$$(1.12) \quad \sup_{x \in D} \phi(x)^{p-1} \int_{(1,\infty)} y^p \Pi(x, dy) < \infty,$$

$$(1.13) \quad \langle \phi^{p-1}, \phi \tilde{\phi} \rangle < \infty,$$

$$(1.14) \quad \left\langle \int_{(1,\infty)} y^2 e^{-w(\cdot)y} \Pi(\cdot, dy), \phi \tilde{\phi} \right\rangle < \infty.$$

The parameter p remains fixed throughout the article. Assumption 3 is satisfied, for example, when ϕ is bounded and $\sup_{x \in D} \int_{(1,\infty)} y^2 \Pi(x, dy) < \infty$. These second moment conditions appeared in the literature (cf. Section 2.1.3), and we will see several examples in Section 4. However, our results are valid under the weaker conditions of Assumption 3. In Sections 2.1.3 and 4.2, we explain that in the case of a quadratic branching mechanism, only (1.10) is needed.

The SLLN has been proved for a large class of branching particle diffusions. Where it has not been established, yet, we assume a SLLN for the skeleton Z . It will be sufficient to assume convergence along lattice times.

ASSUMPTION 4 (Strong law assumption). For all $\mu \in \mathcal{M}_c(D)$, $\delta > 0$ and continuous $f \in p(D)$ with fw/ϕ bounded,

$$\lim_{n \rightarrow \infty} e^{-\lambda_c n \delta} \langle f, Z_{n\delta} \rangle = \langle f, w \tilde{\phi} \rangle W_\infty^{\phi/w}(Z) \quad \mathbf{P}_\mu\text{-almost surely.}$$

At first, Assumption 4 may look like a strong assumption. However, given Assumptions 1–3, the SLLN for the skeleton has been proved under two additional conditions. The first condition controls the spread of the support of the skeleton when started from a single particle; the second condition is a uniformity assumption on the convergence of an associated ergodic motion (the “spine”) to its stationary distribution. See Theorem 2.13 for details. These conditions hold for a wide class of processes, and we demonstrate this for several key examples in Section 4.

1.3. *Statement of the main results.* Before stating the SLLN for superdiffusion X , we relate the limiting random variable of (1.4) to the limit that appears in Assumption 4. In Corollary 2.7 below we show that under Assumption 2, the process

$$W_t^\phi(X) = e^{-\lambda_c t} \langle \phi, X_t \rangle, \quad t \geq 0,$$

is a nonnegative P_μ -martingale for all $\mu \in \mathcal{M}_f^\phi(D) = \{\mu \in \mathcal{M}_f(D) : \langle \phi, \mu \rangle < \infty\}$, and $(W_t^\phi(X))_{t \geq 0}$ has an almost sure limit. To have the notation everywhere, we define $W_\infty^\phi(X) := \liminf_{t \rightarrow \infty} W_t^\phi(X)$.

PROPOSITION 1.1. *Suppose Assumptions 1, 2, (1.10)–(1.12) hold. For all $\mu \in \mathcal{M}_f^\phi(D)$, the martingales $(W_t^\phi(X))_{t \geq 0}$ and $(W_t^{\phi/w}(Z))_{t \geq 0}$ are bounded in $L^p(\mathbf{P}_\mu)$ and*

$$(1.15) \quad W_\infty^\phi(X) = W_\infty^{\phi/w}(Z) \quad \mathbf{P}_\mu\text{-almost surely.}$$

Recall that ℓ denotes the Lebesgue measure on the domain D . Our main theorem is the following.

THEOREM 1.2. *Suppose Assumptions 1–4 hold. For every $\mu \in \mathcal{M}_f^\phi(D)$, there exists a measurable set Ω_0 such that $P_\mu(\Omega_0) = 1$ and, on Ω_0 , for all ℓ -almost everywhere continuous functions $f \in p(D)$ with f/ϕ bounded,*

$$(1.16) \quad \lim_{t \rightarrow \infty} e^{-\lambda c t} \langle f, X_t \rangle = \langle f, \tilde{\phi} \rangle W_\infty^\phi(X).$$

The convergence in (1.16) also holds in $L^1(P_\mu)$. In particular, $P_\mu[W_\infty^\phi(X)] = \langle \phi, \mu \rangle$.

Even though our main interest is almost sure convergence, Theorem 1.2 also implies new results for convergence in probability; see the examples in Section 4. We record the following corollary of Theorem 1.2 to present the result in possibly more familiar terms.

COROLLARY 1.3. *Suppose Assumptions 1–4 hold. In the vague topology, $e^{-\lambda c t} X_t \rightarrow W_\infty^\phi(X) \tilde{\phi}$ P_μ -almost surely as $t \rightarrow \infty$. If, in addition, ϕ is bounded away from zero, then the convergence holds in the weak topology P_μ -almost surely.*

Finally, we present the SLLN as announced in (1.4). This makes the comparison between $\langle f, X_t \rangle$ and its mean explicit.

COROLLARY 1.4. *Suppose that Assumptions 1–4 hold. For all $\mu \in \mathcal{M}_f^\phi(D)$, $\mu \neq 0$, $f \in C_c^+(D)$, $f \neq \mathbf{0}$,*

$$\lim_{t \rightarrow \infty} \frac{\langle f, X_t \rangle}{P_\mu[\langle f, X_t \rangle]} = \frac{1}{\langle \phi, \mu \rangle} W_\infty^\phi(X) \quad P_\mu\text{-almost surely and in } L^1(P_\mu).$$

The weak law of large numbers (WLLN), and even the L^1 -convergence in (1.16), can be obtained without assuming the SLLN for the skeleton as the next theorem reveals.

THEOREM 1.5. *Suppose Assumptions 1, 2, (1.10)–(1.13) hold. For all $\mu \in \mathcal{M}_f^\phi(D)$ and $f \in p(D)$ with f/ϕ bounded, the convergence in (1.16) holds in $L^1(P_\mu)$.*

1.4. Literature review. Terminology in the literature is not always consistent, so let us clarify that we refer to branching particle processes and superprocesses as branching diffusions and superdiffusions, respectively, if the underlying motion is a diffusion. Similar wording is used for other classes of underlying motions.

The limit theory of supercritical branching processes has been studied since the 1960s when sharp statements were established for classical finite-type processes [2, 38]. The first result for branching diffusions was due to Watanabe [60] in 1967, who proved an almost sure convergence result for branching Brownian motion and certain one-dimensional motions. The key ingredient to the proof was Fourier analysis, a technique recently used by Wang [59] and Kouritzin and Ren [40] to establish the SLLN for super-Brownian motion. Super-Brownian motion on \mathbb{R}^d with a spatially independent branching mechanism does not fall into the framework of the current article since $L + \beta - \lambda_c$ is not product L^1 -critical in that case. Rather, $\phi = \tilde{\phi} = \mathbf{1}$, where $\mathbf{1}$ denotes the constant function with value 1, and $e^{-\lambda_c t} P_\mu[\langle f, X_t \rangle]$ converges to zero for all $f \in C_c^+(D)$. The missing scaling factor is $t^{d/2}$ and $P_\mu[\langle f, X_t \rangle] \sim (2\pi t)^{-d/2} e^{\lambda_c t} \langle f, \mathbf{1} \rangle \mu(\mathbb{R}^d)$ for $\mu \in \mathcal{M}_c(\mathbb{R}^d)$. Wang’s [59] SLLN for super-Brownian motion takes the form

$$\lim_{t \rightarrow \infty} \frac{\langle f, X_t \rangle}{P_\mu[\langle f, X_t \rangle]} = \frac{W_\infty^\phi(X)}{\mu(\mathbb{R}^d)} \quad P_\mu\text{-almost surely,}$$

with martingale limit $W_\infty^\phi(X) = \lim_{t \rightarrow \infty} e^{-\lambda_c t} \langle \mathbf{1}, X_t \rangle$, for all nontrivial nonnegative continuous functions with compact support, and for $\mu = \delta_x, x \in \mathbb{R}^d$. Watanabe’s argument is thought to be incomplete because the regularity for his argument is not proven; see [59]. Biggins [6] developed a method to show uniform convergence of martingales for branching random walks. Wang combined these arguments with the compact support property of super-Brownian motion started from $\mu \in \mathcal{M}_c(D)$. Kouritzin and Ren [40] proved the SLLN for super-stable processes of index $\alpha \in (0, 2]$ with spatially independent quadratic branching mechanism. The correct scaling factor in this case is $t^{d/\alpha} e^{-\lambda_c t}$. The authors allow any finite starting measure with finite mean and a class of continuous test functions that decrease sufficiently fast at infinity. Fourier-analytic methods were also used by Grummt and Kolb [34] to prove the SLLN for the two-dimensional super-Brownian motion with a single point source; see [29] for the definition and a proof of existence of this process. Earlier, Engländer [16] established convergence in probability for a class of superdiffusions that do not necessarily satisfy Assumption 2 using a time-dependent h -transform developed in [22].

In the product L^1 -critical case, the dominant method to prove almost sure limit theorems is due to Asmussen and Hering [1]; Kaplan and Asmussen use a similar method in [37]. The main idea is as follows. For $s, t \geq 0$, write $\mathcal{F}_t = \sigma(X_r : r \leq t)$ and

$$\begin{aligned} e^{-\lambda_c(s+t)} \langle f, X_{s+t} \rangle &= e^{-\lambda_c t} P_\mu[e^{-\lambda_c s} \langle f, X_{s+t} \rangle | \mathcal{F}_t] \\ &\quad + (e^{-\lambda_c(s+t)} \langle f, X_{s+t} \rangle - e^{-\lambda_c t} P_\mu[e^{-\lambda_c s} \langle f, X_{s+t} \rangle | \mathcal{F}_t]) \\ &= \text{CE}_f(s, t) + \text{D}_f(s, t). \end{aligned}$$

Here CE stands for “conditional expectation” and D for “difference”. The first step is to show $\text{D}_f(s, t) \rightarrow 0$ as $t \rightarrow \infty$. This is usually done via a Borel–Cantelli

argument and therefore requires a restriction to lattice times $t = n\delta$. The second step is to show that $\text{CE}_f(s, t)$ behaves like the desired limit for s and t large. This is the hardest part of the proof and usually causes most of the assumptions. The third and last step is to extend the result from lattice to continuous time.

Asmussen and Hering control $\text{CE}_f(s, t)$ for branching particle processes by a uniform Perron–Frobenius condition on the semigroup $(S_t)_{t \geq 0}$. Passage to continuous time is obtained under additional continuity assumptions on process and test functions. Recently, their method was generalized by Engländer et al. [17] to show the SLLN for a class of branching diffusions on arbitrary domains. The authors control $\text{CE}_f(s, t)$ by an assumption that restricts the speed at which particles spread in space and a condition on the rate at which a certain ergodic motion (the “spine”) converges to its stationary distribution.

While Asmussen and Hering’s idea for the proof of SLLNs along lattice times is rather robust and (under certain assumptions) feasible also for superprocesses, the argument used for the transition from lattice to continuous time relies heavily on the finite number of particles in the branching diffusion.

A new approach to almost sure limit theorems for branching processes was introduced by Chen and Shiozawa [8] in the setup of branching symmetric Hunt processes. Amongst other assumptions, a spectral gap condition was used to obtain a Poincaré inequality which constitutes the main ingredient in the proof along lattice times. For the transition to continuous times, the argument from Asmussen and Hering was adapted. Chen et al. [7] proved the first SLLN for superprocesses and relied on the same Poincaré inequality and functional analytic methods for the result along lattice times. For the transition to continuous time, Perkins’s Itô formula for superprocesses [48] was used. Even though their proven SLLN holds on the full domain \mathbb{R}^d , the assumptions on motion and branching mechanism are restrictive in the following way: the motion has to be symmetric (and in the diffusive case must have a uniformly elliptic generator), and the coefficients of the branching mechanism have to satisfy a strict Kato class condition.

The idea to use stochastic analysis was brought much further by Liu et al. [46]. The authors gave a proof which is based entirely on the martingale problem for superprocesses and decomposed the process into three martingale measures. Moreover, they introduced a new technique for the transition from lattice to continuous times based on the resolvent operator and estimates for the hitting probabilities of diffusions. The proof by Liu et al. follows again the three steps of Asmussen and Hering. To control the conditional expectation $\text{CE}_f(s, t)$, they assume that the transition density of the underlying motion is intrinsically ultracontractive, and that the domain D is of finite Lebesgue measure. This assumption excludes most of the classical examples; see Section 4.

To complete our review, we mention that the first law of large numbers for superdiffusions was proven by Engländer and Turaev [21] on the domain $D = \mathbb{R}^d$. The authors use analytic tools from the theory of dynamical systems, in particular properties of invariant curves, to show the convergence in distribution. Besides

classical superdiffusions, the 1-dimensional super-Brownian motion with a single point source is studied.

1.5. *Outline of the proof of Theorem 1.2.* The key to our argument is the skeleton decomposition for the supercritical superprocess X . Intuitively, this representation result states that the superprocess is a cloud of subcritical superdiffusive mass immigrating off a supercritical branching diffusion, the skeleton, which governs the large-time behavior of X . It is important to note that we use the skeleton to make a connection between the asymptotic behavior of a branching diffusion and of the superdiffusion, and we do not use any classical approximation of the superprocess by branching particle systems in a high density limit regime.

Broadly speaking, our proof of Theorem 1.2 follows the three steps of Asmussen and Hering outlined in Section 1.4. However, instead of the full process X we consider only the immigration occurring after time t in the decomposition into conditional expectation CE_f and difference D_f . This immigration is a subprocess of X , and we show that the stated convergence for the full process follows when the subprocess converges to the claimed limit.

Using the tree structure of the skeleton, we can split the immigration that occurs after time t according to the different branches of the skeleton at time t . This fact allows us to appeal to discrete techniques for the analysis of the immigration process. To analyze the conditional expectation CE_f for the immigration after time t , we use the SLLN for the skeleton. After exponential rescaling, the immigration along different branches up to a fixed time s is of constant order and the SLLN for the skeleton describes the asymptotic behavior for large t . Taking the observed time frame s to infinity then adjusts only the constants. To replace the limiting random variable $W_\infty^{\phi/w}(Z)$, coming from the SLLN for the skeleton, by $W_\infty^\phi(X)$, we can, as it turns out, reverse the order in which these limits are taken. Taking first the observed time horizon s to infinity for test function ϕ , we recover the martingale for the skeleton as a consequence of the same invariance property of ϕ that makes $(W_t^\phi(X))_{t \geq 0}$ a martingale.

The analysis of D_f for the immigration after time t is fairly standard, and for the transition from lattice to continuous times we adapt the argument by Liu et al. [46], relying again on the skeleton decomposition. The moment estimates needed for our analysis are obtained using a spine decomposition for the superprocess.

1.6. *Overview.* The outline of the paper is as follows. We start in Section 2.1.1 with an analysis of the skeleton assumption (Assumption 1), and give a detailed description of the skeleton decomposition. In the remainder of Section 2.1 we discuss further basic properties of superprocesses and our other three main assumptions, and we compare them to conditions that appeared in the literature. Section 2.2 contains a spine decomposition for the superprocess X , and the proof that the martingale $(W_t^\phi(X))_{t \geq 0}$ is bounded in L^p .

The proofs of the main results are collected in Section 3. First, in Section 3.1, we reduce the SLLN to a statement that focuses on the main technical difficulty. In Section 3.2 we show that the martingale limits for superprocess and skeleton agree and, in Section 3.3 we prove the WLLN stated in Theorem 1.5. The asymptotic behavior of the immigration process is studied in Section 3.4, and the SLLN along lattice times is established. The transition from lattice to continuous times is performed in Section 3.5, and we conclude our main results.

In Section 4 we provide several examples to illustrate our results. Spatially independent branching mechanisms are discussed in Section 4.1; quadratic branching mechanisms are considered in Section 4.2. In Section 4.3 we study the super-Wright–Fisher diffusion and prove a conjecture by Fleischmann and Swart [30].

Some minor statements needed along the way are proved in the Appendix: Appendix A.1 contains Feynman–Kac-type arguments, and Appendix A.2 discusses a generalized version of the mild equation (1.3) and monotonicity of its solution in domain and test function.

2. Preliminaries. This section is split into two parts. In the first part, we discuss our four main assumptions, and in the second part, we prove that the martingale $(W_t^\phi(X))_{t \geq 0}$ converges in L^p .

2.1. *Basic properties.*

2.1.1. *Skeleton decomposition.* In this section, we work under Assumption 1. The skeleton decomposition for supercritical superprocesses offers a pathwise description of the superprocess in terms of a supercritical branching particle process dressed with an immigration process. Heuristically, one can think of the skeleton as the prolific individuals of the branching process, that is, individuals belonging to infinite lines of descent. The function w assigns a small value to regions that prolific individuals should avoid. If $w(x) = -\log P_{\delta_x}(\mathcal{E})$ for some event \mathcal{E} , then the skeleton particles avoid the behavior specified by \mathcal{E} . Classical examples are the event of extinction in finite time $\mathcal{E}_{\text{fin}} = \{\exists t \geq 0: \langle \mathbf{1}, X_t \rangle = 0\}$ (cf. [19, 27]) and the event of weak extinction $\mathcal{E}_{\text{lim}} = \{\lim_{t \rightarrow \infty} \langle \mathbf{1}, X_t \rangle = 0\}$; cf. [3, 5].

For $f \in p(D)$, we let $\tilde{f}(x, t) = f(x)$ for all $(x, t) \in D \times [0, \infty)$. Dynkin [11] derives the superprocess X from *exit measures* that describe the evolution of mass not only in time but also in space. He showed that for any domain $B \subseteq D$ and $t \geq 0$, there exists a random, finite measure \tilde{X}_t^B on $D \times [0, \infty)$ such that for all $\mu \in \mathcal{M}_f(D)$ and $f \in bp(D)$,

$$(2.1) \quad P_\mu[e^{-\langle \tilde{f}, \tilde{X}_t^B \rangle}] = e^{-\langle \tilde{u}_f^B(\cdot, t), \mu \rangle},$$

where \tilde{u}_f^B is the unique, nonnegative solution to the integral equation

$$(2.2) \quad u(x, t) = \mathbb{P}_x[f(\xi_{t \wedge \tau_B})] - \mathbb{P}_x \left[\int_0^{t \wedge \tau_B} \psi_\beta(\xi_s, u(\xi_s, t - s)) ds \right]$$

for all $(x, t) \in D \times [0, \infty)$, and $\tau_B = \inf\{t \geq 0 : \xi_t \notin B\}$. For $f \in p(D)$, there exists a sequence of functions $f_k \in bp(D)$ such that $f_k \uparrow f$ pointwise. By (2.1), $\tilde{u}_{f_k}^B(x, t)$ is monotonically increasing in k , and we denote the limit by $\tilde{u}_f^B(x, t) \in [0, \infty]$. With this notation, the monotone convergence theorem implies that (2.1) is valid for all $f \in p(D)$. The same argument shows that (1.2) holds for all $f \in p(D)$, and (1.6) implies $u_w = w$. Hence, (1.6) holds for all $\mu \in \mathcal{M}_f(D)$. The superprocess X_t is obtained as a projection of \tilde{X}_t^D restricted to $D \times \{t\}$. Writing $\tilde{w}(x, t) = w(x)$ for $(x, t) \in D \times [0, \infty)$, the Markov property (cf. Theorem I.1.3 [11]) and (1.6) yield for all $\mu \in \mathcal{M}_f(D)$,

$$(2.3) \quad P_\mu[e^{-\langle \tilde{w}, \tilde{X}_t^B \rangle}] = P_\mu[e^{-\langle w, X_t \rangle}] = e^{-\langle w, \mu \rangle},$$

and comparing to (2.1), we deduce that $\tilde{u}_w^B = \tilde{w}$. Now let $B \subset\subset D$. If the support of μ , $\text{supp}(\mu)$, is a subset of B , then \tilde{X}_t^B is supported on the boundary of $B \times [0, t)$; if $\text{supp}(\mu) \subseteq D \setminus B$, then $\tilde{X}_t^B = \mu$ almost surely; cf. Theorem I.1.2 in [11]. In particular, (1.5) implies that \tilde{w} in $\langle \tilde{w}, \tilde{X}_t^B \rangle$ can be interpreted as a bounded function, and we combine (2.3) and (2.2) to obtain

$$(2.4) \quad w(x) = \mathbb{P}_x[w(\xi_{t \wedge \tau_B})] - \mathbb{P}_x\left[\int_0^{t \wedge \tau_B} \psi_\beta(\xi_s, w(\xi_s)) ds\right]$$

for all $(x, t) \in \bar{B} \times [0, \infty)$. Since w is bounded on \bar{B} , the continuity of the diffusion ξ yields that w is continuous on B ; see the argument in the last paragraph of page 708 in [19]. Because B was arbitrary, we conclude:

LEMMA 2.1. *The martingale function w is continuous on D .*

Lemma A.1(i) in the Appendix shows that (2.4) can be transformed into

$$w(x) = \mathbb{P}_x\left[w(\xi_{t \wedge \tau_B}) \exp\left(-\int_0^{t \wedge \tau_B} \frac{\psi_\beta(\xi_s, w(\xi_s))}{w(\xi_s)} ds\right)\right]$$

for all $(x, t) \in \bar{B} \times [0, \infty)$. Hence, for any domain $B \subset\subset D$, $x \in B$,

$$(2.5) \quad w(\xi_{t \wedge \tau_B}) \exp\left(-\int_0^{t \wedge \tau_B} \frac{\psi_\beta(\xi_s, w(\xi_s))}{w(\xi_s)} ds\right), \quad t \geq 0, \quad \text{is a } \mathbb{P}_x\text{-martingale.}$$

Since every nonnegative local martingale is a supermartingale, we conclude that for all $x \in D$,

$$\frac{w(\xi_t)}{w(x)} \exp\left(-\int_0^t \frac{\psi_\beta(\xi_s, w(\xi_s))}{w(\xi_s)} ds\right), \quad t \geq 0, \quad \text{is a } \mathbb{P}_x\text{-supermartingale.}$$

In particular, we deduce the following lemma.

LEMMA 2.2. For every $x \in D$, \mathbb{P}_x^w is a well-defined (sub-)probability measure and $(\xi = (\xi_t)_{t \geq 0}; (\mathbb{P}_x^w)_{x \in D})$ is a (possibly non-conservative) Markov process, which we consider as a Markov process in $D \cup \{\dagger\}$.

If w is bounded, the argument leading to (2.5) is valid for $B = D$, and $(\xi; \mathbb{P}^w)$ is conservative.

To give a description of the skeleton decomposition, we use the martingale function w to construct an auxiliary $\mathcal{M}_f(D)$ -valued Markov process. Let for all $x \in D$, $z \geq 0$ and $f \in p(D)$, $\Pi^*(x, dy) := e^{-w(x)y} \Pi(x, dy)$,

$$\beta^*(x) := \beta(x) - 2\alpha(x)w(x) - \int_{(0, \infty)} (1 - e^{-w(x)y})y \Pi(x, dy),$$

$$\psi_0^*(x, z) := \alpha(x)z^2 + \int_{(0, \infty)} (e^{-zy} - 1 + zy) \Pi^*(x, dy).$$

Since $\beta^*(x) \leq \beta(x)$ for all $x \in D$, β^* is bounded from above. However, it is not clear whether β^* is bounded from below. Hence, the branching mechanism $\psi_{\beta^*}^*(x, z) = -\beta^*(x)z + \psi_0^*(x, z)$ might not satisfy the assumptions from Section 1.1. To overcome this problem, set $\beta_+^* = \max\{\beta^*, 0\}$ and $\beta_-^* = \max\{-\beta^*, 0\}$ so that $\beta^* = \beta_+^* - \beta_-^*$ with β_+^* bounded and β_-^* nonnegative. We write for all $f \in p(D)$,

$$(2.6) \quad S_t^* f(x) := \mathbb{P}_x[e^{-\int_0^t \beta_-^*(\xi_s) ds} e^{\int_0^t \beta_+^*(\xi_s) ds} f(\xi_t)] = \mathbb{P}_x[e^{\int_0^t \beta^*(\xi_s) ds} f(\xi_t)],$$

where $(\xi, (\mathbb{P}_x)_{x \in D})$ is the original diffusion process on D with generator L defined in Section 1.1. Dynkin [11], Theorem I.1.1, showed the existence and uniqueness of the superprocess $X^* = (X_t^*)_{t \geq 0}$ whose motion is given by the diffusion with generator L killed at spatially dependent rate β_-^* , branching mechanism $\psi_{\beta_+^*}^*(x, z) = -\beta_+^*(x)z + \psi_0^*(x, z)$ and domain D . Then X^* is an $\mathcal{M}_f(D)$ -valued, time-homogeneous Markov process such that for all $\mu \in \mathcal{M}_f(D)$, $f \in bp(D)$ and $t \geq 0$,

$$P_\mu[e^{-\langle f, X_t^* \rangle}] = e^{-\langle u_f^*(\cdot, t), \mu \rangle},$$

where u_f^* is the unique nonnegative solution to

$$(2.7) \quad u(x, t) = S_t^* f(x) - \int_0^t S_s^* [\psi_0^*(\cdot, u(\cdot, t - s))](x) ds$$

for all $(x, t) \in D \times [0, \infty)$. Comparing (2.7) and (1.3), we refer to X^* as the $(L, \psi_{\beta^*}^*; D)$ -superprocess. In Appendix A.2, we show that, alternatively, X^* can be obtained as a monotone, distributional limit of superprocesses in the setup of Section 1.1. If $w(x) = -\log P_{\delta_x}(\mathcal{E})$ for a tail event \mathcal{E} with $P_\mu(\mathcal{E}) = e^{-\langle w, \mu \rangle}$ for all $\mu \in \mathcal{M}_f(D)$, then X^* can be obtained from X by conditioning on \mathcal{E} ; that is, the distribution of X_t^* is given by $P_\mu(X_t \in \cdot | \mathcal{E})$; cf. [3, 19, 27, 43]. For our analysis it will be enough to know that on compactly supported, continuous functions, the

semigroup $(S_t^*)_{t \geq 0}$ grows more slowly than the semigroup $(S_t)_{t \geq 0}$, and we prove this fact in Lemma 3.5.

The following theorem is a concise version of the skeleton decomposition at the level of detail that is useful to us. It is based on a result from Kyprianou et al. [43]. We denote by $\mathcal{M}_a^{\text{loc}}$ the set of locally finite integer-valued measures on $\mathcal{B}(D)$.

THEOREM 2.3 (Kyprianou et al. [43]). *There exists a probability space with probability measures $\mathbf{P}_{\mu, \nu}$, $\mu \in \mathcal{M}_f(D)$, $\nu \in \mathcal{M}_a^{\text{loc}}(D)$, that carries the following processes:*

(i) $(Z = (Z_t)_{t \geq 0}; \mathbf{P}_{\mu, \nu})$ is a branching diffusion with motion $(\xi; \mathbb{P}^w)$ defined in (1.7), and branching rate q and offspring distribution $(p_k)_{k \geq 2}$ defined by (1.8) and $\mathbf{P}_{\mu, \nu}(Z_0 = \nu) = 1$.

(ii) $(X^* = (X_t^*)_{t \geq 0}; \mathbf{P}_{\mu, \nu})$ is an $\mathcal{M}_f(D)$ -valued time-homogeneous Markov process such that for every $\mu \in \mathcal{M}_f(D)$, $f \in bp(D)$ and $t \geq 0$,

$$\mathbf{P}_{\mu, \nu}[e^{-\langle f, X_t^* \rangle}] = e^{-\langle u_f^*(\cdot, t), \mu \rangle},$$

where u_f^* is the unique solution to (2.7). Moreover, X^* is independent of Z under $\mathbf{P}_{\mu, \nu}$.

(iii) $(I = (I_t)_{t \geq 0}; \mathbf{P}_{\mu, \nu})$ is an $\mathcal{M}_f(D)$ -valued process such that:

(a) $\mathbf{P}_{\mu, \sum_i \delta_{x_i}}[e^{-\langle f, I_t \rangle}] = \prod_i \mathbf{P}_{\mu, \delta_{x_i}}[e^{-\langle f, I_t \rangle}]$ for all $\mu \in \mathcal{M}_f(D)$, $x_i \in D$, $f \in p(D)$. Moreover, $\mathbf{P}_{\mu, \nu}(I \in \cdot)$ does not depend on μ , $\mathbf{P}_{\mu, \nu}(I_0 = 0) = 1$, and, under $\mathbf{P}_{\mu, \nu}$, (Z, I) is independent of X^* .

(b) $((X, Z) := (X^* + I, Z); \mathbf{P}_{\mu, \nu})$ is a Markov process.

(c) $(X = X^* + I; \mathbf{P}_\mu)$ is equal in distribution to $(X; P_\mu)$, where \mathbf{P}_μ denotes the measure $\mathbf{P}_{\mu, \nu}$ with ν replaced by a Poisson random measure with intensity $w(x)\mu(dx)$.

(d) Under \mathbf{P}_μ , conditionally given X_t , the measure Z_t is a Poisson random measure with intensity $w(x)X_t(dx)$.

We call the probability space from Theorem 2.3 the *skeleton space*. The process I is called *immigration process* or simply *immigration*. As the processes $(X; \mathbf{P}_\mu)$ on the skeleton space and $(X; P_\mu)$ on the generic space have the same distribution, we may, without loss of generality, work on the skeleton space whenever it is convenient. Since the distributions of X^* and I under $\mathbf{P}_{\mu, \nu}$ do not depend on ν and μ , respectively, we sometimes write $\mathbf{P}_{\mu, \bullet}$ or $\mathbf{P}_{\bullet, \nu}$.

Kyprianou et al. [43] identify the immigration process explicitly. We need only the properties listed in Theorem 2.3 but, for definiteness, we now give a full characterization of the immigration process.

Dynkin and Kuznetsov [13] showed that on the canonical space of measure-valued càdlàg functions $\mathbb{D}([0, \infty), \mathcal{M}_f(D))$ for every $x \in D$ there is a unique measure \mathbb{N}_x such that for all $f \in bp(D)$, $t \geq 0$,

$$(2.8) \quad -\log P_{\delta_x}[e^{-\langle f, X_t \rangle}] = \mathbb{N}_x[1 - e^{-\langle f, X_t \rangle}].$$

The corresponding measures associated with the superprocess X^* are denoted by \mathbb{N}_x^* , $x \in D$.

To describe the immigration processes, we use the classical Ulam–Harris notation to uniquely refer to individuals in the genealogical tree \mathcal{T} of Z ; see, for example, page 290 in [35]. For each individual $u \in \mathcal{T}$, we write b_u and d_u for its birth and death times, respectively, and $\{z_u(r) : r \in [b_u, d_u]\}$ for its spatial trajectory. The skeleton space carries the following processes:

(iii.1) $(\mathfrak{a}; \mathbf{P}_{\mu, \nu})$ is a random measure, such that conditional on Z , \mathfrak{a} is a Poisson random measure that issues, for every $u \in \mathcal{T}$, $\mathcal{M}_f(D)$ -valued processes $X^{\mathfrak{a}, u, r} = (X_t^{\mathfrak{a}, u, r})_{t \geq 0}$ along the space–time trajectory $\{(z_u(r), r) : r \in (b_u, d_u)\}$ with rate

$$dr \times \left(2\alpha(z_u(r)) d\mathbb{N}_{z_u(r)}^* + \int_{(0, \infty)} \Pi(z_u(r), dy) y e^{-w(z_u(r))y} \times dP_{y\delta_{z_u(r)}}^* \right),$$

where P_μ^* denotes the distribution of X^* started in μ . Since at most countable many processes $X^{\mathfrak{a}, u, r}$ are not equal to the constant zero measure, immigration at time t that occurred in the form of processes $X^{\mathfrak{a}, u, r}$ until time t can be written as

$$I_t^\mathfrak{a} = \sum_{u \in \mathcal{T}} \sum_{b_u < r \leq d_u \wedge t} X_{t-r}^{\mathfrak{a}, u, r}.$$

The processes $(X^{\mathfrak{a}, u, r} : u \in \mathcal{T}, b_u < r \leq d_u)$ are independent given Z and independent of X^* .

(iii.2) $(\mathfrak{b}; \mathbf{P}_{\mu, \nu})$ is a random measure, such that conditional on Z , \mathfrak{b} issues, for every $u \in \mathcal{T}$, at space–time point $(z_u(d_u), d_u)$ process $X^{\mathfrak{b}, u}$ with law $P_{Y_u \delta_{z_u(d_u)}}^*$. Given that u is replaced by k particles at its death time d_u , the independent random variable Y_u is distributed according to the measure

$$\frac{1}{q(x)w(x)p_k(x)} \left(\alpha(x)w(x)^2 \delta_0(dy) \mathbb{1}_{\{k=2\}} + w(x)^k \frac{y^k}{k!} e^{-w(x)y} \Pi(x, dy) \right) \Big|_{x=z_u(d_u)}.$$

The immigration at time t that occurred in the form of processes $X^{\mathfrak{b}, u}$ until time t is denoted by

$$I_t^\mathfrak{b} = \sum_{u \in \mathcal{T}} \mathbb{1}_{\{d_u \leq t\}} X_{t-d_u}^{\mathfrak{b}, u}.$$

The processes $(X^{\mathfrak{b}, u} : u \in \mathcal{T})$ are independent of X^* and, given Z , are mutually independent and independent of \mathfrak{a} .

The full immigration process is given by $I = I^\mathfrak{a} + I^\mathfrak{b}$.

PROOF OF THEOREM 2.3. Theorem 2.3 generalizes Corollary 6.2 in [43] in three ways. First, the authors choose $w(x) = -\log P_{\delta_x}(\mathcal{E}_{\text{fin}})$ but after defining Z and X^* this choice is not used anymore, and their argument goes through without any changes for a general martingale function w satisfying Assumption 1. Second, the authors assume that w is locally bounded away from zero. Since w is continu-

ous by Lemma 2.1, this condition is automatically satisfied. Finally, Kyprianou et al. enforce additional regularity conditions on the underlying motion to use a comparison principle from the literature in the proof of their Lemma 6.1; see also their Footnote 1. The comparison principle allows them to conclude that the solution \tilde{u}_f^B to (2.2) is increasing in the domain B when the support of f is a subset of B . Lemmas A.1(i) and A.6 show that this monotonicity holds in the more general setup of the present article, too. \square

We introduce notation to refer to the different parts of the skeleton decomposition.

NOTATION 2.4 (Notation for Z). For $t \geq 0$, we write $Z_t = \sum_{i=1}^{N_t} \delta_{\xi_i(t)}$, where N_t denotes the number of skeleton particles at time t and $(\xi_i(t) : i = 1, \dots, N_t)$ their (conveniently ordered) locations. Given $Z_0, (Z^{i,0} : i = 1, \dots, N_0)$ denote the independent subtrees of the skeleton obtained by splitting Z according to the ancestor at time 0. The Markov property implies that $Z^{i,0}$ follows the same distribution as $(Z; \mathbf{P}_{\bullet, \delta_{\xi_i(0)}})$, $i = 1, \dots, N_0$. Under \mathbf{P}_μ with $\mu \in \mathcal{M}_c(D)$, $N_0 = (\mathbf{1}, Z_0)$ is a Poisson random variable with mean $\langle w, \mu \rangle$.

For $t \geq 0$, let \mathcal{F}_t denote the σ -algebra generated by the processes X^*, Z and I up to time t . Using the characterization of the immigration process from Theorem 2.3, we obtain for all $\mu \in \mathcal{M}_f(D)$, $\nu \in \mathcal{M}_a^{\text{loc}}(D)$, $f \in p(D)$ and $s, t \geq 0$,

$$(2.9) \quad \begin{aligned} \mathbf{P}_{\mu, \nu} [e^{-\langle f, X_{s+t} \rangle} | \mathcal{F}_t] &\stackrel{(b)}{=} \mathbf{P}_{X_t, Z_t} [e^{-\langle f, X_s \rangle}] = \mathbf{P}_{X_t, Z_t} [e^{-\langle f, X_s^* + I_s \rangle}] \\ &\stackrel{(a)}{=} \mathbf{P}_{X_t, \bullet} [e^{-\langle f, X_s^* \rangle}] \prod_{i=1}^{N_t} \mathbf{P}_{\bullet, \delta_{\xi_i(t)}} [e^{-\langle f, I_s \rangle}] \end{aligned}$$

$\mathbf{P}_{\mu, \nu}$ -almost surely. Since under \mathbf{P}_μ and given X_t, Z_t is a Poisson random measure with intensity $w(x)X_t(dx)$ by (d), (2.9) holds \mathbf{P}_μ -almost surely when $\mathbf{P}_{\mu, \nu}$ on the left-hand side is replaced by \mathbf{P}_μ . To make use of this identity, we split the immigration process according to the immigration that occurred before time t and the immigration that occurred along different branches of Z after time t .

NOTATION 2.5 (Notation for I). For $t \geq 0$, denote by $I_s^{*,t}$ the immigration at time $s + t$ that occurred along the skeleton before time t ; $I^{*,t} = (I_s^{*,t})_{s \geq 0}$. In addition, for $i \in \{1, \dots, N_t\}$, let $I_s^{i,t}$ denote the immigration at time $s + t$ that occurred along the subtree of the skeleton rooted at the i th particle at time t with location $\xi_i(t)$; $I^{i,t} = (I_s^{i,t})_{s \geq 0}$. We have

$$(2.10) \quad X_{s+t} = X_{s+t}^* + I_s^{*,t} + \sum_{i=1}^{N_t} I_s^{i,t} \quad \text{for all } s, t \geq 0.$$

According to (2.9) and by the Markov property, given \mathcal{F}_t , $(X_{s+t}^* + I_s^{*,t})_{s \geq 0}$ follows the same distribution as (X^*, \mathbf{P}_{X_t}) , and $I^{i,t}$ follows the same distribution as $(I; \mathbf{P}_{\bullet, \delta_{\xi_i(t)}})$, $i = 1, \dots, N_t$. Moreover, given \mathcal{F}_t , the processes $(I^{i,t} : i = 1, \dots, N_t)$ are independent and independent of $I^{*,t}$.

We end this section with a note on terminology. Several different phrases have been used in the literature to refer to the skeleton decomposition. Evans and O’Connell [27] proved the first skeleton decomposition for supercritical superprocesses in the case of a conservative motion (not necessarily a diffusion) and a quadratic, spatially independent branching mechanism with $\alpha, \beta \in (0, \infty)$, and call the result “representation theorem”. Their study was motivated by the “immortal particle representation” derived by Evans [26] for critical superprocesses conditioned on non-extinction. This representation is in terms of a single “immortal particle” that throws off pieces of mass. Evans’s article is part of a cluster of papers that study conditioned superprocesses. Salisbury and Verzani [55] condition the exit measure of a super-Brownian motion to hit n fixed, distinct points on the boundary of a bounded smooth domain. The authors show that the resulting process can be described as the sum of a tree with n leaves that throws off mass in a Poissonian way and of a copy of the unconditioned process, and call this decomposition “backbone representation”. In a follow-up article [56] they consider different conditionings and derive an “immortal particle description” where the guiding object is a tree with possibly infinitely many branches that they call “backbone” or “branching backbone”. Salisbury and Sezer [54] describe the super-Brownian motion conditioned on boundary statistics in terms of a “branching backbone” or “branching backbone system”. Etheridge and Williams [24] represent a critical superprocess with infinite variance conditioned to survive until a fixed time as immigration along a Poisson number of “immortal trees”. An overview of decompositions of conditioned superprocesses was offered by Etheridge [23] using the names “skeleton” and “immortal skeleton”. Back in our setup of supercritical superprocesses, Engländer and Pinsky [19] speak about a “decomposition with immigration” and Fleischmann and Swart [31] construct a “trimmed tree”. For the analysis of continuous-state branching processes, Duquesne and Winkel [9] find a “Galton–Watson forest”. In the corresponding superprocess setup, Berestycki et al. [3] identify the “prolific backbone” and call the representation itself a “backbone decomposition”. The latter phrase has been used several times since [43, 44, 47, 52].

We decided to use the term “skeleton decomposition” for the following reasons. Since the words “backbone” and “spine” are used interchangeably in spoken English, using these two words to mean different things might cause confusion. Furthermore, spine/backbone describes one key, supporting element of an object and does not branch. In contrast, a skeleton carries the entire structure and determines the main features of an object. This is the correct intuition for the spine decomposition and the skeleton decomposition of branching processes as well as the distinction between them.

2.1.2. *Product L^1 -criticality.* The first two moments of the superprocess can be expressed in terms of the underlying motion and the branching mechanism. That is (see, e.g., Proposition 2.7 in [28]), for all $\mu \in \mathcal{M}_f(D)$ and $f \in bp(D)$,

$$(2.11) \quad P_\mu[\langle f, X_t \rangle] = \langle S_t f, \mu \rangle,$$

$$(2.12) \quad \text{Var}_\mu(\langle f, X_t \rangle) = \int_0^t \left\langle S_s \left[\left(2\alpha + \int_{(0,\infty)} y^2 \Pi(\cdot, dy) \right) (S_{t-s} f)^2 \right], \mu \right\rangle ds.$$

Here $\text{Var}_\mu(\langle f, X_t \rangle)$ denotes the variance of $\langle f, X_t \rangle$ under P_μ . By the monotone convergence theorem, the boundedness of f in (2.11) is unnecessary, and (2.12) holds for $f \in p(D)$ as soon as $\langle S_t f, \mu \rangle < \infty$. Similarly, under Assumption 1 and for $\mu \in \mathcal{M}_f(D)$, $f \in bp(D)$, the first two moments of $\langle f, X_t^* \rangle$ [see the discussion around (2.6) for the definitions] can be expressed as

$$(2.13) \quad \mathbf{P}_\mu[\langle f, X_t^* \rangle] = \langle S_t^* f, \mu \rangle,$$

$$(2.14) \quad \text{Var}_\mu(\langle f, X_t^* \rangle) = \int_0^t \left\langle S_s^* \left[\left(2\alpha + \int_{(0,\infty)} y^2 \Pi^*(\cdot, dy) \right) (S_{t-s}^* f)^2 \right], \mu \right\rangle ds.$$

The main purpose of this section is to discuss Assumption 2, that enforces conditions on the operator $L + \beta$ and consequently on its semigroup $(S_t)_{t \geq 0}$ which is the expectation semigroup of X by (2.11). Throughout the section, we suppose that Assumptions 1 and 2 hold. Key features of the local behavior of the superdiffusion X are determined by the generalized principal eigenvalue $\lambda_c = \lambda_c(L + \beta)$. If α and Π are sufficiently smooth and $\lambda_c \leq 0$, then the superdiffusion exhibits *weak local extinction*, that is, the total mass assigned to a compact set by the superprocess tends to zero. For quadratic branching mechanisms this was shown by Pinsky [51], Theorem 6; for general branching mechanisms the proof of Theorem 3(i) in [18] gives the result. This is the reason to assume $\lambda_c > 0$.

The assumption of product L^1 -criticality restricts this article to the situation where the expectation semigroup $(S_t)_{t \geq 0}$ scales precisely exponentially on compactly supported, continuous functions. In general, writing $S_t f(x) = e^{\lambda_c t} \omega_{f,x}(t)$, the limit $\omega_{f,x} := \lim_{t \rightarrow \infty} \omega_{f,x}(t)$ exists for all $f \in C_c^+(D)$, $x \in D$. Product L^1 -criticality is equivalent to $\omega_{f,x} > 0$ for all $f \neq \mathbf{0}$. The alternative is $\omega_{f,x} = 0$ for all f and x ; cf. Theorem 7 in [51] and Appendix A in [22]. Some of the relevant literature for this regime was discussed in Section 1.4. The notion of product L^1 -criticality comes from the criticality theory of second order elliptic operators. See Appendix B of [19] for a good summary and Chapter 4 in [50] for a comprehensive treatment.

By Theorem 4.8.6 in [50], criticality implies that the ground state ϕ is an invariant function of $e^{-\lambda_c t} S_t$, that is, $e^{-\lambda_c t} S_t \phi = \phi$, and we define a conservative diffusion $(\xi = (\xi_t)_{t \geq 0}; (\mathbb{P}_x^\phi)_{x \in D})$ by

$$(2.15) \quad \left. \frac{d\mathbb{P}_x^\phi}{d\mathbb{P}_x} \right|_{\sigma(\xi_s : s \leq t)} = \frac{\phi(\xi_t)}{\phi(x)} e^{\int_0^t (\beta(\xi_s) - \lambda_c) ds} \quad \text{on } \{t < \tau_D\},$$

$$\mathbb{P}_x^\phi[g(\xi_t)] = \phi(x)^{-1} e^{-\lambda_c t} S_t[\phi g](x),$$

for all $x \in D, t \geq 0, g \in p(D)$. Product L^1 -criticality is equivalent to $(\xi = (\xi_t)_{t \geq 0}; (\mathbb{P}_x^\phi)_{x \in D})$ being a positive recurrent diffusion with stationary distribution $\phi(x)\tilde{\phi}(x) dx$, and we call it the *ergodic motion* or the *spine* (as we explain in Section 2.2). In particular (see Theorems 4.3.3 and 4.8.6 in [50]),

$$(2.16) \quad \langle \mathbb{P}^\phi [g(\xi_t)], \phi\tilde{\phi} \rangle = \langle g, \phi\tilde{\phi} \rangle \quad \text{for all } g \in p(D),$$

and for every probability measure π on D and $g \in bp(D)$,

$$(2.17) \quad \langle \mathbb{P}^\phi [g(\xi_t)], \pi \rangle \rightarrow \langle g, \phi\tilde{\phi} \rangle \quad \text{as } t \rightarrow \infty.$$

If, in addition, the initial distribution π is of compact support, then (2.17) holds for all $g \in p(D)$ with $\langle g, \phi\tilde{\phi} \rangle < \infty$. Indeed, for g bounded, (2.17) follows from Theorem 4.9.9 in [50] and the dominated convergence theorem. If the support of $\pi, \text{supp}(\pi)$, is compactly embedded in D , choose a domain $B \subset\subset D$ with $\text{supp}(\mu) \subseteq B$. There exists a constant $C > 0$ such that

$$(2.18) \quad p^\phi(x, y, t) \leq C\phi(y)\tilde{\phi}(y) \quad \text{for all } x \in B, y \in D, t > 1,$$

where $p^\phi(x, y, t)$ denotes the transition density of (ξ, \mathbb{P}^ϕ) and

$$\lim_{t \rightarrow \infty} p^\phi(x, y, t) = \phi(y)\tilde{\phi}(y) \quad \text{for every } x, y \in D;$$

cf. (2.12) and Theorem 1.3(ii) of Pinchover [49]. Hence, (2.17) for $\pi \in \mathcal{M}_c(D)$ and $g \in p(D)$ with $\langle g, \phi\tilde{\phi} \rangle < \infty$ follows from the dominated convergence theorem.

LEMMA 2.6 (Many-to-one lemma for X and Z). *For all $\mu \in \mathcal{M}_f(D), \nu \in \mathcal{M}_a^{\text{loc}}(D)$ and $g \in p(D)$,*

$$(2.19) \quad e^{-\lambda c t} P_\mu[\langle \phi g, X_t \rangle] = \langle \mathbb{P}^\phi [g(\xi_t)], \phi\mu \rangle,$$

$$(2.20) \quad e^{-\lambda c t} \mathbf{P}_{\bullet, \nu} \left[\left\langle \frac{\phi}{w} g, Z_t \right\rangle \right] = \left\langle \mathbb{P}^\phi [g(\xi_t)], \frac{\phi}{w} \nu \right\rangle,$$

$$(2.21) \quad e^{-\lambda c t} \mathbf{P}_\mu \left[\left\langle \frac{\phi}{w} g, Z_t \right\rangle \right] = \langle \mathbb{P}^\phi [g(\xi_t)], \phi\mu \rangle.$$

PROOF. Identity (2.19) follows immediately from (2.11) and (2.15). For (2.20), notice that by (1.8) the local growth rate of Z is given by

$$\beta^Z(x) := q(x) \left(\sum_{k=2}^\infty k p_k(x) - 1 \right) = \partial_s G(x, s)|_{s=1} = \frac{\psi_0(x, w(x))}{w(x)}$$

for all $x \in D$. Using the definition of \mathbb{P}_x^w in (1.7), we obtain for all $x \in D$,

$$\begin{aligned} & \mathbb{P}_x^w \left[e^{\int_0^t \beta^Z(\xi_s) ds} \frac{\phi(\xi_t)}{w(\xi_t)} g(\xi_t) \right] \\ &= w(x)^{-1} \mathbb{P}_x \left[\exp \left(\int_0^t \left(\beta^Z(\xi_s) - \frac{\psi_\beta(\xi_s, w(\xi_s))}{w(\xi_s)} \right) ds \right) \phi(\xi_t) g(\xi_t) \right] \\ &= w(x)^{-1} S_t[\phi g](x) \stackrel{(2.15)}{=} \frac{\phi(x)}{w(x)} e^{\lambda c t} \mathbb{P}_x^\phi [g(\xi_t)]. \end{aligned}$$

Hence, the first moment formula for branching diffusions (see, e.g., Theorem 8.5 in [35]) yields

$$\begin{aligned}
 e^{-\lambda c t} \mathbf{P}_{\bullet, \nu} \left[\left\langle \frac{\phi}{w} g, Z_t \right\rangle \right] \\
 = e^{-\lambda c t} \left\langle \mathbb{P}^w \left[e^{\int_0^t \beta^Z(\xi_s) ds} \frac{\phi(\xi_t)}{w(\xi_t)} g(\xi_t) \right], \nu \right\rangle = \left\langle \mathbb{P}^\phi [g(\xi_t)], \frac{\phi}{w} \nu \right\rangle.
 \end{aligned}$$

Since under \mathbf{P}_μ , initial configuration of Z is given by a Poisson random measure with intensity $w(x)\mu(dx)$, (2.21) follows from (2.20). \square

We record the following consequence of Lemma 2.6.

COROLLARY 2.7. *For all $\mu \in \mathcal{M}_f^\phi(D)$, the processes $((W_t^\phi(X))_{t \geq 0}; P_\mu)$ and $((W_t^{\phi/w}(Z))_{t \geq 0}; \mathbf{P}_\mu)$ are martingales with*

$$P_\mu [W_t^\phi(X)] = \mathbf{P}_\mu [W_t^{\phi/w}(Z)] = \langle \phi, \mu \rangle \quad \text{for all } t \geq 0.$$

PROOF. Since (ξ, \mathbb{P}_x^ϕ) is conservative, the formula for the expectations follows immediately from (2.19) and (2.21). The Markov property of X combined with (2.19) gives the claim for X . The Markov property of Z and (2.20) imply that $(W_t^{\phi/w}(Z))_{t \geq 0}$ is a $\mathbf{P}_{\bullet, \nu}$ -martingale for all $\nu \in \mathcal{M}_a^{\text{loc}}(D)$ with $\langle \phi/w, \nu \rangle < \infty$. Replacing ν by a Poisson random measure with intensity $w(x)\mu(dx)$ completes the proof. \square

Let $\mu \in \mathcal{M}_f^\phi(D)$, $\mu \neq 0$. After dividing the right-hand side of (2.19) by $\langle \phi, \mu \rangle$, the expression can be interpreted as the expectation of $g(\xi_t)$, where ξ is the ergodic motion with starting point randomized according to $\frac{\phi\mu}{\langle \phi, \mu \rangle}$. With this motivation, define for all measurable sets A ,

$$(2.22) \quad \mathbb{P}_{\phi\mu}^\phi(A) := \frac{1}{\langle \phi, \mu \rangle} \langle \mathbb{P}^\phi(A), \phi\mu \rangle.$$

We end this section with a remark for the case that the superprocess is deterministic.

REMARK 2.8. If we have $\ell(\{x \in D : \alpha(x) + \Pi(x, (0, \infty)) > 0\}) = 0$, then (2.11)–(2.12) imply that $\langle f, X_t \rangle = \langle S_t f, \mu \rangle$ for all $t \geq 0$, P_μ -almost surely, for all continuous $f \in bp(D)$. Hence, Assumption 1 cannot be satisfied. However, under Assumption 2, (2.15) and (2.17) imply that for continuous $f \in bp(D)$ with f/ϕ bounded, P_μ -almost surely, as $t \rightarrow \infty$,

$$\begin{aligned}
 e^{-\lambda c t} \langle f, X_t \rangle &= e^{-\lambda c t} \langle S_t f, \mu \rangle = \langle \mathbb{P}^\phi [f(\xi_t)/\phi(\xi_t)], \phi\mu \rangle \\
 &\rightarrow \langle f/\phi, \phi\tilde{\phi} \rangle \langle \phi, \mu \rangle = \langle f, \tilde{\phi} \rangle W_\infty^\phi(X).
 \end{aligned}$$

Now a standard approximation argument shows that the conclusion of Theorem 1.2 holds.

2.1.3. *Moment conditions.* In this section, we discuss Assumption 3 and compare it to the conditions used in the literature. We work under Assumptions 1 and 2. While Assumption 3 seems to be the most useful set of conditions, we prove our results under the following weaker moment assumption.

ASSUMPTION 3'. There exists $p \in (1, 2]$, $\varphi_1, \varphi_2 \in p(D)$, $\sigma_1, \sigma_2, \sigma_3 \in [p, 2]$ and $j_1, j_2 \in \{0, 1\}$ such that

$$(2.23) \quad \sup_{x \in D} \phi(x)^{\sigma_1-1} \alpha(x) < \infty,$$

$$(2.24) \quad \sup_{x \in D} \phi(x)^{\sigma_2-1} \int_{(0, \varphi_1(x)]} y^{\sigma_2} \Pi(x, dy) < \infty,$$

$$(2.25) \quad \sup_{x \in D} \phi(x)^{\sigma_3-1} \int_{(\varphi_1(x), \infty)} y^{\sigma_3} \Pi(x, dy) < \infty,$$

$$(2.26) \quad \langle \phi^{p-1}, \phi \tilde{\phi} \rangle < \infty,$$

$$(2.27) \quad \left\langle \phi^{j_1} \int_{(0, \varphi_2(\cdot)]} y^2 e^{-w(\cdot)y} \Pi(\cdot, dy), \phi \tilde{\phi} \right\rangle < \infty,$$

$$(2.28) \quad \left\langle \phi^{j_2} \int_{(\varphi_2(\cdot), \infty)} y^2 e^{-w(\cdot)y} \Pi(\cdot, dy), \phi \tilde{\phi} \right\rangle < \infty.$$

Assumption 3 is the special case $\varphi_1 = \varphi_2 = \mathbf{1}$, $\sigma_1 = \sigma_2 = 2$, $\sigma_3 = p$ and $j_1 = j_2 = 0$ of Assumption 3'. Notice that with this choice, condition (2.27) trivially holds since $\langle \phi, \tilde{\phi} \rangle < \infty$ and $x \mapsto \int_{(0, 1]} y^2 \Pi(x, dy)$ is a bounded function by the model assumptions in Section 1.1. Therefore, the following theorem generalizes Proposition 1.1 and Theorems 1.2 and 1.5.

THEOREM 2.9. *Suppose Assumptions 1, 2, (2.23)–(2.25) hold and $\mu \in \mathcal{M}_f^\phi(D)$.*

(i) *The martingales $(W_t^\phi(X))_{t \geq 0}$ and $(W_t^{\phi/w}(Z))_{t \geq 0}$ are bounded in $L^p(\mathbf{P}_\mu)$ and $W_\infty^\phi(X) = W_\infty^{\phi/w}(Z)$ \mathbf{P}_μ -almost surely.*

(ii) *Suppose that, in addition, (2.26) holds. For all $f \in p(D)$ with f/ϕ bounded, we have in $L^1(\mathbf{P}_\mu)$*

$$(2.29) \quad \lim_{t \rightarrow \infty} e^{-\lambda ct} \langle f, X_t \rangle = \langle f, \tilde{\phi} \rangle W_\infty^\phi(X).$$

(iii) *If, in addition, Assumptions 3' and 4 hold, then there exists a measurable set Ω_0 with $P_\mu(\Omega_0) = 1$ and, on Ω_0 , for all ℓ -almost everywhere continuous functions $f \in p(D)$ with f/ϕ bounded, the convergence in (2.29) holds.*

The first three moment conditions, (1.10)–(1.12) or (2.23)–(2.25), are used to guarantee that the martingale $(W_t^\phi(X))_{t \geq 0}$ is bounded in L^p ; see Theorem 2.15 below. To the best of our knowledge, even though these conditions may not be opti-

mal, they are the best conditions obtained so far to guarantee L^p -boundedness, $p \in (1, 2)$, for general superprocesses. For the case of a super-Brownian motion, similar conditions were found in [42]. Condition (1.10) appeared as the main moment assumption in [21] and [22] to establish the convergence (2.29) in distribution and in probability, respectively. The two articles that study almost sure convergence in the product L^1 -critical regime (i.e., under Assumption 2) are by Chen et al. [7] and Liu et al. [46]. In both papers, α and ϕ are bounded; hence, (1.10) holds.

The article [7] is restricted to quadratic branching mechanisms; that is, $\Pi \equiv 0$, and (1.11)–(1.12) are trivially satisfied. Liu et al. [46] do not require Π to have a p th moment. The authors show that under their assumptions [D of finite Lebesgue measure and $(S_t)_{t \geq 0}$ intrinsically ultracontractive], the martingale limit $W_\infty^\phi(X)$ is nontrivial if and only if $\langle \int_{(1,\infty)} y \log y \Pi(\cdot, y/\phi), \tilde{\phi} \rangle < \infty$, and they establish their result under this condition. In the alternative case, the martingale limit is zero almost surely, and the stated convergence (1.16) holds trivially.

The fourth assumption, (1.13) or (2.26), is a technical condition. It is only used in Proposition 3.11 to compare the immigration after a large time t , $\sum_{i=1}^{N_t} \langle f, I_s^{i,t} \rangle$, to its expectation $\sum_{i=1}^{N_t} \mathbf{P}_\mu[\langle f, I_s^{i,t} \rangle | \mathcal{F}_t]$. In the previous articles on the SLLN [7, 46], Assumption (1.13) holds since ϕ is bounded.

The technical condition can be avoided using an h -transform. The h -transform for measure-valued diffusions was introduced by Engländer and Pinsky in [19]. For $h \in C^{2,\eta}(D)$, $h > 0$, let

$$(2.30) \quad \begin{aligned} L_0^h &= L + a \frac{\nabla h}{h} \cdot \nabla, & \beta^h(x) &= \frac{(L + \beta)h(x)}{h(x)}, \\ \psi_0^h(x, z) &= \frac{\psi_0(x, h(x)z)}{h(x)}. \end{aligned}$$

If $\beta^h, \alpha h$ and $x \mapsto \int_{(0,\infty)} (y \wedge h(x)y^2) \Pi(x, dy)$ belong to $b(D)$, then $\psi_{\beta^h}^h(x, z) := -\beta^h(x)z + \psi_0^h(x, z)$ satisfies the assumptions from Section 1.1. We denote the space of such functions h by $\mathbb{H}(\psi_\beta)$. An $(L_0^h, \psi_{\beta^h}^h; D)$ -superprocess X^h started in $h(x)\mu(dx)$ can be obtained from an $(L, \psi; D)$ -superprocess X started in μ by setting $X_t^h(dx) := h(x)X_t(dx)$. This result follows immediately from a comparison of the Laplace transforms using the mild equation (1.3) and Corollary 4.1.2 in [50]; see [19] for the computation in the quadratic case. In the following, we superscript all quantities derived from X^h with an h . Clearly, the $(L, \psi; D)$ -superprocess can be recovered from the $(L_0^h, \psi_{\beta^h}^h; D)$ -superprocess by a transform with $1/h$.

LEMMA 2.10. *Let $h \in \mathbb{H}(\psi_\beta)$ and $\mu \in \mathcal{M}_f^\phi(D)$.*

(i) *The operator $L_0^h + \beta^h$ satisfies Assumption 2 with $\phi^h = \phi/h$, $\tilde{\phi}^h = \tilde{\phi}h$ and $\lambda_c^h = \lambda_c$, and the process $(W_t^{\phi^h}(X^h) = e^{-\lambda_c^h t} \langle \phi^h, X_t^h \rangle : t \geq 0; P_{h\mu}^h)$ is a martingale with almost sure limit $W_\infty^{\phi^h}(X^h)$.*

(ii) Suppose (2.29) holds P_μ -almost surely for some $f \in p(D)$. Then

$$(2.31) \quad \lim_{t \rightarrow \infty} e^{-\lambda_c^h t} \langle f/h, X_t^h \rangle = \langle f/h, \tilde{\phi}^h \rangle W_\infty^{\phi^h}(X^h) \quad P_{h\mu}^h\text{-almost surely.}$$

If (2.29) holds in $L^1(P_\mu)$ instead, then (2.31) holds in $L^1(P_{h\mu}^h)$.

PROOF. The first part of the claim was proved by Pinsky [50], Chapter 4. Setting $X^h := hX$, we immediately obtain $W_t^{\phi^h}(X^h) = e^{-\lambda_c^h t} \langle \phi^h, X_t^h \rangle = W_t^\phi(X)$ and, P_μ -almost surely [in $L^1(P_\mu)$, resp.],

$$e^{-\lambda_c^h t} \langle f/h, X_t^h \rangle = e^{-\lambda_c t} \langle f, X_t \rangle \rightarrow \langle f, \tilde{\phi} \rangle W_\infty^\phi(X) = \langle f/h, \tilde{\phi}^h \rangle W_\infty^{\phi^h}(X^h)$$

as $t \rightarrow \infty$. \square

Lemma 2.10 states that Assumption 2 and our results are invariant under h -transforms. The same is true for Assumptions 1 and 4.

LEMMA 2.11. Let $h \in \mathbb{H}(\psi_\beta)$. The $(L_0^h, \psi_{\beta^h}^h; D)$ -superprocess X^h satisfies Assumption 1 with martingale function $w^h = w/h$ and the distribution of the skeleton Z^h under $\mathbf{P}_{h\mu}^h$ agrees with the distribution of Z under \mathbf{P}_μ for all $\mu \in \mathcal{M}_c(D)$. In particular, if X satisfies Assumption 4, then X^h satisfies Assumption 4.

PROOF. The claim follows immediately from the definitions. \square

Exploiting the invariance under h -transforms, we can prove our main results under the following moment assumption.

ASSUMPTION 3''. There exists $p \in (1, 2]$ such that conditions (1.10)–(1.12) and (2.28) for $j_2 = 1, \varphi_2 = \mathbf{1}$ hold and

$$(2.32) \quad \sup_{x \in D} \int_{(1/\phi(x), \infty)} y \Pi(x, dy) < \infty.$$

Crucially, Assumption 3'' does not require $\langle \phi^p, \tilde{\phi} \rangle < \infty$. In the case of a quadratic branching mechanism, only boundedness of $\phi\alpha$ is needed. Condition (2.32) is needed to guarantee that $\phi \in \mathbb{H}(\psi_\beta)$.

THEOREM 2.12. Suppose Assumptions 1, 2, (1.10)–(1.12) and (2.32) hold, and let $\mu \in \mathcal{M}_f^\phi(D)$.

(i) For all $f \in p(D)$ with f/ϕ bounded, the convergence in (2.29) holds in $L^1(P_\mu)$.

(ii) If, in addition, Assumptions 3'' and 4 hold, then there exists a measurable set Ω_0 with $P_\mu(\Omega_0) = 1$ and, on Ω_0 , for all ℓ -almost everywhere continuous functions $f \in p(D)$ with f/ϕ bounded, the convergence in (2.29) holds.

Theorem 2.12 is a consequence of Theorem 2.9, and will not be used until Section 4, after Theorem 2.9 has been proved.

PROOF OF THEOREM 2.12. Part (i): since we have $\beta^\phi = \lambda_c$, $\alpha^\phi = \phi\alpha$ and $\Pi^\phi(x, dy) = \frac{1}{\phi(x)}\Pi(x, dy/\phi(x))$, conditions (1.10), (1.11), (2.32) and the model assumptions in Section 1.1 guarantee that $\phi \in \mathbb{H}(\psi_\beta)$. By Lemma 2.10(i), X^ϕ satisfies Assumption 2 and $\phi^\phi = \mathbf{1}$. Thus (1.10)–(1.12) imply that X^ϕ satisfies (2.23)–(2.26) with $\varphi_1 = \phi$, $\sigma_2 = 2$, $\sigma_3 = p$ and $\sigma_1 \in [p, 2]$ arbitrary. Using Lemma 2.11, we deduce that Theorem 2.9(ii) applies to X^ϕ , and the claim follows from Lemma 2.10(ii).

Part (ii): X^ϕ satisfies (2.27)–(2.28) with $\varphi_2 = \phi$ and arbitrary $j_1, j_2 \in \{0, 1\}$, and Assumption 4 by Lemma 2.11. Hence, Theorem 2.9(iii) applies to X^ϕ and Lemma 2.10(ii) completes the proof for fixed functions f . The existence of a common set Ω_0 will be proved in Lemma 3.4 below. \square

Engländer and Winter [22] proved the convergence (2.29) in probability under the assumption of a quadratic branching mechanisms and (1.10). Their argument can easily be extended to general branching mechanisms. Since the proof relies on an h -transform with $h = \phi$ and second moment estimates, the additional conditions needed for this generalization are (1.11), (1.12) with $p = 2$, and (2.32).

The freedom to choose $p \in (1, 2]$ allows us to analyze processes where $(W_t^\phi(X))_{t \geq 0}$ is bounded in L^p for $p \in (1, 2)$ but not in L^2 . Examples of such processes are given in Section 4. In these cases, not only our almost sure convergence result is new but also the implied convergence in probability result. The main tool to deal with non-integer moments is a spine decomposition presented in Section 2.2, and we are not aware of any other way to obtain these conditions.

The final conditions (2.27)–(2.28) simplify to (1.14) in the case $j_1 = j_2 = 0$, $\varphi_2 = \mathbf{1}$. These assumptions guarantee that the process X^* from the skeleton decomposition has finite second moments (2.14), a fact which is only used in the transition from lattice to continuous times. In particular, the SLLN along lattice times in Theorem 3.13 holds without it. If w is bounded away from zero, for instance when the branching mechanism is spatially independent and the motion is conservative (see Section 4.1), then (1.14) holds automatically. Since Chen et al. [7] consider a quadratic branching mechanism, the conditions automatically hold in their article. In contrast, Liu et al. [46] have no conditions of this type.

In summary, our moment conditions are weaker than those used in [7], but compared to [46], we impose stricter assumptions on the Lévy measure Π , yet allow a much larger class of underlying motions ξ and domains D .

2.1.4. *The strong law of large numbers for the skeleton.* Throughout this section, we suppose that Assumptions 1, 2 and 3' hold. Assumption 4 may look like a strong assumption on first glance. However, we argue that this is not so. The

skeleton decomposition shows that the large-time behavior of the superprocess is guided by the skeleton. This suggests that the total mass the superprocess assigns to a compact ball, will be asymptotically well-behaved if and only if the skeleton carrying the superprocess has asymptotically a well-behaved number of particles in that ball. We write $\mathcal{B}_0(D) := \{B \in \mathcal{B}(D) : \ell(\partial B) = 0\}$. To show that Assumption 4 holds, it suffices to prove that for all $\mu \in \mathcal{M}_c(D)$, $B \in \mathcal{B}_0(D)$,

$$\liminf_{n \rightarrow \infty} e^{-\lambda_c n \delta} \left\langle \frac{\phi}{w} \mathbb{1}_B, Z_{n\delta} \right\rangle \geq \langle \phi \mathbb{1}_B, \tilde{\phi} \rangle W_\infty^{\phi/w}(Z) \quad \mathbf{P}_\mu\text{-almost surely}$$

as we will see in Lemma 3.1(ii) below. Often it is a much easier task to prove the convergence along lattice times than along continuous times.

There are good results in the literature proving SLLNs for branching diffusions. Some of the relevant literature was reviewed in Section 1.4. A nice argument to obtain almost sure asymptotics for spatial branching particle processes from related asymptotic behavior of the spine was found recently by Harris and Roberts [36]. However, they assume a convergence for the spine which usually does not hold in our setup. The theorem we use to verify several examples in Section 4 is based on a result from Engländer et al. [17]. The authors prove the convergence for strictly dyadic branching diffusions along continuous times. We require only convergence along lattice times, but allow a more general branching generator. The following theorem is a version of their result as our proof reveals.

THEOREM 2.13. *Let $\mu \in \mathcal{M}_c(D)$, and assume that for every x in the support of μ the following conditions hold:*

(i) *There is a family of sets $D_t \in \mathcal{B}(D)$, $t \geq 0$, such that for all $\delta > 0$,*

$$\mathbf{P}_{\bullet, \delta_x}(\exists n_0 \in \mathbb{N} : \text{supp}(Z_{n\delta}) \subseteq D_{n\delta} \text{ for all } n \geq n_0) = 1.$$

(ii) *For every $B \subset\subset D$, there exists a constant $K > 0$ such that*

$$(2.33) \quad \sup_{y \in D_t} |\mathbb{P}_y^\phi[\mathbb{1}_B(\xi_{Kt})] - \langle \phi \mathbb{1}_B, \tilde{\phi} \rangle| \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Then, for all $\delta > 0$, $f \in p(D)$ with $f w / \phi$ bounded,

$$\lim_{n \rightarrow \infty} e^{-\lambda_c n \delta} \langle f, Z_{n\delta} \rangle = \langle f, w \tilde{\phi} \rangle W_\infty^{\phi/w}(Z) \quad \mathbf{P}_\mu\text{-almost surely.}$$

PROOF. Using Notation 2.4, we have $Z = \sum_{i=1}^{N_0} Z^{i,0}$, where given \mathcal{F}_0 , $(Z^{i,0} : i = 1, \dots, N_0)$ are independent and $(Z^{i,0}; \mathbf{P}_\mu(\cdot | \mathcal{F}_0))$ is equal in distribution to $(Z; \mathbf{P}_{\bullet, \delta_{\xi_i(0)}})$. In particular, $W_\infty^{\phi/w}(Z) = \sum_{i=1}^{N_0} W_\infty^{\phi/w}(Z^{i,0})$ and

$$\begin{aligned} & \mathbf{P}_\mu \left(\lim_{n \rightarrow \infty} e^{-\lambda_c n \delta} \langle f, Z_{n\delta} \rangle = \langle f, w \tilde{\phi} \rangle W_\infty^{\phi/w}(Z) \right) \\ & \geq \mathbf{P}_\mu \left(\bigcap_{i=1}^{N_0} \left\{ \lim_{n \rightarrow \infty} e^{-\lambda_c n \delta} \langle f, Z_{n\delta}^{i,0} \rangle = \langle f, w \tilde{\phi} \rangle W_\infty^{\phi/w}(Z^{i,0}) \right\} \right) \end{aligned}$$

$$= \mathbf{P}_\mu \left[\prod_{i=1}^{N_0} \mathbf{P}_{\bullet, \delta_{\xi_i(0)}} \left(\lim_{n \rightarrow \infty} e^{-\lambda_c n \delta} \langle f, Z_{n\delta} \rangle = \langle f, w\tilde{\phi} \rangle W_\infty^{\phi/w}(Z) \right) \right].$$

It remains to argue that under the stated assumptions,

$$\mathbf{P}_{\bullet, \delta_x} \left(\lim_{n \rightarrow \infty} e^{-\lambda_c n \delta} \langle f, Z_{n\delta} \rangle = \langle f, w\tilde{\phi} \rangle W_\infty^{\phi/w}(Z) \right)$$

equals 1 for every x in the support of μ . Engländer et al. [17] give a proof of this result for strictly dyadic branching diffusions in two steps. The argument can be generalized as follows. The first step is to show that with $(s_n)_{n \geq 0}$ nonnegative and nondecreasing, and $U_n = e^{-\lambda_c(s_n + \delta n)} \langle f, Z_{s_n + \delta n} \rangle$, the sequence $D_f(s_n, \delta n) = |U_n - \mathbf{P}_{\bullet, \delta_x}[U_n | \sigma(Z_r : r \leq n\delta)]|$ converges to zero. The key to this result is an upper bound on the p th moment of $W_t^{\phi/w}(Z)$ and is obtained via a spine decomposition of the branching diffusion. This would be possible even in our more general setup but is not needed since the required bound follows easily from Theorem 2.9(i). The second step is to show the convergence of $\text{CE}_f(s_n, \delta n) = \mathbf{P}_{\bullet, \delta_x}[U_n | \sigma(Z_r : r \leq n\delta)]$ for $s_n = K\delta n$ to $\langle f, w\tilde{\phi} \rangle W_\infty^{\phi/w}(Z)$. Their assumptions for this convergence are conditions (iii) and (iv) in their Definition 4. Condition (iii) is our condition (i) in Theorem 2.13. From the proof in [17] it is easy to see that their condition (iv) in Definition 4 can be relaxed to our condition (ii), a fact that has also been used in the verification of some examples in [17]. \square

The following lemma is useful in the verification of the conditions of Theorem 2.13 and has been proved by Engländer et al. [17]. We give the main argument for completeness. We denote by $\|\cdot\|$ the ℓ^2 -norm on \mathbb{R}^d .

LEMMA 2.14. *Suppose that for $x \in D$ there are a continuous function $a : [0, \infty) \rightarrow [0, \infty)$ and some $\varepsilon > 0$ such that*

$$(2.34) \quad \mathbb{P}_x^\phi [\mathbb{1}_{\{\|\xi_t\| \geq a(t)\}} w(\xi_t) / \phi(\xi_t)] \leq e^{-(\lambda_c + \varepsilon)t} \quad \text{for all } t \text{ sufficiently large.}$$

Then condition (i) in Theorem 2.13 holds with $D_t = \{y \in D : \|y\| < a(t)\}$. If, in addition, for every $B \subset\subset D$, there is $K > 0$ such that

$$(2.35) \quad \sup_{\|y_1\| < a(t), y_2 \in B} \left| \frac{p^\phi(y_1, y_2, Kt)}{\phi(y_2)\tilde{\phi}(y_2)} - 1 \right| \rightarrow 0 \quad \text{as } t \rightarrow \infty,$$

where p^ϕ denotes the transition density of $(\xi; \mathbb{P}^\phi)$, then condition (ii) in Theorem 2.13 is satisfied.

PROOF. Markov’s inequality and (2.20) yield for all $t \geq 0$,

$$\begin{aligned} \mathbf{P}_{\bullet, \delta_x}(\text{supp}(Z_t) \not\subseteq D_t) &\leq \mathbf{P}_{\bullet, \delta_x}[\langle \mathbb{1}_{D_t^c}, Z_t \rangle] \\ &= e^{\lambda_c t} \frac{\phi(x)}{w(x)} \mathbb{P}_x^\phi [\mathbb{1}_{\{\|\xi_t\| \geq a(t)\}} w(\xi_t) / \phi(\xi_t)]. \end{aligned}$$

The Borel–Cantelli lemma yields the first part of the lemma. The second part follows immediately from the definitions and from $\langle \phi, \tilde{\phi} \rangle < \infty$. \square

We will see in Section 4 that for many of the main examples of superdiffusions the SLLN for the skeleton already follows from Theorem 2.13. For those processes where Assumption 4 has not been proved yet, we believe that the particle nature of the skeleton will make it easier to obtain the SLLN for the skeleton than to derive further convergence statements in the superprocess setup. This article will then allow us to carry results for the branching diffusion over to the superdiffusion. We emphasize that the SLLN for the skeleton is only needed along lattice times and for compactly supported starting measures.

2.2. Spine decomposition. In this section, we use a spine decomposition of X to identify $(W_t^\phi(X))_{t \geq 0}$ as an L^p -bounded martingale, where $p \in (1, 2]$ is determined by Assumption 3'. A similar decomposition has been used for other purposes by Engländer and Kyprianou [18] on bounded subdomains for quadratic branching mechanisms and by Liu et al. [45] in the case $\alpha = \mathbf{0}$. For the one-dimensional super-Brownian motion the spine decomposition was used by Kyprianou et al. [41, 42] to establish L^p -boundedness of martingales closely related to $(W_t^\phi(X))_{t \geq 0}$. Similar arguments have been used in the setup of branching diffusions in [17, 35]. See [18] for an overview of the history of spine decompositions for branching processes. Throughout this section, we suppose that Assumption 2 holds. Further conditions used are stated explicitly. Recall that $\mathcal{M}_f^\phi(D) = \{\mu \in \mathcal{M}_f(D) : \langle \phi, \mu \rangle < \infty\}$.

THEOREM 2.15. *Suppose assumptions (2.23)–(2.25) hold. For all $\mu \in \mathcal{M}_f^\phi(D)$, $((W_t^\phi(X))_{t \geq 0}; P_\mu)$ is an L^p -bounded martingale. In particular, $((W_t^\phi(X))_{t \geq 0}; P_\mu)$ converges in $L^p(P_\mu)$.*

Let $\mu \in \mathcal{M}_f^\phi(D)$, $\mu \neq 0$. We already showed in Corollary 2.7 that $(W_t^\phi(X))_{t \geq 0}$ is a martingale. Hence, it suffices to prove L^p -boundedness, and we can define a new probability measure Q_μ by

$$\frac{dQ_\mu}{dP_\mu} \Big|_{\sigma(X_s : s \in [0, t])} = \frac{W_t^\phi(X)}{\langle \phi, \mu \rangle} \quad \text{for all } t \geq 0.$$

Recall from (2.22) that $(\xi = (\xi_t)_{t \geq 0}; \mathbb{P}_{\phi\mu}^\phi)$ is the ergodic motion with randomized starting point and use (2.15) to obtain

$$(2.36) \quad \mathbb{P}_{\phi\mu}^\phi(A) = \frac{e^{-\lambda c t}}{\langle \phi, \mu \rangle} \langle \mathbb{P} \cdot [e^{\int_0^t \beta(\xi_s) ds} \phi(\xi_t) \mathbb{1}_A], \mu \rangle$$

for all $A \in \sigma(\xi_s : 0 \leq s \leq t)$.

LEMMA 2.16. For all $\mu \in \mathcal{M}_f^\phi(D)$, $\mu \neq 0$, $f, g \in bp(D)$, $t \geq 0$,

$$(2.37) \quad \begin{aligned} Q_\mu \left[e^{-\langle f, X_t \rangle} \frac{\langle \phi g, X_t \rangle}{\langle \phi, X_t \rangle} \right] \\ = P_\mu [e^{-\langle f, X_t \rangle}] \mathbb{P}_{\phi\mu}^\phi \left[g(\xi_t) \exp \left(- \int_0^t \partial_z \psi_0(\xi_s, u_f(\xi_s, t-s)) ds \right) \right]. \end{aligned}$$

Notice that by definition, $\langle \phi, X_t \rangle > 0$, Q_μ -almost surely.

PROOF OF LEMMA 2.16. We prove (2.37) only for g compactly supported since the general case then follows from the monotone convergence theorem. The continuity of ϕ implies that $f + \theta\phi g \in bp(D)$ for all $\theta \geq 0$. We use the definition of Q_μ and interchange differentiation and integration using the dominated convergence theorem to obtain

$$\begin{aligned} Q_\mu \left[e^{-\langle f, X_t \rangle} \frac{\langle \phi g, X_t \rangle}{\langle \phi, X_t \rangle} \right] &= - \frac{e^{-\lambda_c t}}{\langle \phi, \mu \rangle} P_\mu [\partial_\theta |_{\theta=0} e^{-\langle f + \theta\phi g, X_t \rangle}] \\ &= \frac{e^{-\lambda_c t}}{\langle \phi, \mu \rangle} e^{-\langle u_f(\cdot, t), \mu \rangle} \partial_\theta |_{\theta=0} \langle u_{f + \theta\phi g}(\cdot, t), \mu \rangle. \end{aligned}$$

By (1.2), the definition of ψ_β , and (2.36) the claim follows when we have shown that

$$(2.38) \quad \begin{aligned} h_{f,g}(x, t) &:= \partial_\theta |_{\theta=0} u_{f + \theta\phi g}(x, t) \\ &= \mathbb{P}_x \left[\phi(\xi_t) g(\xi_t) \exp \left(- \int_0^t \partial_z \psi_\beta(\xi_s, u_f(\xi_s, t-s)) ds \right) \right] \end{aligned}$$

since integration with respect to μ and differentiation can be interchanged using the dominated convergence theorem. By (1.3), for any $\theta > 0$,

$$\begin{aligned} \frac{u_{f + \theta\phi g}(x, t) - u_f(x, t)}{\theta} \\ = S_t[\phi g](x) - \int_0^t S_s \left[\frac{\psi_0(\cdot, u_{f + \theta\phi g}(\cdot, t-s)) - \psi_0(\cdot, u_f(\cdot, t-s))}{\theta} \right](x) ds. \end{aligned}$$

The Laplace exponent $\theta \mapsto v(\theta) := u_{f + \theta\phi g}(x, t) = -\log P_{\delta_x} [e^{-\langle f + \theta\phi g, X_t \rangle}]$ is increasing, concave and nonnegative. In particular, $\frac{v(\theta) - v(0)}{\theta}$ is decreasing in θ . Moreover, $z \mapsto \psi_0(x, z)$ is increasing, convex and nonnegative. Hence, for all $(x, t) \in D \times [0, \infty)$,

$$0 \leq \frac{v(\theta) - v(0)}{\theta} = \frac{u_{f + \theta\phi g}(x, t) - u_f(x, t)}{\theta} \leq S_t[\phi g](x) \leq \|\phi g\|_\infty e^{\tilde{\beta}t},$$

where $\bar{\beta} = \sup_{x \in D} \beta(x)$ and $\|\cdot\|_\infty$ denotes the supremum norm. A Taylor expansion of ψ_0 yields for every $(x, t) \in D \times [0, \infty)$ some $\tilde{\theta} \in (0, \theta)$ such that

$$\begin{aligned} & \frac{\psi_0(x, v(\theta)) - \psi_0(x, v(0))}{\theta} \\ &= \partial_z \psi_0(x, v(0)) \frac{v(\theta) - v(0)}{\theta} \\ & \quad + (\partial_z \psi_0(x, v(\tilde{\theta})) - \partial_z \psi_0(x, v(0))) \frac{v(\theta) - v(0)}{\theta}. \end{aligned}$$

The first term on the right-hand side is nonnegative and increases as $\theta \downarrow 0$; the second term is dominated and tends to zero. Hence

$$(2.39) \quad \begin{aligned} h_{f,g}(x, t) &= S_t[\phi g](x) \\ & \quad - \int_0^t S_s[\partial_z \psi_0(\cdot, u_f(\cdot, t-s)) h_{f,g}(\cdot, t-s)](x) ds. \end{aligned}$$

Lemma A.1(ii) below applied to the functions $g_1(x, t) = -\partial_z \psi_\beta(x, u_f(x, t))$, $g_2(x, t) = \partial_z \psi_0(x, u_f(x, t))$, $f_1 = \phi g$ and $f_2(x, t) = -\partial_z \psi_0(x, u_f(x, t)) \times h_{f,g}(x, t)$ shows that the unique solution to (2.39) is given by the right-hand side of (2.38). \square

Recall the definition of Dynkin and Kuznetsov’s \mathbb{N}_x -measures from (2.8), and let $\mu \in \mathcal{M}_f^\phi(D)$, $\mu \neq 0$. On a suitable probability space with measure $P_{\mu,\phi}$, we define the following processes:

(i) $(\xi = (\xi_t)_{t \geq 0}; P_{\mu,\phi})$ is equal in distribution to $(\xi_t : t \geq 0; \mathbb{P}_{\phi\mu}^\phi)$, that is an ergodic diffusion. We refer to this process as the *spine*.

(ii) *Continuous immigration*: $(\mathbf{n}; P_{\mu,\phi})$ a random measure such that, given ξ , \mathbf{n} is a Poisson random measure which issues $\mathcal{M}_f(D)$ -valued processes $X^{\mathbf{n},t} = (X_s^{\mathbf{n},t})_{s \geq 0}$ at space–time point (ξ_t, t) with rate $2\alpha(\xi_t) dt \times d\mathbb{N}_{\xi_t}$. The almost surely countable set of immigration times is denoted by $\mathcal{D}^{\mathbf{n}}$; $\mathcal{D}_t^{\mathbf{n}} := \mathcal{D}^{\mathbf{n}} \cap (0, t]$. Given ξ , the processes $(X^{\mathbf{n},t} : t \in \mathcal{D}^{\mathbf{n}})$ are independent.

(iii) *Discontinuous immigration*: $(\mathbf{m}; P_{\mu,\phi})$ a random measure such that, given ξ , \mathbf{m} is a Poisson random measure which issues $\mathcal{M}_f(D)$ -valued processes $X^{\mathbf{m},t}$ at space–time point (ξ_t, t) with rate $dt \times \int_{(0,\infty)} \Pi(\xi_t, dy) y \times dP_{y\delta_{\xi_t}}$. The almost surely countable set of immigration times is denoted by $\mathcal{D}^{\mathbf{m}}$; $\mathcal{D}_t^{\mathbf{m}} = \mathcal{D}^{\mathbf{m}} \cap (0, t]$. Given ξ , the processes $(X^{\mathbf{m},t} : t \in \mathcal{D}^{\mathbf{m}})$ are independent and independent of \mathbf{n} and $(X^{\mathbf{n},t} : t \in \mathcal{D}^{\mathbf{n}})$.

(iv) $(X = (X_t)_{t \geq 0}; P_{\mu,\phi})$ is equal in distribution to $(X = (X_t)_{t \geq 0}; P_\mu)$, that is, it is a copy of the original process. Moreover, X is independent of ξ , \mathbf{n} , \mathbf{m} and all immigration processes.

We denote by

$$X_t^n = \sum_{s \in \mathcal{D}_t^n} X_{t-s}^{n,s} \quad \text{and} \quad X_t^m = \sum_{s \in \mathcal{D}_t^m} X_{t-s}^{m,s}$$

the continuous and discontinuous immigration processes, respectively. We write $\Gamma_t := X_t + X_t^n + X_t^m$ for all $t \geq 0$, and $\stackrel{d}{=}$ denotes distributional equality.

PROPOSITION 2.17 (Spine decomposition). *For all $\mu \in \mathcal{M}_f^\phi(D)$, $\mu \neq 0$,*

$$(X_t : t \geq 0; Q_\mu) \stackrel{d}{=} (\Gamma_t = X_t + X_t^n + X_t^m : t \geq 0; P_{\mu,\phi}).$$

The proof of Proposition 2.17 is very similar to the proof of Theorem 5.2 in [41], and we omit long computations.

PROOF OF PROPOSITION 2.17. Using the definitions and Campbell’s formula for Poisson random measures, one easily checks that the marginal distributions agree. By definition, $((\Gamma_t, \xi_t)_{t \geq 0}; P_{\mu,\phi})$ is a time-homogeneous Markov process, and when we show that

$$P_{\mu,\phi}(\xi_t \in dx | \Gamma_t) = \frac{1}{\langle \phi, \Gamma_t \rangle} \phi(x) \Gamma_t(dx) \quad \text{for all } t \geq 0,$$

then $((\Gamma_t)_{t \geq 0}; P_{\mu,\phi})$ is a time-homogeneous Markov process (by the argument given on page 21 of [41]). Using the definition, $(\Gamma_t; P_{\mu,\phi}) \stackrel{d}{=} (X_t; Q_\mu)$ and Lemma 2.16, we find that for all $f, g \in bp(D)$,

$$P_{\mu,\phi}[e^{-\langle f, \Gamma_t \rangle} P_{\mu,\phi}[g(\xi_t) | \Gamma_t]] = P_{\mu,\phi}\left[e^{-\langle f, \Gamma_t \rangle} \frac{\langle \phi g, \Gamma_t \rangle}{\langle \phi, \Gamma_t \rangle}\right],$$

and the claim follows. \square

For all $t \geq 0$, let \mathcal{G}_t be the σ -algebra generated by ξ up to time t and by \mathbf{n} and \mathbf{m} restricted in the time component to $[0, t]$.

LEMMA 2.18. *For all $\mu \in \mathcal{M}_f^\phi(D)$, $\mu \neq 0$ and $t \geq 0$, $P_{\mu,\phi}$ -almost surely,*

$$P_{\mu,\phi}[e^{-\lambda c t} \langle \phi, \Gamma_t \rangle | \mathcal{G}_t] = \langle \phi, \mu \rangle + \sum_{s \in \mathcal{D}_t^n} e^{-\lambda c s} \phi(\xi_s) + \sum_{s \in \mathcal{D}_t^m} e^{-\lambda c s} \mathcal{I}_s^m \phi(\xi_s),$$

where $(\mathcal{I}_t^m := \langle \mathbf{1}, X_0^{m,t} \rangle : t \geq 0; P_{\mu,\phi})$ is, given ξ , a Poisson point process with intensity measure $dt \times \int_{(0,\infty)} \Pi(\xi_t, dy)y$.

PROOF. Proposition 1.1 of [13] states that, for all test functions $f \in p(D)$ with $P_{\delta_x}[\langle f, X_t \rangle] < \infty$,

$$(2.40) \quad \mathbb{N}_x[\langle f, X_t \rangle] = P_{\delta_x}[\langle f, X_t \rangle].$$

Using first the definition of Γ_t , and then (2.40) and the branching property of X , we obtain

$$\begin{aligned} P_{\mu,\phi} [e^{-\lambda c t} \langle \phi, \Gamma_t \rangle | \mathcal{G}_t] &= P_\mu [W_t^\phi(X)] + \sum_{s \in \mathcal{D}_t^n} e^{-\lambda c t} \mathbb{N}_{\xi_s} [\langle \phi, X_{t-s} \rangle] + \sum_{s \in \mathcal{D}_t^m} e^{-\lambda c t} P_{\mathcal{I}_s^m \delta_{\xi_s}} [\langle \phi, X_{t-s} \rangle] \\ &= P_\mu [W_t^\phi(X)] + \sum_{s \in \mathcal{D}_t^n} e^{-\lambda c t} P_{\delta_{\xi_s}} [\langle \phi, X_{t-s} \rangle] + \sum_{s \in \mathcal{D}_t^m} e^{-\lambda c t} \mathcal{I}_s^m P_{\delta_{\xi_s}} [\langle \phi, X_{t-s} \rangle]. \end{aligned}$$

Since $W_t^\phi(X) = e^{-\lambda c t} \langle \phi, X_t \rangle, t \geq 0$, is a P_μ - and P_{δ_x} -martingale for all $x \in D$, the claim follows. \square

Finally, everything is prepared for the proof of Theorem 2.15. Throughout the article, we use the letters c and C for generic constants in $(0, \infty)$, and their value can change from line to line. Important constants are marked by an index indicating the order in which they occur.

PROOF OF THEOREM 2.15. The martingale property was proved in Corollary 2.7. We have to show the L^p -boundedness. If $\mu \equiv 0$, then $X_t(D) = 0$ for all $t \geq 0$, and the statement is trivially true. Let $\mu \in \mathcal{M}_f^\phi(D)$, $\mu \not\equiv 0$. We write $W_t^\phi(\Gamma) = e^{-\lambda c t} \langle \phi, \Gamma_t \rangle$ and $\bar{p} = p - 1 \in (0, 1]$. Then $x \mapsto x^{\bar{p}}$ is concave and $(x + y)^{\bar{p}} \leq x^{\bar{p}} + y^{\bar{p}}$ for $x, y \geq 0$. Hence, the definition of Q_μ , Proposition 2.17, Jensen's inequality and Lemma 2.18, yield

$$\begin{aligned} \frac{P_\mu [W_t^\phi(X)^p]}{\langle \phi, \mu \rangle} &= Q_\mu [W_t^\phi(X)^{\bar{p}}] = P_{\mu,\phi} [P_{\mu,\phi} [W_t^\phi(\Gamma)^{\bar{p}} | \mathcal{G}_t]] \\ &\leq P_{\mu,\phi} [P_{\mu,\phi} [W_t^\phi(\Gamma) | \mathcal{G}_t]^{\bar{p}}] \\ &\leq P_{\mu,\phi} \left[\langle \phi, \mu \rangle^{\bar{p}} + \left(\sum_{s \in \mathcal{D}_t^n} e^{-\lambda c s} \phi(\xi_s) \right)^{\bar{p}} \right] \\ &\quad + P_{\mu,\phi} \left[\left(\sum_{s \in \mathcal{D}_t^m, \mathcal{I}_s^m \leq \varphi_1(\xi_s)} e^{-\lambda c s} \mathcal{I}_s^m \phi(\xi_s) \right)^{\bar{p}} \right. \\ &\quad \left. + \left(\sum_{s \in \mathcal{D}_t^m, \mathcal{I}_s^m > \varphi_1(\xi_s)} e^{-\lambda c s} \mathcal{I}_s^m \phi(\xi_s) \right)^{\bar{p}} \right], \end{aligned}$$

where φ_1 is determined by Assumption 3'. The first term is deterministic. For the remaining three terms we first use that $x^{\bar{p}} \leq 1 + x^{\bar{\sigma}}$ for all $\bar{\sigma} \geq \bar{p}$, then $(x + y)^{\bar{\sigma}} \leq x^{\bar{\sigma}} + y^{\bar{\sigma}}$ for all $\bar{\sigma} \in [0, 1]$, and finally apply Campbell's formula to obtain

$$P_{\mu,\phi} \left[\left(\sum_{s \in \mathcal{D}_t^n} e^{-\lambda c s} \phi(\xi_s) \right)^{\bar{p}} \right] \leq 1 + \int_0^t 2e^{-\lambda c \bar{\sigma}_1 s} \mathbb{P}_{\phi\mu}^\phi [\phi(\xi_s)^{\bar{\sigma}_1} \alpha(\xi_s)] ds,$$

$$\begin{aligned}
 & P_{\mu, \phi} \left[\left(\sum_{s \in \mathcal{D}_t^{\mathbf{m}}, \mathcal{I}_s^{\mathbf{m}} \leq \varphi_1(\xi_s)} e^{-\lambda_c s} \mathcal{I}_s^{\mathbf{m}} \phi(\xi_s) \right)^{\bar{p}} \right] \\
 & \leq 1 + \int_0^t e^{-\lambda_c \bar{\sigma}_2 s} \mathbb{P}_{\phi \mu}^\phi \left[\int_{(0, \varphi_1(\xi_s))} \phi(\xi_s)^{\bar{\sigma}_2} y^{\bar{\sigma}_2+1} \Pi(\xi_s, dy) \right] ds, \\
 & P_{\mu, \phi} \left[\left(\sum_{s \in \mathcal{D}_t^{\mathbf{m}}, \mathcal{I}_s^{\mathbf{m}} > \varphi_1(\xi_s)} e^{-\lambda_c s} \mathcal{I}_s^{\mathbf{m}} \phi(\xi_s) \right)^{\bar{p}} \right] \\
 & \leq 1 + \int_0^t e^{-\lambda_c \bar{\sigma}_3 s} \mathbb{P}_{\phi \mu}^\phi \left[\int_{(\varphi_1(\xi_s), \infty)} \phi(\xi_s)^{\bar{\sigma}_3} y^{\bar{\sigma}_3+1} \Pi(\xi_s, dy) \right] ds,
 \end{aligned}$$

where $\bar{\sigma}_i = \sigma_i - 1 \in [\bar{p}, 1]$ with σ_i defined in Assumption 3', $i \in \{1, 2, 3\}$. We conclude that if assumptions (2.23)–(2.25) hold, then there exists a constant $C_1 \in (0, \infty)$ independent of μ and t such that

$$(2.41) \quad \frac{P_\mu[W_t^\phi(X)^p]}{\langle \phi, \mu \rangle} \leq \langle \phi, \mu \rangle^{\bar{p}} + C_1 \quad \text{for all } t \geq 0,$$

and $((W_t^\phi(X))_{t \geq 0}; P_\mu)$ is an L^p -bounded martingale. Doob's inequality yields the stated L^p -convergence. \square

In Section 3.3 the following lemma will be used in the comparison between immigration process and its conditional expectation.

LEMMA 2.19. *Suppose assumptions (2.23)–(2.26) hold. For every $\mu \in \mathcal{M}_c(D)$, $\mu \neq 0$, there exists a time $T > 0$ and a constant $C_2 \in (0, \infty)$ such that*

$$\mathbb{P}_{\phi \mu}^\phi [\phi(\xi_t)^{-1} P_{\delta_{\xi_t}} [W_s^\phi(X)^p]] \leq C_2 \quad \text{for all } s \geq 0, t \geq T.$$

PROOF. According to (2.41), for all $s, t \geq 0$,

$$\mathbb{P}_{\phi \mu}^\phi [\phi(\xi_t)^{-1} P_{\delta_{\xi_t}} [W_s^\phi(X)^p]] \leq \mathbb{P}_{\phi \mu}^\phi [\phi(\xi_t)^{p-1}] + C_1.$$

Since $\mu \in \mathcal{M}_c(D)$ and $\langle \phi^{p-1}, \phi \tilde{\phi} \rangle < \infty$ by assumption (2.26), (2.17) implies that $\mathbb{P}_{\phi \mu}^\phi [\phi(\xi_t)^{p-1}]$ converges to $\langle \phi^{p-1}, \phi \tilde{\phi} \rangle$, and the claim follows. \square

3. Proofs of the main results.

3.1. *Reduction to a core statement.* In this section, we work under Assumption 2. We first show that it suffices to consider test functions $f = \phi \mathbb{1}_B =: \phi|_B$ for Borel sets $B \in \mathcal{B}_0(D) = \{B \in \mathcal{B}(D) : \ell(\partial B) = 0\}$ and that we only have to prove that $\liminf_{t \rightarrow \infty} e^{-\lambda_c t} \langle f, X_t \rangle \geq \langle f, \tilde{\phi} \rangle W_\infty^\phi(X)$ instead of the full convergence. The proof is based on standard approximation theory combined with an idea that appeared in Lemma 9 of [1]. We denote by $C_\ell^+(D)$ the space of nonnegative, measurable, ℓ -almost everywhere continuous functions on D .

LEMMA 3.1. *Let $\mu \in \mathcal{M}_f^\phi(D)$ and either $\mathbb{T} = [0, \infty)$ or $\mathbb{T} = \delta\mathbb{N}$ for some $\delta > 0$. In addition, let either $\mathcal{A} = \mathcal{B}_0(D)$ and $\mathcal{A}^\phi = \{f \in C_\ell^+(D) : f/\phi \in b(D)\}$, or $\mathcal{A} = \mathcal{B}(D)$ and $\mathcal{A}^\phi = \{f \in p(D) : f/\phi \in b(D)\}$. We define $\mathcal{A}^{\phi/w}$ like \mathcal{A}^ϕ where ϕ is replaced by ϕ/w .*

(i) *If for all $B \in \mathcal{A}$,*

$$(3.1) \quad \liminf_{\mathbb{T} \ni t \rightarrow \infty} e^{-\lambda ct} \langle \phi|_B, X_t \rangle \geq \langle \phi|_B, \tilde{\phi} \rangle W_\infty^\phi(X) \quad P_\mu\text{-almost surely,}$$

then $\lim_{\mathbb{T} \ni t \rightarrow \infty} e^{-\lambda ct} \langle f, X_t \rangle = \langle f, \tilde{\phi} \rangle W_\infty^\phi(X)$ P_μ -almost surely for all $f \in \mathcal{A}^\phi$.

(ii) *If for all $B \in \mathcal{A}$, $\liminf_{\mathbb{T} \ni t \rightarrow \infty} e^{-\lambda ct} \langle \frac{\phi}{w} \mathbb{1}_B, Z_t \rangle \geq \langle \phi|_B, \tilde{\phi} \rangle W_\infty^{\phi/w}(Z)$ P_μ -almost surely, then, for all $f \in \mathcal{A}^{\phi/w}$, $\lim_{\mathbb{T} \ni t \rightarrow \infty} e^{-\lambda ct} \langle f, Z_t \rangle = \langle f, w\tilde{\phi} \rangle W_\infty^{\phi/w}(Z)$ P_μ -almost surely.*

PROOF. We show only part (i); the proof of part (ii) is similar. Let $\mathcal{S} = \{\sum_{i=1}^k c_i \phi|_{B_i} : k \in \mathbb{N}, c_i \in [0, \infty), B_i \in \mathcal{A}\}$ and $f \in \mathcal{A}^\phi$. There exists a sequence of functions $f_k \in \mathcal{S}$ such that $0 \leq f_k \leq f$ and $f_k \uparrow f$ pointwise. Using (3.1) and the monotone convergence theorem, we deduce that P_μ -almost surely,

$$\begin{aligned} \liminf_{\mathbb{T} \ni t \rightarrow \infty} e^{-\lambda ct} \langle f, X_t \rangle &\geq \sup_{k \in \mathbb{N}} \liminf_{\mathbb{T} \ni t \rightarrow \infty} e^{-\lambda ct} \langle f_k, X_t \rangle \\ &\geq \sup_{k \in \mathbb{N}} \langle f_k, \tilde{\phi} \rangle W_\infty^\phi(X) \\ &= \langle f, \tilde{\phi} \rangle W_\infty^\phi(X). \end{aligned}$$

Let $c = \sup_{x \in D} f(x)/\phi(x)$. Since $0 \leq c\phi - f \leq c\phi$, the same argument can be applied to $c\phi - f$, and we conclude that P_μ -almost surely

$$\begin{aligned} \limsup_{\mathbb{T} \ni t \rightarrow \infty} e^{-\lambda ct} \langle f, X_t \rangle &= \limsup_{\mathbb{T} \ni t \rightarrow \infty} (cW_t^\phi(X) - e^{-\lambda ct} \langle c\phi - f, X_t \rangle) \\ &\leq cW_\infty^\phi(X) - \liminf_{\mathbb{T} \ni t \rightarrow \infty} e^{-\lambda ct} \langle c\phi - f, X_t \rangle \\ &\leq c \langle \phi, \tilde{\phi} \rangle W_\infty^\phi(X) - \langle c\phi - f, \tilde{\phi} \rangle W_\infty^\phi(X) \\ &= \langle f, \tilde{\phi} \rangle W_\infty^\phi(X). \quad \square \end{aligned}$$

In the next step, we use the branching property of the superprocess to restrict ourselves to compactly supported starting measures.

LEMMA 3.2. *Let $\mathbb{T} = [0, \infty)$ or $\mathbb{T} = \delta\mathbb{N}_0$ for some $\delta > 0$, and let $\mathcal{A}^\phi = \{f \in C_\ell^+(D) : f/\phi \in b(D)\}$ or $\mathcal{A}^\phi = \{f \in p(D) : f/\phi \in b(D)\}$.*

(i) *If for all $\mu \in \mathcal{M}_c(D)$ and $f \in \mathcal{A}^\phi$,*

$$(3.2) \quad \lim_{\mathbb{T} \ni t \rightarrow \infty} e^{-\lambda ct} \langle f, X_t \rangle = \langle f, \tilde{\phi} \rangle W_\infty^\phi(X) \quad P_\mu\text{-almost surely,}$$

then (3.2) holds for all $\mu \in \mathcal{M}_f^\phi(D)$.

(ii) If convergence (3.2) holds in $L^1(P_\mu)$ for all $\mu \in \mathcal{M}_c(D)$, then it holds for all $\mu \in \mathcal{M}_f^\phi(D)$.

PROOF. Let $\mu \in \mathcal{M}_f^\phi(D)$, and take a sequence of domains $B_k \subset\subset D$, $B_k \subseteq B_{k+1}$, with $D = \bigcup_{k=1}^\infty B_k$; $\widehat{B}_k := B_k \setminus B_{k-1}$, where $B_0 := \emptyset$. On a suitable probability space, let $X^{\widehat{B}_k}$, $k \in \mathbb{N}$, be independent $(L, \psi_\beta; D)$ -superprocesses, where $X^{\widehat{B}_k}$ is started in $\mathbb{1}_{\widehat{B}_k}\mu$. By the branching property, $X^{B_k} := \sum_{l=1}^k X^{\widehat{B}_l}$, $X^{D \setminus B_k} := \sum_{l=k+1}^\infty X^{\widehat{B}_l}$ and $X := X^{B_k} + X^{D \setminus B_k}$ are $(L, \psi_\beta; D)$ -superprocesses with starting measures $\mathbb{1}_{B_k}\mu$, $\mathbb{1}_{D \setminus B_k}\mu$ and μ , respectively. In particular,

$$W_t^\phi(X) = e^{-\lambda c t} \langle \phi, X_t^{B_k} + X_t^{D \setminus B_k} \rangle = W_t^\phi(X^{B_k}) + W_t^\phi(X^{D \setminus B_k}),$$

and the martingale limits $W_\infty^\phi(X^{D \setminus B_k}) := \liminf_{t \rightarrow \infty} W_t^\phi(X^{D \setminus B_k})$, $k \in \mathbb{N}$, are decreasing in k . Fatou's lemma yields

$$\begin{aligned} P_\mu[W_\infty^\phi(X^{D \setminus B_k})] &= P_\mu\left[\liminf_{t \rightarrow \infty} W_t^\phi(X^{D \setminus B_k})\right] \\ &\leq \liminf_{t \rightarrow \infty} P_\mu[W_t^\phi(X^{D \setminus B_k})] \\ &= \langle \phi, \mathbb{1}_{D \setminus B_k}\mu \rangle. \end{aligned}$$

In particular, $\langle \phi, \mu \rangle < \infty$ implies that $(W_\infty^\phi(X^{D \setminus B_k}) : k \in \mathbb{N})$ converges to zero in $L^1(P_\mu)$ as $k \rightarrow \infty$, and since the sequence is monotonically decreasing, almost sure convergence follows. We conclude that $\lim_{k \rightarrow \infty} W_\infty^\phi(X^{B_k}) = W_\infty^\phi(X)$ almost surely and in $L^1(P_\mu)$.

By Lemma 3.1(i), for part (i) it suffices to show that for all $f \in \mathcal{A}^\phi$,

$$\liminf_{t \rightarrow \infty} e^{-\lambda c t} \langle f, X_t \rangle \geq \langle f, \tilde{\phi} \rangle W_\infty^\phi(X) \quad P_\mu\text{-almost surely.}$$

Since $\mathbb{1}_{B_k}\mu \in \mathcal{M}_c(D)$, the assumption implies that for all $k \in \mathbb{N}$,

$$\liminf_{t \rightarrow \infty} e^{-\lambda c t} \langle f, X_t \rangle \geq \liminf_{t \rightarrow \infty} e^{-\lambda c t} \langle f, X_t^{B_k} \rangle \geq \langle f, \tilde{\phi} \rangle W_\infty^\phi(X^{B_k})$$

P_μ -almost surely, and taking $k \rightarrow \infty$ yields the claim.

To show part (ii), let $c = \sup_{x \in D} f(x)/\phi(x)$ and estimate for fixed $k \in \mathbb{N}$,

$$\begin{aligned} P_\mu[|e^{-\lambda c t} \langle f, X_t \rangle - \langle f, \tilde{\phi} \rangle W_\infty^\phi(X)|] \\ \leq c P_\mu[e^{-\lambda c t} \langle \phi, X_t^{D \setminus B_k} \rangle] + P_\mu[|e^{-\lambda c t} \langle f, X_t^{B_k} \rangle - \langle f, \tilde{\phi} \rangle W_\infty^\phi(X^{B_k})|] \\ + \langle f, \tilde{\phi} \rangle P_\mu[W_\infty^\phi(X^{D \setminus B_k})]. \end{aligned}$$

The second term on the right-hand side tends to zero as $t \rightarrow \infty$ by assumption. The first term is equal to $c \langle \phi, \mathbb{1}_{D \setminus D_k}\mu \rangle$ and, therefore, tends to zero as $k \rightarrow \infty$, and so does the third term. \square

Let $\mathcal{M}(D)$ be the set of all σ -finite measures on D .

REMARK 3.3. The superprocess X can be defined for starting measures $\mu \in \mathcal{M}(D)$ via the branching property; see also Section 1.4.4.1 in [12]. The proof of Lemma 3.2 then shows that (3.2) for all $\mu \in \mathcal{M}_c(D)$ implies (3.2) for all $\mu \in \mathcal{M}(D)$ with $\langle \phi, \mu \rangle < \infty$.

Finally, we show that it suffices to consider fixed test functions. The argument is borrowed from Chen and Shiozawa [8], Theorem 3.7.

LEMMA 3.4 (Chen and Shiozawa [8]). *Let $\mu \in \mathcal{M}_f^\phi(D)$. If for every $B \in \mathcal{B}_0(D)$, P_μ -almost surely, $\lim_{t \rightarrow \infty} e^{-\lambda ct} \langle \phi|_B, X_t \rangle = \langle \phi|_B, \tilde{\phi} \rangle W_\infty^\phi(X)$, then there exists a measurable set Ω_0 such that $P_\mu(\Omega_0) = 1$ and, on Ω_0 , the convergence in (1.16) holds for all $f \in C_\ell^+(D)$ with f/ϕ bounded.*

PROOF. Take a countable base $(B_k)_{k \in \mathbb{N}}$ of $\mathcal{B}_0(D)$ which is closed under finite unions, and let

$$\Omega_0 = \left\{ \lim_{t \rightarrow \infty} e^{-\lambda ct} \langle \phi|_{B_k}, X_t \rangle = \langle \phi|_{B_k}, \tilde{\phi} \rangle W_\infty^\phi(X) \text{ for all } k \in \mathbb{N} \right\}.$$

Then $P_\mu(\Omega_0) = 1$ by assumption. On $\{W_\infty^\phi(X) = 0\}$, convergence (1.16) trivially holds for all $f \in p(D)$ with f/ϕ bounded. On $\{W_\infty^\phi(X) > 0\} \cap \Omega_0$, we define

$$\chi_t(A) := e^{-\lambda ct} \frac{\langle \phi|_A, X_t \rangle}{W_\infty^\phi(X)} \quad \text{and} \quad \chi(A) = \langle \phi|_A, \tilde{\phi} \rangle \quad \text{for all } A \in \mathcal{B}(D).$$

Then χ_t converges vaguely to χ as $t \rightarrow \infty$ and, since $\lim_{t \rightarrow \infty} \chi_t(D) = \chi(D) = 1$, the convergence holds also in the weak sense; cf. Theorem 13.35 in [39]. For every $f \in C_\ell^+(D)$ with f/ϕ bounded, $g := f/\phi \in bp(D)$ is ℓ -almost everywhere continuous and, since χ is absolutely continuous with respect to ℓ , $\lim_{t \rightarrow \infty} \langle g, \chi_t \rangle = \langle g, \chi \rangle$, which is equivalent to

$$\lim_{t \rightarrow \infty} e^{-\lambda ct} \langle f, X_t \rangle = W_\infty^\phi(X) \langle f, \tilde{\phi} \rangle \quad \text{on } \Omega_0 \cap \{W_\infty^\phi(X) > 0\}. \quad \square$$

3.2. *Martingale limits.* In this section, we prove Proposition 1.1; that is, we show that the martingale limits for the superprocess and its skeleton agree almost surely. We assume only Assumptions 1 and 2 throughout this section. Recall from (2.13) that $(S_t^*)_{t \geq 0}$ denotes the expectation semigroup of X^* .

LEMMA 3.5. *Let $f \in p(D)$ with f/ϕ bounded. For all $x \in D$,*

$$\theta_t^*(x) := e^{-\lambda ct} S_t^* f(x) / \phi(x) \rightarrow 0 \quad \text{as } t \rightarrow \infty,$$

and for $t > 0$, the function $x \mapsto \theta_t^(x)$ is continuous. Moreover, $\theta_t^*(x)$ is uniformly bounded in t and x , and if $f = \phi$, $\theta_t^*(x)$ is nonincreasing in t .*

PROOF. Let $c = \sup_{x \in D} f(x)/\phi(x)$. By (2.6) and (2.15), for all $(x, t) \in D \times [0, \infty)$,

$$0 \leq \theta_t^*(x) = \mathbb{P}_x^\phi \left[e^{\int_0^t [\beta^*(\xi_s) - \beta(\xi_s)] ds} f(\xi_t)/\phi(\xi_t) \right] \leq c e^{-\lambda c t} S_t^* \phi(x)/\phi(x).$$

Since $\beta^* - \beta \leq \mathbf{0}$, $e^{-\lambda c t} S_t^* \phi(x)/\phi(x)$ is nonincreasing in t , and Theorem 7.2.4 in [58] (see also Theorem 4.9.7 in [50]) implies that $\theta_t^*(x)$ is continuous in x for $t > 0$. The dominated convergence theorem yields

$$\lim_{t \rightarrow \infty} e^{-\lambda c t} S_t^* \phi(x)/\phi(x) = \mathbb{P}_x^\phi \left[\exp \left(\int_0^\infty [\beta^*(\xi_s) - \beta(\xi_s)] ds \right) \right].$$

By Assumption 2, the diffusion $(\xi = (\xi_t)_{t \geq 0}; \mathbb{P}_x^\phi)$ is positive recurrent, and Theorem 4.9.5(ii) in [50] yields \mathbb{P}_x^ϕ -almost surely,

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t \min\{\beta(\xi_s) - \beta^*(\xi_s), 1\} ds = \langle \min\{\beta - \beta^*, 1\}, \phi \tilde{\phi} \rangle > 0,$$

where the limit is positive since $\ell(\{x \in D : \alpha(x) + \Pi(x, (0, \infty)) > 0\}) > 0$ by Remark 2.8. Hence $\int_0^\infty [\beta^*(\xi_s) - \beta(\xi_s)] ds = -\infty$ holds \mathbb{P}_x^ϕ -almost surely, and the claim is established. \square

The following lemma gives a useful bound for the \hat{p} th moment of $\langle f, I_t \rangle$. We will apply the bound to $\hat{p} = 1$ and to $\hat{p} = p$ with p from Assumption 3'.

LEMMA 3.6. For $\hat{p} \geq 1$, $f \in p(D)$, $x \in D$ and $t \geq 0$,

$$\mathbf{P}_{\bullet, \delta_x}[\langle f, I_t \rangle^{\hat{p}}] \leq w(x)^{-1} \mathbf{P}_{\delta_x}[\langle f, I_t \rangle^{\hat{p}}],$$

where the inequality is an equality in the case $\hat{p} = 1$.

PROOF. Using Notation 2.5, $I_t = \sum_{i=1}^{N_0} I_t^{i,0}$, where under \mathbf{P}_{δ_x} , N_0 is Poisson distributed with mean $w(x)$ and \mathcal{F}_0 -measurable, and $(I_t^{i,0}; \mathbf{P}_{\delta_x}(\cdot | \mathcal{F}_0))$ is equal in distribution to $(I_t; \mathbf{P}_{\bullet, \delta_x})$. Using the monotonicity of the $\ell^{\hat{p}}$ -norm, we derive

$$\begin{aligned} \mathbf{P}_{\delta_x}[\langle f, I_t \rangle^{\hat{p}}] &= \mathbf{P}_{\delta_x} \left[\left(\sum_{i=1}^{N_0} \langle f, I_t^{i,0} \rangle \right)^{\hat{p}} \right] \geq \mathbf{P}_{\delta_x} \left[\sum_{i=1}^{N_0} \langle f, I_t^{i,0} \rangle^{\hat{p}} \right] \\ &= \mathbf{P}_{\delta_x} [N_0 \mathbf{P}_{\bullet, \delta_x}[\langle f, I_t \rangle^{\hat{p}}]] = w(x) \mathbf{P}_{\bullet, \delta_x}[\langle f, I_t \rangle^{\hat{p}}]. \end{aligned}$$

For $\hat{p} = 1$ the inequality is an equality. Rearranging terms completes the proof. \square

We now come to the main part of this section. First, we employ the skeleton decomposition to compute the conditional expectation of $\langle f, X_{s+t} \rangle$.

PROPOSITION 3.7. For all $\mu \in \mathcal{M}_f^\phi(D)$, $f \in p(D)$ with f/ϕ bounded, and $s, t \geq 0$,

$$\mathbf{P}_\mu[\langle f, X_{s+t} \rangle | \mathcal{F}_t] = \langle S_s^* f, X_t \rangle + \left\langle \frac{S_s f}{w}, Z_t \right\rangle - \left\langle \frac{S_s^* f}{w}, Z_t \right\rangle \quad \mathbf{P}_\mu\text{-almost surely.}$$

PROOF. By Notation 2.5, $\langle f, X_{s+t} \rangle = \langle f, X_{s+t}^* + I_s^{*,t} \rangle + \sum_{i=1}^{N_t} \langle f, I_s^{i,t} \rangle$, where $(X_{s+t}^* + I_s^{*,t}; \mathbf{P}_\mu(\cdot | \mathcal{F}_t))$ is equal in distribution to $(X_s^*; \mathbf{P}_{X_t})$ and $(I_s^{i,t}; \mathbf{P}_\mu(\cdot | \mathcal{F}_t))$ to $(I_s; \mathbf{P}_{\bullet, \delta_{\xi_i(t)}})$, $i = 1, \dots, N_t$. Hence, \mathbf{P}_μ -almost surely,

$$\begin{aligned} \mathbf{P}_\mu[\langle f, X_{s+t} \rangle | \mathcal{F}_t] &= \mathbf{P}_\mu \left[\langle f, X_{s+t}^* + I_s^{*,t} \rangle + \sum_{i=1}^{N_t} \langle f, I_s^{i,t} \rangle \middle| \mathcal{F}_t \right] \\ (3.3) \qquad &= \mathbf{P}_{X_t}[\langle f, X_s^* \rangle] + \sum_{i=1}^{N_t} \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle]. \end{aligned}$$

The first term on the right can be rewritten using (2.13) to obtain $\mathbf{P}_{X_t}[\langle f, X_s^* \rangle] = \langle S_s^* f, X_t \rangle$. For the second term, we use Lemma 3.6 and Theorem 2.3 to derive

$$(3.4) \qquad \sum_{i=1}^{N_t} \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle] = \sum_{i=1}^{N_t} w(\xi_i(t))^{-1} \mathbf{P}_{\delta_{\xi_i(t)}}[\langle f, X_s - X_s^* \rangle].$$

Since f/ϕ is bounded and $\mu \in \mathcal{M}_f^\phi(D)$, $\mathbf{P}_{\delta_{\xi_i(t)}}[\langle f, X_s \rangle]$ is finite, and (3.3), (3.4), (2.11) and (2.13) yield \mathbf{P}_μ -almost surely,

$$(3.5) \qquad \mathbf{P}_\mu \left[\sum_{i=1}^{N_t} \langle f, I_s^{i,t} \rangle \middle| \mathcal{F}_t \right] = \sum_{i=1}^{N_t} \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle] = \left\langle \frac{S_s f}{w}, Z_t \right\rangle - \left\langle \frac{S_s^* f}{w}, Z_t \right\rangle$$

as required. \square

PROOF OF PROPOSITION 1.1 AND THEOREM 2.9(i). Proposition 3.7 yields, \mathbf{P}_μ -almost surely,

$$\begin{aligned} \mathbf{P}_\mu[W_{s+t}^\phi(X) | \mathcal{F}_t] &= e^{-\lambda c t} \langle e^{-\lambda c s} S_s^* \phi, X_t \rangle + e^{-\lambda c t} \left\langle e^{-\lambda c s} \frac{S_s \phi}{w}, Z_t \right\rangle \\ (3.6) \qquad &\quad - e^{-\lambda c t} \left\langle e^{-\lambda c s} \frac{S_s^* \phi}{w}, Z_t \right\rangle, \end{aligned}$$

and we are interested in the limit as $s \rightarrow \infty$. The first and last term tend to zero \mathbf{P}_μ -almost surely by Lemma 3.5 and the dominated convergence theorem. The second term is independent of s since $e^{-\lambda c s} S_s \phi = \phi$. Hence, the right-hand side of (3.6) converges to $W_t^{\phi/w}(Z)$. According to Theorem 2.15, $((W_t^\phi(X))_{t \geq 0}; \mathbf{P}_\mu)$ is

an L^p -bounded martingale, and we can interchange on the left-hand side of (3.6) the limit $s \rightarrow \infty$ with the integration to obtain \mathbf{P}_μ -almost surely,

$$(3.7) \quad \mathbf{P}_\mu[W_\infty^\phi(X)|\mathcal{F}_t] = \lim_{s \rightarrow \infty} \mathbf{P}_\mu[W_{s+t}^\phi(X)|\mathcal{F}_t] = W_t^{\phi/w}(Z).$$

Since $W_\infty^\phi(X)$ is measurable with respect to $\mathcal{F}_\infty = \sigma(\cup_{t \geq 0} \mathcal{F}_t)$, (1.15) follows by taking $t \rightarrow \infty$ in (3.7). Moreover, (3.7) shows that $(W_t^{\phi/w}(Z))_{t \geq 0}$ is a uniformly integrable martingale and since $W_\infty^\phi(X) = W_\infty^{\phi/w}(Z)$ is in $L^p(\mathbf{P}_\mu)$, L^p -boundedness of $(W_t^{\phi/w}(Z))_{t \geq 0}$ follows by Jensen’s inequality. \square

3.3. *Convergence in $L^1(P_\mu)$.* In this section, we prove the WLLN in the form of Theorem 1.5. We suppose that Assumptions 1, 2 and (2.23)–(2.26) hold and begin with an L^p -estimate for the immigration that occurred after a large time t . Recall Notations 2.4 and 2.5.

PROPOSITION 3.8. *For every $\mu \in \mathcal{M}_c(D)$ and $f \in p(D)$ with f/ϕ bounded, there exists a time $T > 0$ and a constant $C_3 \in (0, \infty)$ such that for all $s \geq 0, t \geq T$,*

$$e^{-\lambda_c p(s+t)} \mathbf{P}_\mu \left[\left| \sum_{i=1}^{N_t} (\langle f, I_s^{i,t} \rangle - \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle]) \right|^p \right] \leq C_3 e^{-\lambda_c(p-1)t}.$$

PROOF. For $\mu \equiv 0$ the claim is trivial. Let $\mu \not\equiv 0$. It was noted in [6], Lemma 1, that for $p \in [1, 2], n \in \mathbb{N}$ and $(Y_i : i \in \{1, \dots, n\})$ independent, centered random variables (or martingale differences)

$$P \left[\left| \sum_{i=1}^n Y_i \right|^p \right] \leq 2^p \sum_{i=1}^n P[|Y_i|^p].$$

For $s, t \geq 0$, we first apply this inequality to $\mathbf{P}_\mu[\cdot|\mathcal{F}_t], n = N_t$ and $Y_i = \langle f, I_s^{i,t} \rangle - \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle]$, and then use $|x - y|^p \leq x^p + y^p$ for $x, y \geq 0, (I_s^{i,t}; \mathbf{P}_\mu(\cdot|\mathcal{F}_t)) \stackrel{d}{=} (I_s; \mathbf{P}_{\bullet, \delta_{\xi_i(t)}})$ and Jensen’s inequality to obtain

$$\begin{aligned} & \mathbf{P}_\mu \left[\left| \sum_{i=1}^{N_t} (\langle f, I_s^{i,t} \rangle - \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle]) \right|^p \middle| \mathcal{F}_t \right] \\ & \leq 2^p \sum_{i=1}^{N_t} \mathbf{P}_\mu [|\langle f, I_s^{i,t} \rangle - \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle]|^p | \mathcal{F}_t] \\ & \leq 2^p \sum_{i=1}^{N_t} (\mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle^p] + \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle]^p) \\ & \leq 2^{p+1} \sum_{i=1}^{N_t} \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle^p]. \end{aligned}$$

Lemma 3.6, the identity $X_s = X_s^* + I_s$ under $\mathbf{P}_{\delta_{\xi_i(t)}}$ and the monotonicity of $x \mapsto x^p$ on $[0, \infty)$ yield

$$\begin{aligned} & \mathbf{P}_\mu \left[\left| \sum_{i=1}^{N_t} (\langle f, I_s^{i,t} \rangle - \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle]) \right|^p \middle| \mathcal{F}_t \right] \\ & \leq 2^{p+1} \sum_{i=1}^{N_t} \frac{\mathbf{P}_{\delta_{\xi_i(t)}}[\langle f, X_s \rangle^p]}{w(\xi_i(t))} = 2^{p+1} \left\langle \frac{\mathbf{P}_\delta[\langle f, X_s \rangle^p]}{w}, Z_t \right\rangle. \end{aligned}$$

Writing $c = \sup_{x \in D} f(x)/\phi(x) < \infty$, the many-to-one lemma for Z , that is, (2.21), yields

$$\begin{aligned} & e^{-\lambda_c p(s+t)} \mathbf{P}_\mu \left[\left| \sum_{i=1}^{N_t} (\langle f, I_s^{i,t} \rangle - \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle]) \right|^p \right] \\ & \leq 2^{p+1} c^p e^{-\lambda_c p t} \mathbf{P}_\mu \left[\left\langle \frac{\mathbf{P}_\delta[W_s^\phi(X)^p]}{w}, Z_t \right\rangle \right] \\ & = 2^{p+1} c^p e^{-\lambda_c(p-1)t} \langle \phi, \mu \rangle \mathbb{P}_{\phi\mu}^\phi[\phi(\xi_t)^{-1} \mathbf{P}_{\delta_{\xi_t}}[W_s^\phi(X)^p]]. \end{aligned}$$

Since $\mu \in \mathcal{M}_c(D)$, Lemma 2.19 yields a time $T > 0$ and a constant $C_2 \in (0, \infty)$ such that the right-hand side is bounded by $2^{p+1} c^p \langle \phi, \mu \rangle C_2 e^{-\lambda_c(p-1)t}$ for all $t \geq T$, and the proof is complete. \square

We are now in the position to prove Theorems 1.5 and 2.9(ii).

PROOF OF THEOREMS 1.5 AND 2.9(ii). By Lemma 3.2(ii) it suffices to consider $\mu \in \mathcal{M}_c(D)$, and without loss of generality, we work on the skeleton space. Using the skeleton decomposition in the form of (2.10), we write for $s, t \geq 0$,

$$\begin{aligned} & e^{-\lambda_c(s+t)} \langle f, X_{s+t} \rangle - \langle f, \tilde{\phi} \rangle W_\infty^\phi(X) \\ & = e^{-\lambda_c(s+t)} \langle f, X_{s+t}^* + I_s^{*,t} \rangle + e^{-\lambda_c(s+t)} \sum_{i=1}^{N_t} (\langle f, I_s^{i,t} \rangle - \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle]) \\ & \quad + \left(e^{-\lambda_c(s+t)} \sum_{i=1}^{N_t} \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle] - \langle f, \tilde{\phi} \rangle W_t^{\phi/w}(Z) \right) \\ & \quad + (\langle f, \tilde{\phi} \rangle W_t^{\phi/w}(Z) - \langle f, \tilde{\phi} \rangle W_\infty^\phi(X)) \\ & =: \Xi_1(s, t) + \Xi_2(s, t) + \Xi_3(s, t) + \Xi_4(s, t). \end{aligned}$$

It suffices to show that

$$(3.8) \quad \limsup_{s \rightarrow \infty} \limsup_{t \rightarrow \infty} \mathbf{P}_\mu[|\Xi_i(s, t)|] = 0 \quad \text{for all } i \in \{1, \dots, 4\}.$$

We begin with Ξ_1 : since $(X_{s+t}^* + I_s^{*,t}; \mathbf{P}_\mu(\cdot|\mathcal{F}_t)) \stackrel{d}{=} (X_s^*; \mathbf{P}_{X_t})$, the first moment formulas (2.13), (2.11) and (2.15) yield

$$\begin{aligned} \mathbf{P}_\mu[|\Xi_1(s, t)|] &= e^{-\lambda c(s+t)} \mathbf{P}_\mu[\mathbf{P}_{X_t}[\langle f, X_s^* \rangle]] = e^{-\lambda c(s+t)} \langle S_t S_s^* f, \mu \rangle \\ &= \langle \mathbb{P}^\phi[\theta_s^*(\xi_t)], \phi \mu \rangle, \end{aligned}$$

where $\theta_s^*(x) = e^{-\lambda c s} S_s^* f(x) / \phi(x)$. By Lemma 3.5, $\theta_s^*(x)$ is uniformly bounded in s , and x , and converges to zero as $s \rightarrow \infty$. Using the ergodicity of $(\xi; \mathbb{P}^\phi)$ [cf. (2.17)] and the dominated convergence theorem, we conclude

$$\lim_{s \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbf{P}_\mu[|\Xi_1(s, t)|] = \lim_{s \rightarrow \infty} \langle \theta_s^*, \phi \tilde{\phi} \rangle \langle \phi, \mu \rangle = 0.$$

Proposition 3.8 implies that $\Xi_2(s, t)$ converges to zero in $L^p(\mathbf{P}_\mu)$ as $t \rightarrow \infty$ for every fixed $s > 0$. By monotonicity of norms, (3.8) for $i = 2$ follows.

For Ξ_3 , we use (3.5) and (2.15) to rewrite

$$\begin{aligned} e^{-\lambda c(s+t)} \sum_{i=1}^{N_t} \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle f, I_s \rangle] &= e^{-\lambda c(s+t)} \left\langle \frac{S_s f - S_s^* f}{w}, Z_t \right\rangle \\ &= e^{-\lambda c t} \left\langle \mathbb{P}^\phi[f(\xi_s) / \phi(\xi_s)] - \theta_s^*, \frac{\phi}{w} Z_t \right\rangle. \end{aligned}$$

Let $\Upsilon_s(x) := \mathbb{P}_x^\phi[f(\xi_s) / \phi(\xi_s)] - \theta_s^*(x)$. Since f/ϕ is bounded, Υ is uniformly bounded in s and x , and by (2.17) and Lemma 3.5, $\lim_{s \rightarrow \infty} \Upsilon_s(x) = \langle f, \tilde{\phi} \rangle$. Moreover, $\Xi_3(s, t) = e^{-\lambda c t} \langle \Upsilon_s - \langle f, \tilde{\phi} \rangle, \frac{\phi}{w} Z_t \rangle$ by the definition of $W_t^{\phi/w}(Z)$. The many-to-one lemma for Z , that is, (2.21), yields

$$\mathbf{P}_\mu[|\Xi_3(s, t)|] \leq e^{-\lambda c t} \mathbf{P}_\mu \left[\left| \langle \Upsilon_s - \langle f, \tilde{\phi} \rangle, \frac{\phi}{w} Z_t \right| \right] = \langle \mathbb{P}^\phi[|\Upsilon_s(\xi_t) - \langle f, \tilde{\phi} \rangle|], \phi \mu \rangle.$$

Since Υ_s is bounded and $\phi(x)\mu(dx)$ is a finite measure, (2.17) implies that the right-hand side converges to $\langle |\Upsilon_s - \langle f, \tilde{\phi} \rangle|, \phi \tilde{\phi} \rangle \langle \phi, \mu \rangle$ as $t \rightarrow \infty$ and this expression converges to zero as $s \rightarrow \infty$ by the dominated convergence theorem.

Finally, since $(W_t^{\phi/w}(Z))_{t \geq 0}$ is an $L^p(\mathbf{P}_\mu)$ -bounded martingale by Theorem 2.9(i), it converges to $W_\infty^{\phi/w}(Z) = W_\infty^\phi(X)$ in $L^1(\mathbf{P}_\mu)$. Hence, (3.8) for $i = 4$ holds, and the proof is complete. \square

3.4. *Asymptotics for the immigration process and the SLLN along lattice times.*

In this section, we analyze the asymptotic behavior of the immigration process I . According to Lemma 3.1(i), the process X in $e^{-\lambda c t} \langle f, X_t \rangle$ can be replaced by the immigration process when $e^{-\lambda c t} \langle f, I_t \rangle$ converges to the correct limit. To show this, we begin with the conditional expectation of the immigration after a large time t studied in (3.5). We now work under Assumptions 1, 2, 3' and 4.

LEMMA 3.9. For all $\mu \in \mathcal{M}_c(D)$, $s, \delta > 0$ and all $f \in p(D)$ with f/ϕ bounded, \mathbf{P}_μ -almost surely,

$$\lim_{\delta \mathbb{N} \ni t \rightarrow \infty} e^{-\lambda c t} \sum_{i=1}^{N_t} \mathbf{P}_{\bullet, \delta \xi_i(t)} [\langle f, I_s \rangle] = e^{\lambda c s} (\langle f, \tilde{\phi} \rangle - \langle e^{-\lambda c s} S_s^* f, \tilde{\phi} \rangle) W_\infty^\phi(X).$$

PROOF. We apply the SLLN for the skeleton (Assumption 4) to the functions f_1, f_2 given by

$$f_1(x) := \frac{S_s f(x)}{w(x)} = e^{\lambda c s} \mathbb{P}_x^\phi [f(\xi_s)/\phi(\xi_s)] \frac{\phi(x)}{w(x)},$$

$$f_2(x) := \frac{S_s^* f(x)}{w(x)} = e^{\lambda c s} \theta_s^*(x) \frac{\phi(x)}{w(x)}.$$

By Lemmas 2.1 and 3.5 and Theorem 4.9.7 in [50], f_1 and f_2 are continuous. Hence, (3.5) and Assumption 4 yield \mathbf{P}_μ -almost surely,

$$\begin{aligned} & \lim_{\delta \mathbb{N} \ni t \rightarrow \infty} e^{-\lambda c t} \sum_{i=1}^{N_t} \mathbf{P}_{\bullet, \delta \xi_i(t)} [\langle f, I_s \rangle] \\ &= \left\langle \frac{S_s f}{w}, w \tilde{\phi} \right\rangle W_\infty^{\phi/w}(Z) - \left\langle \frac{S_s^* f}{w}, w \tilde{\phi} \right\rangle W_\infty^{\phi/w}(Z). \end{aligned}$$

By (2.15) and (2.16), $\langle S_s f, \tilde{\phi} \rangle = e^{\lambda c s} \langle f, \tilde{\phi} \rangle$ and Theorem 2.9(i) completes the proof. \square

PROPOSITION 3.10. For all $\mu \in \mathcal{M}_c(D)$, $\delta > 0$ and all $f \in p(D)$ with f/ϕ bounded, \mathbf{P}_μ -almost surely,

$$\lim_{\delta \mathbb{N} \ni s \rightarrow \infty} \lim_{\delta \mathbb{N} \ni t \rightarrow \infty} e^{-\lambda c (s+t)} \sum_{i=1}^{N_t} \mathbf{P}_{\bullet, \delta \xi_i(t)} [\langle f, I_s \rangle] = \langle f, \tilde{\phi} \rangle W_\infty^\phi(X).$$

PROOF. The claim follows immediately from Lemmas 3.9 and 3.5 and the dominated convergence theorem. \square

Recall from Notation 2.5 that, given \mathcal{F}_t , $I^{i,t}$ denotes the immigration occurring along the skeleton descending from particle i at time t .

PROPOSITION 3.11. For all $\mu \in \mathcal{M}_c(D)$, $s, \delta > 0$ and all $f \in p(D)$ with f/ϕ bounded,

$$\lim_{\delta \mathbb{N} \ni t \rightarrow \infty} e^{-\lambda c (s+t)} \left| \sum_{i=1}^{N_t} (\langle f, I_s^{i,t} \rangle - \mathbf{P}_{\bullet, \delta \xi_i(t)} [\langle f, I_s \rangle]) \right| = 0 \quad \mathbf{P}_\mu\text{-almost surely.}$$

PROOF. By the Borel–Cantelli lemma, it is sufficient to show that for all $\varepsilon > 0$, there is a $n_0 \in \mathbb{N}$ such that

$$(3.9) \quad \sum_{n=n_0}^{\infty} \mathbf{P}_{\mu} \left(e^{-\lambda_c(s+n\delta)} \left| \sum_{i=1}^{N_{n\delta}} \langle f, I_s^{i,n\delta} \rangle - \mathbf{P}_{\bullet, \delta \xi_i(n\delta)} [\langle f, I_s \rangle] \right| > \varepsilon \right) < \infty.$$

Proposition 3.8 yields a time $T > 0$ and a constant $C_3 \in (0, \infty)$ such that for $n_0 \geq T$, the left-hand side in (3.9) is bounded by

$$\begin{aligned} & \varepsilon^{-p} \sum_{n=n_0}^{\infty} e^{-\lambda_c p(s+n\delta)} \mathbf{P}_{\mu} \left[\left| \sum_{i=1}^{N_{n\delta}} \langle f, I_s^{i,n\delta} \rangle - \mathbf{P}_{\bullet, \delta \xi_i(n\delta)} [\langle f, I_s \rangle] \right|^p \right] \\ & \leq \varepsilon^{-p} C_3 \sum_{n=n_0}^{\infty} e^{-\lambda_c(p-1)n\delta} < \infty, \end{aligned}$$

where we used Markov’s inequality in the first estimate. \square

Combining Propositions 3.10 and 3.11, we conclude that the immigration process alone has the asymptotic behavior which we expect from the superprocess.

COROLLARY 3.12. For all $\mu \in \mathcal{M}_c(D)$, $\delta > 0$ and all $f \in p(D)$ with f/ϕ bounded,

$$\lim_{\delta \mathbb{N} \ni s \rightarrow \infty} \lim_{\delta \mathbb{N} \ni t \rightarrow \infty} e^{-\lambda_c(s+t)} \sum_{i=1}^{N_t} \langle f, I_s^{i,t} \rangle = \langle f, \tilde{\phi} \rangle W_{\infty}^{\phi}(X) \quad \mathbf{P}_{\mu}\text{-almost surely.}$$

Now we are in the position to prove the SLLN along lattice times.

THEOREM 3.13. For all $\mu \in \mathcal{M}_f^{\phi}(D)$, $\delta > 0$ and all $f \in p(D)$ with f/ϕ bounded,

$$\lim_{n \rightarrow \infty} e^{-\lambda_c n \delta} \langle f, X_{n\delta} \rangle = \langle f, \tilde{\phi} \rangle W_{\infty}^{\phi}(X) \quad P_{\mu}\text{-almost surely.}$$

PROOF. By Lemma 3.2(i), it suffices to consider $\mu \in \mathcal{M}_c(D)$. Moreover, without loss of generality, we work on the skeleton space from Theorem 2.3. Corollary 3.12 yields \mathbf{P}_{μ} -almost everywhere,

$$\begin{aligned} \liminf_{\delta \mathbb{N} \ni t \rightarrow \infty} e^{-\lambda_c t} \langle f, X_t \rangle &= \liminf_{\delta \mathbb{N} \ni s \rightarrow \infty} \liminf_{\delta \mathbb{N} \ni t \rightarrow \infty} e^{-\lambda_c(s+t)} \langle f, X_{s+t} \rangle \\ &\geq \liminf_{\delta \mathbb{N} \ni s \rightarrow \infty} \liminf_{\delta \mathbb{N} \ni t \rightarrow \infty} e^{-\lambda_c(s+t)} \sum_{i=1}^{N_t} \langle f, I_s^{i,t} \rangle \\ &= \langle f, \tilde{\phi} \rangle W_{\infty}^{\phi}(X). \end{aligned}$$

Lemma 3.1(i) yields the claim. \square

3.5. *Transition from lattice to continuous times.* In this section, we extend the convergence along lattice times in Theorem 3.13 to convergence along continuous times and conclude our main results. We work under Assumptions 1, 2, 3' and 4. For $\kappa > 0$, let U^κ be the resolvent operator in integral form, that is,

$$U^\kappa f(x) := \int_0^\infty e^{-\kappa t} \mathbb{P}_x^\phi[f(\xi_t)] dt \quad \text{for all } f \in bp(D), x \in D.$$

The argument for the transition from lattice to continuous times proceeds in two steps. First we use the resolvent operator to bring the semigroup of $(\xi; (\mathbb{P}_x^\phi)_{x \in D})$ into the argument. The semigroup property gives us a martingale which, combined with Doob's L^p -inequality, enables us to control the behavior between time $n\delta$ and $(n + 1)\delta$. Second, we remove the resolvent operator by taking $\kappa \rightarrow \infty$ in $\kappa U^\kappa f(x)$. It is an analysis of hitting times for diffusion processes which allows us to control the convergences in this step. The main idea for the proof is borrowed from [46], but we employ the skeleton decomposition to replace the stochastic analysis and the martingale measures used there.

PROPOSITION 3.14. *For all $\mu \in \mathcal{M}_c(D)$, $\kappa > 0$ and $f \in bp(D)$,*

$$\lim_{t \rightarrow \infty} e^{-\lambda c t} \langle \phi \kappa U^\kappa f, X_t \rangle = \langle \phi f, \tilde{\phi} \rangle W_\infty^\phi(X) \quad \mathbf{P}_\mu\text{-almost surely.}$$

PROOF. Without loss of generality, we assume that $\mu \neq 0$ and work on the skeleton space. Since κU^κ is linear with $\kappa U^\kappa \mathbf{1} = \mathbf{1}$ the same argument that led to Lemma 3.1 shows that it suffices to prove that for all $f \in bp(D)$,

$$(3.10) \quad \liminf_{t \rightarrow \infty} e^{-\lambda c t} \langle \phi \kappa U^\kappa f, X_t \rangle \geq \langle \phi f, \tilde{\phi} \rangle W_\infty^\phi(X) \quad \mathbf{P}_\mu\text{-almost surely.}$$

Let $f, g \in bp(D)$ with $\kappa U^\kappa f \geq g$, $\delta, t > 0$, and let n be such that $n\delta \leq t < (n + 1)\delta$. Then

$$\begin{aligned} & e^{-\lambda c t} \langle \phi \kappa U^\kappa f, X_t \rangle \\ & \geq (e^{-\lambda c t} \langle \phi \kappa U^\kappa f, X_t \rangle - e^{-\lambda c t} \langle \phi \mathbb{P}^\phi[\kappa U^\kappa f(\xi_{(n+1)\delta-t}], X_t \rangle) \\ (3.11) \quad & + (e^{-\lambda c t} \langle \phi \mathbb{P}^\phi[g(\xi_{(n+1)\delta-t}], X_t \rangle - e^{-\lambda c n \delta} \langle \phi \mathbb{P}^\phi[g(\xi_\delta)], X_{n\delta} \rangle) \\ & + (e^{-\lambda c n \delta} \langle \phi \mathbb{P}^\phi[g(\xi_\delta)], X_{n\delta} \rangle - \langle \phi g, \tilde{\phi} \rangle W_\infty^\phi(X)) + \langle \phi g, \tilde{\phi} \rangle W_\infty^\phi(X) \\ & =: \Theta_{1, \kappa U^\kappa f}(n, \delta, t) + \Theta_{2, g}(n, \delta, t) + \Theta_{3, g}(n, \delta) + \langle \phi g, \tilde{\phi} \rangle W_\infty^\phi(X). \end{aligned}$$

If we show, for all $f, g \in bp(D)$, g of compact support, that \mathbf{P}_μ -almost surely,

$$(3.12) \quad \limsup_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \sup_{t \in [n\delta, (n+1)\delta]} |\Theta_{1, \kappa U^\kappa f}(n, \delta, t)| = 0,$$

$$(3.13) \quad \limsup_{n \rightarrow \infty} \sup_{t \in [n\delta, (n+1)\delta]} |\Theta_{2, g}(n, \delta, t)| = 0 \quad \text{for all } \delta > 0,$$

$$(3.14) \quad \limsup_{n \rightarrow \infty} |\Theta_{3, g}(n, \delta)| = 0 \quad \text{for all } \delta > 0,$$

then we can choose $g = \mathbb{1}_B \kappa U^\kappa f$ for $B \subset\subset D$ in (3.11) to obtain

$$\liminf_{t \rightarrow \infty} e^{-\lambda c t} \langle \phi \kappa U^\kappa f, X_t \rangle \geq \langle \phi \mathbb{1}_B \kappa U^\kappa f, \tilde{\phi} \rangle W_\infty^\phi(X) \quad \mathbf{P}_\mu\text{-almost surely.}$$

Choosing a sequence $B_j \subset\subset D$, $B_j \subseteq B_{j+1}$, $D = \bigcup_{j=1}^\infty B_j$, the factor $\langle \phi \mathbb{1}_{B_j} \kappa U^\kappa f, \tilde{\phi} \rangle$ increases, as $j \rightarrow \infty$, to

$$\begin{aligned} \langle \phi \kappa U^\kappa f, \tilde{\phi} \rangle &= \int_0^\infty \kappa e^{-\kappa t} \langle \phi \mathbb{P}^\phi[f(\xi_t)], \tilde{\phi} \rangle dt \\ &\stackrel{(2.16)}{=} \int_0^\infty \kappa e^{-\kappa t} \langle \phi f, \tilde{\phi} \rangle dt \\ &= \langle \phi f, \tilde{\phi} \rangle, \end{aligned}$$

and (3.10) follows. We begin with the proof of (3.12). Fubini’s theorem and the Markov property of $(\xi; \mathbb{P}^\phi)$ yield for all $x \in D$ and $s > 0$,

$$\begin{aligned} &|\kappa U^\kappa f(x) - \mathbb{P}_x^\phi[\kappa U^\kappa f(\xi_s)]| \\ &= \left| \int_0^\infty \kappa e^{-\kappa t} \mathbb{P}_x^\phi[f(\xi_t)] dt - \int_0^\infty \kappa e^{-\kappa t} \mathbb{P}_x^\phi[f(\xi_{t+s})] dt \right| \\ &\leq 2(1 - e^{-\kappa s}) \|f\|_\infty. \end{aligned}$$

Using the linearity of integration and the definition of $W_t^\phi(X)$, we obtain

$$\sup_{t \in [n\delta, (n+1)\delta]} \Theta_{1, \kappa U^\kappa f}(n, \delta, t) \leq 2(1 - e^{-\kappa \delta}) \|f\|_\infty \sup_{t \in [n\delta, (n+1)\delta]} W_t^\phi(X).$$

Since the martingale $(W_t^\phi(X))_{t \geq 0}$ has a finite limit, (3.12) is established. For the proof of (3.13), let $g \in bp(D)$ be compactly supported. By (2.15),

$$\Theta_{2, g}(n, \delta, t) = e^{-\lambda c (n+1)\delta} (\langle S_{(n+1)\delta-t}[\phi g], X_t \rangle - \langle S_\delta[\phi g], X_{n\delta} \rangle)$$

for all $t \in [0, (n+1)\delta]$, and the Markov property of X and (2.11) imply that $(\Theta_{2, g}(n, \delta, t) : t \in [n\delta, (n+1)\delta]; \mathbf{P}_\mu)$ is a martingale. Hence, (3.13) follows from the Borel–Cantelli lemma, Doob’s L^p -inequality (cf. Theorem II.1.7 in [53]) and $\mathbf{P}_\mu[\langle \phi g, X_{(n+1)\delta} \rangle | \mathcal{F}_{n\delta}] = \langle S_\delta[\phi g], X_{n\delta} \rangle$ when we prove that for sufficiently large n_0 ,

$$(3.15) \quad \sum_{n=n_0}^\infty e^{-\lambda c p(n+1)\delta} \mathbf{P}_\mu[|\langle \phi g, X_{(n+1)\delta} \rangle - \mathbf{P}_\mu[\langle \phi g, X_{(n+1)\delta} \rangle | \mathcal{F}_{n\delta}]|^p] < \infty.$$

By (2.10) and (3.3), we can use the skeleton decomposition to obtain, for all $s, t > 0$, \mathbf{P}_μ -almost surely,

$$\begin{aligned} &\langle \phi g, X_{s+t} \rangle - \mathbf{P}_\mu[\langle \phi g, X_{s+t} \rangle | \mathcal{F}_t] \\ (3.16) \quad &= \langle \phi g, X_{s+t}^* + I_s^{*,t} \rangle - \mathbf{P}_{X_t}[\langle \phi g, X_s^* \rangle] \\ &\quad + \sum_{i=1}^{N_t} (\langle \phi g, I_s^{i,t} \rangle - \mathbf{P}_{\bullet, \delta_{\xi_i(t)}}[\langle \phi g, I_s \rangle]). \end{aligned}$$

The monotonicity of L^p -norms and $(X_{s+t}^* + I_s^{*,t}; \mathbf{P}_\mu(\cdot|\mathcal{F}_t)) \stackrel{d}{=} (X_s^*; \mathbf{P}_{X_t})$ imply

$$(3.17) \quad \mathbf{P}_\mu[|(\phi g, X_{s+t}^* + I_s^{*,t}) - \mathbf{P}_{X_t}[(\phi g, X_s^*)]|^p] \leq \mathbf{P}_\mu[\text{Var}_{X_t}((\phi g, X_s^*))]^{p/2}.$$

Denote

$$c_1^*(x) = 2\alpha(x), \quad c_2^* = \int_{(0, \varphi_2(x)]} y^2 \Pi^*(x, dy),$$

$$c_3^*(x) = \int_{(\varphi_2(x), \infty)} y^2 \Pi^*(x, dy), \quad c^*(x) = \sum_{i=1}^3 c_i^*(x)$$

for all $x \in D$, where φ_2 is determined by Assumption 3'. We notice that $\beta^* \leq \beta$ implies that $S_t^* f \leq S_t f$ for all $f \in p(D)$. Using (2.14), (2.11), and the semigroup property of S , we obtain

$$\mathbf{P}_\mu[\text{Var}_{X_t}((\phi g, X_s^*))]$$

$$\leq \int_0^s \langle S_t S_r^* [c^*(S_{s-r}^*[\phi g])^2], \mu \rangle dr \leq \int_0^s \langle S_{t+r} [c^*(S_{s-r}[\phi g])^2], \mu \rangle dr.$$

Recall the definition of $\mathbb{P}_{\phi\mu}^\phi$ from (2.22), and use (2.15) to deduce

$$(3.18) \quad \mathbf{P}_\mu[\text{Var}_{X_t}((\phi g, X_s^*))]$$

$$\leq \langle \phi, \mu \rangle \|g\|_\infty \int_0^s e^{\lambda c(s+t)} \mathbb{P}_{\phi\mu}^\phi [c^*(\xi_{t+r}) S_{s-r}[\phi g](\xi_{t+r})] dr.$$

Writing $\bar{\beta} = \sup_{x \in D} \beta(x)$, we notice that $S_{s-r}[\phi g](x) \leq e^{\bar{\beta}s} \|\phi g\|_\infty$. Further, (2.15) implies $S_{s-r}[\phi g](x) \leq e^{\lambda c s} \|g\|_\infty \phi(x)$. Hence, for all $i \in \{1, 2, 3\}$,

$$(3.19) \quad \mathbb{P}_{\phi\mu}^\phi [c_i^*(\xi_{t+r}) S_{s-r}[\phi g](\xi_{t+r})]$$

$$\leq \min\{e^{\bar{\beta}s} \|\phi g\|_\infty \mathbb{P}_{\phi\mu}^\phi [c_i^*(\xi_{t+r})], e^{\lambda c s} \|g\|_\infty \mathbb{P}_{\phi\mu}^\phi [c_i^*(\xi_{t+r}) \phi(\xi_{t+r})]\}.$$

Since α is bounded, c_1^* is bounded. For c_2^* and c_3^* , the right-hand side of (3.19) is bounded for large t according to (2.17) and conditions (2.27) and (2.28), respectively. Combining (3.17), (3.18) and (3.19), we obtain a time $T > 0$ and a constant $C \in (0, \infty)$, which may depend on s, g and μ , such that for all $t \geq T$,

$$(3.20) \quad e^{-\lambda c p(s+t)} \mathbf{P}_\mu[|(\phi g, X_{s+t}^* + I_s^{*,t}) - \mathbf{P}_{X_t}[(\phi g, X_s^*)]|^p] \leq C e^{-\lambda c p t/2}.$$

Since $|x + y|^p \leq 2^p(|x|^p + |y|^p)$ for all $x, y \in \mathbb{R}$, (3.16), (3.20) and Proposition 3.8 yield (3.15).

It remains to prove (3.14). Since, $\langle \phi \mathbb{P}^\phi [g(\xi_\delta)], \tilde{\phi} \rangle = \langle \phi g, \tilde{\phi} \rangle$ by (2.16), (3.14) follows from Theorem 3.13. \square

In the second step we remove the resolvent operator from Proposition 3.14. The proof is essentially borrowed from the lower bound in Theorem 2.2 in [46]. We present the argument here for completeness. Recall that $\mathcal{B}_0(D) = \{B \in \mathcal{B}(D) : \ell(\partial B) = 0\}$ and $\phi|_B = \phi \mathbb{1}_B$.

PROPOSITION 3.15. For all $\mu \in \mathcal{M}_c(D)$ and $B \in \mathcal{B}_0(D)$,

$$(3.21) \quad \liminf_{t \rightarrow \infty} e^{-\lambda ct} \langle \phi|_B, X_t \rangle \geq \langle \phi|_B, \tilde{\phi} \rangle W_\infty^\phi(X) \quad P_\mu\text{-almost surely.}$$

PROOF. The claim is trivial when $\ell(B) = 0$. When (3.21) is proved for $B \in \mathcal{B}_0(D)$ with $B \subset\subset D$, then, for arbitrary $B \in \mathcal{B}_0(D)$, we choose a sequence of sets $B_k \in \mathcal{B}_0(D)$, with $B_k \subset\subset D$, $B_k \subseteq B_{k+1}$ and $B = \bigcup_{k \in \mathbb{N}} B_k$, and the monotone convergence theorem yields P_μ -almost surely,

$$\liminf_{t \rightarrow \infty} e^{-\lambda ct} \langle \phi|_B, X_t \rangle \geq \sup_{k \in \mathbb{N}} \liminf_{t \rightarrow \infty} e^{-\lambda ct} \langle \phi|_{B_k}, X_t \rangle \geq \langle \phi|_B, \tilde{\phi} \rangle W_\infty^\phi(X).$$

Hence, let $B \in \mathcal{B}_0(D)$, $B \subset\subset D$, contain a non-empty, open ball. For small $\varepsilon > 0$, let $B_\varepsilon = \{x \in B : \text{dist}(x, \partial B) \geq \varepsilon\} \neq \emptyset$ and denote by $\sigma_{B_\varepsilon} = \inf\{t > 0 : \xi_t \in B_\varepsilon\}$ the hitting time of B_ε . We write $U^\kappa(x, A) = U^\kappa \mathbb{1}_A(x)$ for all $A \in \mathcal{B}(D)$. Since $\{\xi_t \in B_\varepsilon\} \subseteq \{\sigma_{B_\varepsilon} \leq t\}$, for all $x \in D$,

$$\begin{aligned} \kappa U^\kappa(x, B_\varepsilon) &\leq \int_0^\infty \kappa e^{-\kappa t} \mathbb{P}_x^\phi(\sigma_{B_\varepsilon} \leq t) dt = \mathbb{P}_x^\phi[e^{-\kappa \sigma_{B_\varepsilon}}] \\ &\leq \mathbb{1}_B(x) + \mathbb{1}_{B^c}(x) \mathbb{P}_x^\phi[e^{-\kappa \sigma_{B_\varepsilon}}], \end{aligned}$$

where $B^c := D \setminus B$. In particular,

$$e^{-\lambda ct} \langle \phi|_B, X_t \rangle \geq e^{-\lambda ct} \langle \phi \kappa U^\kappa \mathbb{1}_{B_\varepsilon}, X_t \rangle - e^{-\lambda ct} \langle \phi|_{B^c} \mathbb{P}^\phi[e^{-\kappa \sigma_{B_\varepsilon}}], X_t \rangle,$$

and Proposition 3.14 yields, P_μ -almost surely,

$$(3.22) \quad \begin{aligned} &\liminf_{t \rightarrow \infty} e^{-\lambda ct} \langle \phi|_B, X_t \rangle \\ &\geq \langle \phi|_{B_\varepsilon}, \tilde{\phi} \rangle W_\infty^\phi(X) - \limsup_{t \rightarrow \infty} e^{-\lambda ct} \langle \phi|_{B^c} \mathbb{P}^\phi[e^{-\kappa \sigma_{B_\varepsilon}}], X_t \rangle. \end{aligned}$$

The first term on the right converges to $\langle \phi|_B, \tilde{\phi} \rangle W_\infty^\phi(X)$ as $\varepsilon \rightarrow 0$. Thus we have to show that the second term vanishes as $\varepsilon \rightarrow 0$. Heuristically, if the SLLN holds, then the lim sup is a limit with value

$$\langle \phi|_{B^c} \mathbb{P}^\phi[e^{-\kappa \sigma_{B_\varepsilon}}], \tilde{\phi} \rangle W_\infty^\phi(X).$$

Since B_ε has positive distance to B^c , this value converges to zero as $\kappa \rightarrow \infty$. Hence, we first take $\kappa \rightarrow \infty$ and then $\varepsilon \rightarrow 0$. Of course, we do not know the SLLN, yet. Thus, we artificially reintroduce the resolvent operator in order to apply Proposition 3.14.

Continuing rigorously, let $B'_\varepsilon := \{x \in B : \text{dist}(x, \partial B_\varepsilon) \leq \varepsilon/2\}$. The situation is sketched in Figure 1.

When ξ starts outside B , then $\xi_{\sigma_{B_\varepsilon}} \in \partial B_\varepsilon$, and we obtain for all $x \in B^c$,

$$(3.23) \quad \begin{aligned} \mathbb{P}_x^\phi[e^{-\kappa \sigma_{B_\varepsilon}}] &= \mathbb{P}_x^\phi \left[e^{-\kappa \sigma_{B_\varepsilon}} \frac{U^\kappa(\xi_{\sigma_{B_\varepsilon}}, B'_\varepsilon)}{U^\kappa(\xi_{\sigma_{B_\varepsilon}}, B'_\varepsilon)} \right] \\ &\leq \frac{1}{\inf_{y \in \partial B_\varepsilon} U^\kappa(y, B'_\varepsilon)} \mathbb{P}_x^\phi[e^{-\kappa \sigma_{B_\varepsilon}} U^\kappa(\xi_{\sigma_{B_\varepsilon}}, B'_\varepsilon)]. \end{aligned}$$




FIG. 1. The big ball with thick boundary is B , the small, hatched ball is B_ϵ and the shaded area denotes B'_ϵ . The diffusion is started in $x \in B^c$.

For $t \geq 0$, let $\mathcal{H}_t := \sigma(\xi_s : 0 \leq s \leq t)$. By the Markov property of ξ , the second factor on the right-hand side of (3.23) can be estimated by

$$\begin{aligned}
 & \mathbb{P}_x^\phi [e^{-\kappa\sigma_{B_\epsilon}} U^\kappa(\xi_{\sigma_{B_\epsilon}}, B'_\epsilon)] \\
 (3.24) \quad &= \mathbb{P}_x^\phi \left[e^{-\kappa\sigma_{B_\epsilon}} \mathbb{P}_x^\phi \left[\int_0^\infty e^{-\kappa t} \mathbb{1}\{\xi_{t+\sigma_{B_\epsilon}} \in B'_\epsilon\} dt \mid \mathcal{H}_{\sigma_{B_\epsilon}} \right] \right] \\
 &= \mathbb{P}_x^\phi \left[\int_{\sigma_{B_\epsilon}}^\infty e^{-\kappa t} \mathbb{1}\{\xi_t \in B'_\epsilon\} dt \right] \leq U^\kappa(x, B'_\epsilon).
 \end{aligned}$$

Writing $\Phi(\kappa, \epsilon) := \inf_{y \in \partial B_\epsilon} \kappa U^\kappa(y, B'_\epsilon)$, (3.23) and (3.24) yield

$$\mathbb{1}_{B^c}(x) \mathbb{P}_x^\phi [e^{-\kappa\sigma_{B_\epsilon}}] \leq \kappa U^\kappa(x, B'_\epsilon) / \Phi(\kappa, \epsilon),$$

and Proposition 3.14 entails, P_μ -almost surely,

$$(3.25) \quad \limsup_{t \rightarrow \infty} e^{-\lambda_c t} \langle \phi |_{B^c} \mathbb{P}^\phi [e^{-\kappa\sigma_{B_\epsilon}}, X_t] \leq \frac{1}{\Phi(\kappa, \epsilon)} \langle \phi |_{B'_\epsilon}, \tilde{\phi} \rangle W_\infty^\phi(X).$$

Clearly, $\langle \phi |_{B'_\epsilon}, \tilde{\phi} \rangle W_\infty^\phi(X)$ converges to zero as $\epsilon \rightarrow 0$. Thus, it remains to bound $\kappa U^\kappa(y, B'_\epsilon)$ for $y \in \partial B_\epsilon$, and therefore $\Phi(\kappa, \epsilon)$, away from zero. We write $b_0(x)$ for the vector whose j th component is given by $b_j(x) + \sum_{i=1}^d \partial_{x_i} a_{i,j}(x)$, $j \in \{1, \dots, d\}$, $x \in D$. Since $B \subset\subset D$,

$$\begin{aligned}
 c(B, \phi) &:= \frac{\inf_{x \in B} \phi(x)}{\sup_{x \in B} \phi(x)}, & \tilde{\beta} &:= \sup_{x \in B} |\beta(x)|, \\
 \tilde{b}_0 &:= \sup_{x \in B} |b_0(x)|, & \tilde{a} &:= \sup_{x \in B} \sup_{|v|=1} \langle v, a(x)v \rangle,
 \end{aligned}$$

satisfy $c(B, \phi), \tilde{a} \in (0, \infty)$ and $\tilde{\beta}, \tilde{b}_0 \in [0, \infty)$. For all $T > 0$, we have

$$\begin{aligned}
 \kappa U^\kappa(y, B'_\epsilon) &= \int_0^\infty \kappa e^{-\kappa t} \mathbb{P}_y^\phi(\xi_t \in B'_\epsilon) dt \\
 (3.26) \quad &= \int_0^\infty e^{-t} \mathbb{P}_y^\phi(\xi_{t/\kappa} \in B'_\epsilon) dt \\
 &\geq \int_0^T e^{-t} \mathbb{P}_y^\phi(\xi_{t/\kappa} \in B'_\epsilon) dt,
 \end{aligned}$$

and for $y \in \partial B_\varepsilon$, we use the definition of B'_ε and (2.15) to estimate

$$\begin{aligned}
 \mathbb{P}_y^\phi(\xi_{t/\kappa} \in B'_\varepsilon) &\geq \mathbb{P}_y^\phi\left(\sup_{0 \leq s \leq t/\kappa} |\xi_s - y| \leq \varepsilon/2\right) \\
 (3.27) \qquad &\geq c(B, \phi)e^{-(\lambda_c + \tilde{\beta})t/\kappa} \mathbb{P}_y\left(\sup_{0 \leq s \leq t/\kappa} |\xi_s - y| \leq \varepsilon/2\right).
 \end{aligned}$$

To estimate the probability on the right-hand side, we use Theorem 2.2.2 in [50]. Since this theorem is stated for a diffusion generator in non-divergence form, we introduced the function b_0 . In particular, for κ so large that $t\tilde{b}_0/\kappa \leq \varepsilon/4$ and for all $y \in \partial B_\varepsilon$, we deduce

$$(3.28) \qquad \mathbb{P}_y\left(\sup_{0 \leq s \leq t/\kappa} |\xi_s - y| \leq \varepsilon/2\right) \geq 1 - 2d \exp\left(-\frac{\varepsilon^2 \kappa}{32\tilde{a}td}\right).$$

Combining (3.26)–(3.28), we obtain for all $\varepsilon > 0$,

$$\liminf_{\kappa \rightarrow \infty} \Phi(\kappa, \varepsilon) \geq \int_0^T e^{-t} c(B, \phi) dt > 0.$$

Since the right-hand side does not depend on ε , taking first $\kappa \rightarrow \infty$ and then $\varepsilon \rightarrow 0$ in (3.25) and (3.22) completes the proof. \square

We are now in the position to conclude our main results.

PROOF OF THEOREM 2.9(iii). The P_μ -almost sure convergence in (2.29) for every given ℓ -almost everywhere continuous $f \in p(D)$ with f/ϕ bounded follows from Proposition 3.15 and Lemmas 3.1(i) and 3.2(i). The existence of a common set Ω_0 for all such test functions follows from Lemma 3.4. \square

PROOF OF THEOREM 1.2. The $L^1(P_\mu)$ -convergence in (1.16) was proved in Theorem 1.5, the remainder follows from Theorem 2.9(iii). \square

PROOF OF COROLLARY 1.4. The claim follows immediately from Theorem 1.2, (2.19) and (2.17). \square

4. Examples. In this section, we explore our assumptions by verifying them for many classical examples of superdiffusions from the literature. Moreover, we give several examples to illustrate the implications and boundaries of the SLLN. For all examples considered, this article proves the SLLN, and for some even the WLLN, for the first time.

4.1. *Spatially independent branching mechanisms.* In this section, we consider superdiffusions with a conservative motion and a spatially independent branching mechanism and write $\psi(z) = \psi_\beta(x, z)$ to simplify notation. Under these conditions, the total mass process $((\mathbf{1}, X_t) : t \geq 0)$ is a continuous state branching process (CSBP) with branching mechanism ψ ; cf. [3, 57]. We exclude the trivial case of a linear branching mechanism $\psi(z) = -\beta z$; see Remark 2.8 for the result in this situation. Since ψ is strictly convex, $\psi(\infty) := \lim_{z \rightarrow \infty} \psi(z)$ exists in $[-\infty, 0) \cup \{\infty\}$. Writing $z^* = \sup\{z \geq 0 : \psi(z) \leq 0\}$, we have $z^* \in (0, \infty)$ if and only if $\beta > 0$ and $\psi(\infty) = \infty$, and in this case (cf. Proposition 1.1 in [57]),

$$P_\mu[e^{-z^*(\mathbf{1}, X_t)}] = e^{-z^*(\mathbf{1}, \mu)} \quad \text{for all } \mu \in \mathcal{M}_f(D), t \geq 0.$$

In particular, Assumption 1 is satisfied with $w(x) = z^*$ for all $x \in D$. In this CSBP context, the skeleton decomposition was proved by Berestycki et al. in [3] a few years before [43]. The martingale function z^* is related to the event of weak extinction $\mathcal{E}_{\text{lim}} = \{\lim_{t \rightarrow \infty} (\mathbf{1}, X_t) = 0\}$ by the identity $P_{\delta_x}(\mathcal{E}_{\text{lim}}) = e^{-z^*}$ which holds even if $\beta \leq 0$ or $\psi(\infty) < 0$. To compare the martingale function $w(x) = -\log P_{\delta_x}(\mathcal{E}_{\text{lim}})$ to the classical choice $w(x) = -\log P_{\delta_x}(\mathcal{E}_{\text{fin}})$, where \mathcal{E}_{fin} denotes the event of extinction after finite time, notice that $\mathcal{E}_{\text{fin}} \subseteq \mathcal{E}_{\text{lim}}$, and for all $\mu \in \mathcal{M}_f(D)$,

$$(4.1) \quad \begin{aligned} P_\mu(\mathcal{E}_{\text{fin}}) &= P_\mu(\mathcal{E}_{\text{lim}}) = e^{-z^*(\mathbf{1}, \mu)} > 0 \\ &\text{if } \psi(\infty) = \infty \text{ and } \int^\infty \frac{1}{\psi(z)} dz < \infty. \end{aligned}$$

Otherwise, $P_\mu(\mathcal{E}_{\text{fin}}) = 0$ and on \mathcal{E}_{lim} the total mass of X drifts to zero while staying positive at all finite times; cf. [33, 57].

From now on, assume $\beta > 0$, $\psi(\infty) = \infty$ and $w(x) = z^*$. In this case, Assumption 3 simplifies to

$$\phi \text{ bounded} \quad \text{and} \quad \int_{(1, \infty)} y^p \Pi(dy) < \infty \quad \text{for some } p \in (1, 2).$$

In the following, we present two families of superprocesses for which the SLLN is proved by Theorem 1.2. As far as we know, these results are new. Apart from the intrinsic interest, the results are very useful since the analyzed processes are frequently employed to obtain further examples of superprocesses with interesting properties via h -transform. For those examples the SLLN follows from Lemma 2.10.

We begin with the inward Ornstein–Uhlenbeck process (OU-process) which has attracted a wide interest in the literature. Specifically, its asymptotic behavior is the subject of recent research articles [47, 52].

EXAMPLE 4.1 (Inward OU-process). Let $d \geq 1$, $D = \mathbb{R}^d$, $L = \frac{1}{2} \Delta - \gamma x \cdot \nabla$ with $\gamma > 0$, ψ spatially independent with $\beta \in (0, \infty)$, $\psi(\infty) = \infty$ and

$\int_{(1,\infty)} y^p \Pi(dy) < \infty$ for some $p \in (1, 2]$. Then Theorem 1.2 applies with $\phi = \mathbf{1}$, $\tilde{\phi}(x) = (\gamma/\pi)^{d/2} e^{-\gamma\|x\|^2}$ and $\lambda_c = \beta$.

The generator L corresponds to the positive recurrent inward OU-process with transition density

$$(4.2) \quad p_{\text{in-OU}}(x, y, t) = \left(\frac{\gamma}{\pi(1 - e^{-2\gamma t})} \right)^{d/2} \exp\left(-\frac{\gamma}{1 - e^{-2\gamma t}} \|y - e^{-\gamma t} x\|^2 \right)$$

for all $x, y \in \mathbb{R}^d, t > 0$. Hence, $\lambda_c = \lambda_c(L + \beta) = \beta > 0$, L is product L^1 -critical, $\phi = \mathbf{1}$ and $\tilde{\phi}(x) = (\gamma/\pi)^{d/2} e^{-\gamma\|x\|^2}$; cf. Chapter 4 in [50] or Example 3 in [51]. Thus Assumptions 1–3 are satisfied. Using the estimate for $p^\phi = p_{\text{in-OU}}$ in (2.18), we obtain that condition (2.34) holds for $a(t) = \sqrt{(\lambda_c/\gamma + \delta)t}$ with $\delta > 0$ (cf. Example 10 in [17]) and using (4.2), we deduce that (2.35) holds with $K = 1$. Hence, Theorem 2.13 applies and Assumption 4 is satisfied.

EXAMPLE 4.2 (Outward OU-process). Let $d \geq 1, D = \mathbb{R}^d, L = \frac{1}{2}\Delta + \gamma x \cdot \nabla$ with $\gamma > 0, \psi$ spatially independent with $\beta \in (\gamma d, \infty), \psi(\infty) = \infty$ and $\int_{(1,\infty)} y^p \Pi(dy) < \infty$ for some $p \in (1, 2]$. Then Theorem 1.2 applies with $\phi(x) = (\gamma/\pi)^{d/2} e^{-\gamma\|x\|^2}, \tilde{\phi} = \mathbf{1}$ and $\lambda_c = \beta - \gamma d$.

The generator L corresponds to the conservative, transient outward OU-process. The operator $L_1 := L + \gamma d$ is the formal adjoint of the inward OU-process with parameter γ . Hence, L_1 is critical with ground states $\phi_1(x) = (\gamma/\pi)^{d/2} e^{-\gamma\|x\|^2}$ and $\tilde{\phi}_1 = \mathbf{1}$ by Example 4.1; see Theorem 4.3.3 in [50] or Example 2 in [51]. Writing $L_1 = L + \beta - (\beta - \gamma d)$, we deduce that Assumptions 1–3 hold and $\phi, \tilde{\phi}$, and λ_c have been correctly identified. The corresponding ergodic motion is the inward OU-process with parameter γ . Thus conditions (2.34) and (2.35) can be verified using (4.2), (2.18), $a(t) = e^{\gamma(1+\delta)t}$ for some $\delta > 0$ and $K > 1 + \delta$, and Theorem 2.13 implies that Assumption 4 holds.

The SLLN describes the asymptotic behavior of the mass in compact sets. In general one cannot draw conclusions for the scaling of the total mass from the local behavior; cf. [18, 20]. Example 4.2 illustrates this fact. Since the total mass process is a CSBP with branching mechanism $\psi, Y_t = e^{-\beta t} \langle \mathbf{1}, X_t \rangle$ converges to a finite random variable Y_∞ with $P_\mu(Y_\infty = 0) = P_\mu(\mathcal{E}_{\text{lim}})$ if $\beta > 0$ and $\int_{(1,\infty)} y \log y \Pi(dy) < \infty$; cf. [33]. In particular, in Example 4.2, the local growth rate $\lambda_c = \beta - \gamma d$ is strictly smaller than the global growth rate β . The reason is the transient nature of the underlying diffusion which allows mass to leave compact sets permanently and is reflected in the decay of ϕ at infinity. In particular, the function $\mathbf{1}$ is not an allowed test function in Theorem 1.2, but the focus is on functions of the form $\mathbb{1}_B$ for B a compact set.

We call the diffusion corresponding to the generator $L = \frac{1}{2}\nabla \cdot a\nabla + b \cdot \nabla$ symmetric if $b = a\nabla Q$ for some $Q \in C^{2,\eta}(D)$. The inward and outward OU-processes

constitute examples of symmetric diffusions with $Q(x) = -\frac{\gamma}{2}\|x\|^2$ and $Q(x) = \frac{\gamma}{2}\|x\|^2$, respectively. Chen et al. [7] studied superdiffusions with a symmetric motion but insisted that Q is bounded. Hence, their results are not applicable to Examples 4.1 and 4.2. The result from Liu et al. [46] is not applicable since the domain is not of finite Lebesgue measure.

Engländer and Winter [22] proved convergence in probability in (1.16) for the situation of a quadratic branching mechanism. It is straightforward to extend their argument to general branching mechanisms but the method requires second moments. Hence, if $\int_{(1,\infty)} y^p \Pi(dy) < \infty$ for some $p \in (1, 2)$ but not for $p = 2$, then even the convergence in probability in Examples 4.1 and 4.2 is new.

4.2. Quadratic branching mechanisms. In this section, we consider the classical situation of a quadratic branching mechanism studied by Engländer, Pinsky and Winter [19, 22] and Chen, Ren and Wang [7]. Our assumptions on the branching mechanism in this section are $\alpha, \beta \in C^\eta(D)$, $\alpha(x) > 0$ for all $x \in D$, $\lambda_c := \lambda_c(L + \beta) < \infty$ and $\Pi \equiv 0$. We write $\psi(x, z) = -\beta(x)z + \alpha(x)z^2$ and call ψ a *generalised quadratic branching mechanism* (GQBM). In Section 1.1 we insisted that α and β are bounded. This assumption can be relaxed as follows. First suppose that β is bounded from above but not necessarily from below. Engländer and Pinsky [19] showed that there is a unique $\mathcal{M}_f(D)$ -valued Markov process $X = (X_t)_{t \geq 0}$ such that for all continuous $f \in bp(D)$ and all $\mu \in \mathcal{M}_f(D)$,

$$P_\mu[e^{-\langle f, X_t \rangle}] = e^{-\langle u_f(\cdot, t), \mu \rangle},$$

where u_f is the minimal, nonnegative solution $u \in C(D \times [0, \infty))$, $(x, t) \mapsto u(x, t)$ twice continuously differentiable in $x \in D$ and once in $t \in (0, \infty)$, to

$$\begin{aligned} \partial_t u(x, t) &= Lu(x, t) - \psi(x, u(x, t)) && \text{for all } (x, t) \in D \times (0, \infty), \\ (4.3) \quad u(x, 0) &= f(x) && \text{for all } x \in D. \end{aligned}$$

If α and β are bounded, the minimal solution of (4.3) equals the unique solution to (1.3) by Lemma A1 in [19]. Hence, the two definitions are consistent.

Now let $\beta \in C^\eta(D)$ with $\lambda_c = \lambda_c(L + \beta) < \infty$ be not necessarily bounded from above. By definition (1.9), there exists $\lambda \in \mathbb{R}$ and $h \in C^{2,\eta}(D)$, $h > 0$, such that $(L + \beta)h = \lambda h$. Recall the definition of h -transforms from Section 2.1.3. An $(L, \psi; D)$ -superprocess can be defined by $X = \frac{1}{h}X^h$, where X^h is the $(L_0^h, \psi^h; D)$ superprocess with $\beta^h = \lambda$ and $\alpha^h = \alpha h$; cf. [19]. Since h is not necessarily bounded from below, the process X may take values in the space of σ -finite measures $\mathcal{M}(D)$. While we have considered mainly finite measure-valued processes in this article, it is natural to consider also processes with values in the space $\mathcal{M}(D)$ via the branching property, and, as noted in Remark 3.3, in our results the space of starting measures $\mathcal{M}_f^\phi(D)$ can be enlarged to the space of all $\mu \in \mathcal{M}(D)$ with $\langle \phi, \mu \rangle < \infty$.

Engländer and Pinsky [19] proved the skeleton decomposition for supercritical superdiffusions with GQBM long before [43]. We only record the existence of a martingale function in the following lemma. Recall that \mathcal{E}_{fin} denotes the event of extinction after a finite time.

LEMMA 4.3 (Engländer and Pinsky [19]). *Let ψ be a GQBM and $\lambda_c > 0$. The function $x \mapsto w(x) := -\log P_{\delta_x}(\mathcal{E}_{\text{fin}})$ is strictly positive, belongs to $C^{2,\eta}(D)$ and satisfies (1.6).*

PROOF. By Theorem 3.1 and Corollary 4.2 in [19], $w \in C^{2,\eta}(D)$, $w(x) > 0$ for all $x \in D$ and

$$(4.4) \quad P_\mu(\mathcal{E}_{\text{fin}}) = e^{-\langle w, \mu \rangle} \quad \text{for all } \mu \in \mathcal{M}_c(D).$$

Recall the notation from the beginning of Section 2.1.1. Let $B \subset\subset D$ be a domain and $\mu \in \mathcal{M}_f(D)$ with $\text{supp}(\mu) \subseteq B$. Then the support of the exit measure \tilde{X}_t^B is P_μ -almost surely compact; see the discussion following (2.3). Since \mathcal{E}_{fin} is a tail event, the Markov property and (4.4) yield $P_\mu[e^{-\langle \tilde{w}, \tilde{X}_t^B \rangle}] = e^{-\langle w, \mu \rangle}$. Choose a sequence of functions $w_j \in C_c^+(D)$ with $w_j \uparrow w$ pointwise and a sequence of domains $B_k \subset\subset D$, $B_k \subseteq B_{k+1}$, $D = \bigcup_{k=1}^\infty B_k$. By Lemma A1 in [19] and Lemma A.6 in the present article, $\tilde{u}_{w_j}^{B_k}$ is increasing in j and k with $\lim_{k \rightarrow \infty} \tilde{u}_{w_j}^{B_k} = u_{w_j}$ pointwise, and we obtain for all $\mu \in \mathcal{M}_c(D)$,

$$\begin{aligned} P_\mu[e^{-\langle w, X_t \rangle}] &= \lim_{j \rightarrow \infty} \lim_{k \rightarrow \infty} P_\mu[e^{-\langle \tilde{w}_j, \tilde{X}_t^{B_k} \rangle}] = \lim_{k \rightarrow \infty} \lim_{j \rightarrow \infty} P_\mu[e^{-\langle \tilde{w}_j, \tilde{X}_t^{B_k} \rangle}] \\ &= \lim_{k \rightarrow \infty} P_\mu[e^{-\langle \tilde{w}, \tilde{X}_t^{B_k} \rangle}] = e^{-\langle w, \mu \rangle}. \end{aligned} \quad \square$$

In the remainder of this section, we choose w to be the function $w(x) = -\log P_{\delta_x}(\mathcal{E}_{\text{fin}})$ and let $Z = (Z_t)_{t \geq 0}$ be a strictly dyadic branching particle diffusion, where the spatial motion is defined by (1.7) and the branching rate is given by $q = \alpha w$ [in accordance with (1.8)].

One advantage of allowing unbounded α and β is that the setup is now invariant under h -transforms: for any $h \in C^{2,\eta}(D)$, $h > 0$, ψ^h is a GQBM. Moreover,

$$\begin{aligned} w^h(x) &:= -\log P_{\delta_x}^h(\exists t \geq 0 : \langle \mathbf{1}, X_t^h \rangle = 0) \\ (4.5) \quad &= -\log P_{h(x)^{-1}\delta_x}(\exists t \geq 0 : \langle h, X_t \rangle = 0) \\ &= w(x)/h(x), \end{aligned}$$

and Lemmas 2.10 and 2.11 remain valid for GQBM and $\mathbb{H}(\psi) = \{h \in C^{2,\eta}(D) : h > 0\}$. We record the following result.

THEOREM 4.4. *Let ψ be a GQBM and suppose Assumption 2 holds and $\phi\alpha$ is bounded. Let $\mu \in \mathcal{M}_f^\phi(D)$.*

(i) For all $f \in p(D)$ with f/ϕ bounded, the convergence in (1.16) holds in $L^1(P_\mu)$.

(ii) If, in addition, Assumption 4 holds, then there exists a measurable set Ω_0 with $P_\mu(\Omega_0) = 1$, and on Ω_0 , the convergence in (1.16) holds for all ℓ -almost everywhere continuous $f \in p(D)$ with f/ϕ bounded.

PROOF. Let X^ϕ be an $(L_0^\phi, \psi^\phi; D)$ -superprocess. Since $\beta^\phi = \lambda_c$ and $\alpha^\phi = \phi\alpha$ are bounded, X^ϕ satisfies the assumptions of Section 1.1. Moreover, X^ϕ satisfies Assumption 1 by Lemma 4.3, Assumption 2 with $\phi^\phi = \mathbf{1}$ by Lemma 2.10(i) and Assumption 3''. Hence, Theorem 2.12(i) and Lemma 2.10(ii) yield the first part of the claim. Lemmas 2.11, 2.10(ii), 3.4 and Theorem 2.12(ii) yield the second part. \square

The h -transforms are one way to relate two superprocesses to each other, another is monotonicity.

LEMMA 4.5. Let ψ_β and $\hat{\psi}_{\hat{\beta}}$ be two branching mechanisms as defined in Section 1.1 with $\psi_\beta \geq \hat{\psi}_{\hat{\beta}}$. Let X and \hat{X} be $(L, \psi_\beta; D)$ - and $(L, \hat{\psi}_{\hat{\beta}}; D)$ -superprocesses, respectively.

- (i) For all $\mu \in \mathcal{M}_f(D)$, $f \in bp(D)$, $t \geq 0$, $P_\mu[e^{-\langle f, X_t \rangle}] \geq P_\mu[e^{-\langle f, \hat{X}_t \rangle}]$.
- (ii) Let $w(x) = -\log P_{\delta_x}(\exists t \geq 0: \langle \mathbf{1}, X_t \rangle = 0)$ and $\hat{w}(x) = -\log P_{\delta_x}(\exists t \geq 0: \langle \mathbf{1}, \hat{X}_t \rangle = 0)$ for all $x \in D$. Then $w \leq \hat{w}$.

PROOF. Part (i) is proved in Appendix A.2. Part (ii) follows from part (i) and the identity $w(x) = \lim_{t \rightarrow \infty} \lim_{\theta \rightarrow \infty} -\log P_{\delta_x}[e^{-\theta \langle \mathbf{1}, X_t \rangle}]$. \square

We saw in Example 4.2 that the SLLN describes the asymptotics of the mass in compact sets, not necessarily the global growth. A second distinction between the local and global behavior can be observed on the event $\{W_\infty^\phi(X) = 0\} \setminus \mathcal{E}_{\text{fin}}$. Engländer and Turaev [21], Problem 14, raised the question whether this event can have positive probability. Suppose Assumption 2 holds. Engländer [14] observed that if $\lim_{t \rightarrow \infty} e^{-\lambda_c t} \langle f, X_t \rangle = \langle f, \tilde{\phi} \rangle W_\infty^\phi(X)$ in distribution for all $f \in C_c^+(D)$, $\mu \in \mathcal{M}_c(D)$, and if the support of X is transient, then

$$(4.6) \quad P_\mu(W_\infty^\phi(X) = 0) > P_\mu(\mathcal{E}_{\text{fin}}) \quad \text{for all } \mu \in \mathcal{M}_c(D), \mu \neq 0.$$

Here the support of $(X; P_\mu)$ is recurrent if

$$P_\mu(X_t(B) > 0 \text{ for some } t \geq 0 | \mathcal{E}_{\text{fin}}^c) = 1 \quad \text{for every open } B \subseteq D, B \neq \emptyset,$$

and transient otherwise. See [19] for a detailed discussion of recurrence and transience of the support of superdiffusions.

We study three examples in this section. In the first example, α and β are bounded, but w is unbounded. In the second example α is bounded, but β, ϕ

and w are unbounded. Both examples are based on a recurrent motion but while the support of the superprocess is recurrent in the second, it is transient in the first example. The third example considers a large class of processes containing super-Brownian motion with compactly supported growth rate β and instances of non-symmetric underlying motions.

The domain for all these examples is $D = \mathbb{R}^d$, and therefore, none of them is covered in Liu et al.'s [46] article. Chen et al.'s [7] article is not applicable to the first two examples since they are based on the inward-OU process as underlying motion (because, as in Section 4.1, Q is unbounded) and not to the third because the motion is non-symmetric (for some processes in the considered class), and the variance parameter α is unbounded, whereas [7] requires α to be bounded.

The motivation for the first example comes from Example 5.1 in [19].

EXAMPLE 4.6. Let $d \geq 1$, $D = \mathbb{R}^d$, $L = \frac{1}{2}\Delta - \gamma x \cdot \nabla$ with $\gamma > 0$, $\beta \in (0, \infty)$ constant, $\alpha(x) = e^{-\gamma\|x\|^2}$, $\Pi \equiv 0$. Then Theorem 1.2 applies with $\phi = \mathbf{1}$, $\tilde{\phi}(x) = (\gamma/\pi)^{d/2}e^{-\gamma\|x\|^2}$ and $\lambda_c = \beta$. Moreover, $w(x) = (\beta + \gamma d)e^{\gamma\|x\|^2}$, the support of X is transient and (4.6) holds.

There are two ways to prove (1.16) for this example. First, we perform an h -transform with $h(x) = (\gamma/\pi)^{-d/2}e^{\gamma\|x\|^2}$ to obtain

$$L_0^h = \frac{1}{2}\Delta + \gamma x \cdot \nabla, \quad \beta^h = \beta + \gamma d, \quad \alpha^h = (\pi/\gamma)^{d/2}.$$

In Example 4.2, we showed that Theorem 1.2 applies to the $(L_0^h, \psi^h; \mathbb{R}^d)$ -superprocess. The $(L, \psi; \mathbb{R}^d)$ -superprocess can be recovered by an h -transform with $h_2 = 1/h$, and Lemma 2.10 yields that Assumption 2 is satisfied with the stated ϕ , $\tilde{\phi}$ and λ_c , and that (1.16) holds. Alternatively, we can deduce (1.16) by a direct application of Theorems 1.2 and 2.13. Assumption 1 holds by Lemma 4.3, and Assumption 3 holds since α and ϕ are bounded. To verify Assumption 4, we notice that $w^h(x) = \beta^h/\alpha^h = (\beta + \gamma d)(\gamma/\pi)^{d/2}$ by (4.1), and (4.5) yields

$$w(x) = w^h(x)h(x) = (\beta + \gamma d)e^{\gamma\|x\|^2} \quad \text{for all } x \in \mathbb{R}^d.$$

Thus w is not bounded from above. The verification of (2.34) and (2.35) is the same as in Example 4.2 since w/ϕ is of the same order and the ergodic motion is the same. Hence the conditions hold with $a(t) = e^{\gamma(1+\delta)t}$ for some $\delta > 0$ and $K > 1 + \delta$.

To see that the support of X is transient, notice that the support is invariant under h -transforms and the support of the $(L_0^h; \psi^h; \mathbb{R}^d)$ -superprocess is transient by Theorem 4.6 in [50] and Example 2 in [51].

Example 4.6 should be compared to Example 4.2 for a quadratic branching mechanism. In both examples, the support of the superprocess is transient and the event $\{W_\infty^\phi(X) = 0\} \setminus \mathcal{E}_{\text{fin}}$ has positive probability. Hence, in both examples,

mass can escape to infinity which is reflected in the SLLN by virtue of the fact that $W_\infty^\phi(X) = 0$. However, the motion in Example 4.6 is recurrent and the SLLN captures not only the local but also the global growth of mass.

The unbounded w in Example 4.6 can be interpreted as follows. Heuristically, since the local growth rate β is bounded away from zero, on average a large population is generated everywhere in space. Risk for the branching process comes from areas of a relatively large variance for the total mass process. In contrast, when the variance parameter α is very small, then extinction is unlikely and w becomes large.

The motivation for the next example comes from Example 10 in [17]. For $B \in \mathcal{B}(D)$, $f_1, f_2 \in p(B)$, we write $f_1 \asymp f_2$ if there are constants $0 < c \leq C < \infty$ such that $cf_1(x) \leq f_2(x) \leq Cf_1(x)$ for all $x \in B$.

EXAMPLE 4.7. Let $d \geq 1$, $D = \mathbb{R}^d$, $L = \frac{1}{2}\Delta - \gamma x \cdot \nabla$, $\beta(x) = c_1 \|x\|^2 + c_2$, where $\gamma, c_1, c_2 > 0$, $\gamma > \sqrt{2c_1}$, and write $\vartheta := \frac{1}{2}(\gamma - \sqrt{\gamma^2 - 2c_1})$. Then Assumption 2 holds with $\lambda_c = c_2 + d\vartheta$, $\phi(x) = e^{\vartheta \|x\|^2}$ and $\tilde{\phi}(x) = ce^{(\vartheta - \gamma)\|x\|^2}$, where $c = (\frac{\gamma - 2\vartheta}{\pi})^{d/2}$. Suppose that $\Pi \equiv 0$ and $\alpha \in C^\eta(D)$ with $\alpha \asymp 1/\phi$ on \mathbb{R}^d . Then Theorem 4.4 applies, $w \asymp \phi$ and the support of X is recurrent.

Let $h(x) = e^{\vartheta \|x\|^2}$. Using $-\gamma + 2\vartheta = -\sqrt{\gamma^2 - 2c_1}$ and $\vartheta^2 - \gamma\vartheta + \frac{c_1}{2} = 0$, we observe that on \mathbb{R}^d ,

$$L_0^h = \frac{1}{2}\Delta - \sqrt{\gamma^2 - 2c_1}x \cdot \nabla, \quad \beta^h = \vartheta d + c_2, \quad \alpha^h(x) = h(x)\alpha(x) \asymp 1.$$

The $(L_0^h, \psi^h; \mathbb{R}^d)$ -superprocess, denoted by X^h , satisfies Assumption 2 by Example 4.1 with $\phi^h = \mathbf{1}$, $\tilde{\phi}^h \asymp e^{(2\vartheta - \gamma)\|x\|^2}$ and $\lambda_c^h = \vartheta d + c_2$. Hence, Lemma 2.10(i) shows that X satisfies Assumption 2, $\phi, \tilde{\phi}$ and λ_c have been correctly identified and $\phi\alpha$ is bounded. When we have verified Assumption 4 for X^h , then Lemma 2.11 will yield Assumption 4 for X , and the claim is established.

To this end, choose constants $c_3, c_4 > 0$ such that $c_3/h \leq \alpha \leq c_4/h$. Let $\overline{\psi}(x, z) := -\beta^h z + c_3 z^2$ and $\underline{\psi}(x, z) := -\beta^h z + c_4 z^2$, and denote by \overline{w} and \underline{w} the martingale functions corresponding to the event of extinction after finite time for the $(L_0^h, \overline{\psi}; \mathbb{R}^d)$ and $(L_0^h, \underline{\psi}; \mathbb{R}^d)$ -superprocesses, respectively. Since $\overline{\psi} \leq \psi^h \leq \underline{\psi}$, Lemma 4.5(ii) and (4.1) imply

$$\beta^h/c_3 = \overline{w}(x) \geq w^h(x) \geq \underline{w}(x) = \beta^h/c_4 \quad \text{for all } x \in D.$$

Hence, $w^h \asymp \mathbf{1} = \phi^h$. Now the verification of (2.34) and (2.35) for X^h is the same as in Example 4.1, and we can choose $a(t) = \sqrt{(\lambda_c/\gamma + \delta)t}$ for some $\delta > 0$ and $K = 1$. Theorem 2.13 verifies Assumption 4.

The support of X is recurrent since the support is invariant under h -transforms and the support of the $(L_0^h, \psi^h; \mathbb{R}^d)$ -superprocess is recurrent according to Theorem 4.4(b) in [19].

The next example covers a large class of processes. The underlying motion is a Brownian motion with or without a compactly supported drift term. Depending on the choice of that drift, the motion can be symmetric or non-symmetric. For a choice of b which makes L non-symmetric, see Example 13 in [17]. The article by Chen et al. [7] excludes non-symmetric motions. The example is motivated by Example 22 in [21] and Examples 12 and 13 in [17].

EXAMPLE 4.8. Let $d \in \{1, 2\}$, $D = \mathbb{R}^d$, $L = \frac{1}{2}\Delta + b \cdot \nabla$ where all components of b belong to $C^{1,\eta}(\mathbb{R}^d)$ for some $\eta \in (0, 1]$ and are of compact support, $\beta_0 \in C^\eta(\mathbb{R}^d)$ nonnegative and of compact support, $\beta_0 \neq \mathbf{0}$. There exists $\theta > 0$ such that $\lambda_c(L + \theta\beta_0) > 0$, and we let $\beta = \theta\beta_0$, $\lambda_c = \lambda_c(L + \beta)$. We further write

$$\varrho(x) = \|x\|^{(1-d)/2} e^{-\sqrt{2\lambda_c}\|x\|} \quad \text{for all } x \in \mathbb{R}^d \setminus \{0\}.$$

Let $\alpha \in C^\eta(D)$, $\alpha(x) > 0$ for all $x \in \mathbb{R}^d$ and $\alpha \asymp 1/\varrho$ on $\mathbb{R}^d \setminus B$ for an open ball B around the origin. Then Theorem 4.4 applies with $\phi, \tilde{\phi}, w \asymp \varrho$ on $\mathbb{R}^d \setminus B$, and the support of X is recurrent.

The existence of θ is proved in Theorems 4.6.3 and 4.6.4 of Pinsky’s book [50] and $L + \beta - \lambda_c$ is critical by Theorem 4.6.7 in the same book. Denote by G the Green’s function corresponding to the operator $L - \lambda_c$. Then $\phi \asymp G(\cdot, 0)$ on $\mathbb{R}^d \setminus B$ by Theorems 4.6.3 and 7.3.8 in [50]. Pinsky showed in Example 7.3.11 that the Green’s function G_1 of $\frac{1}{2}\Delta - \lambda_c$ satisfies $G_1(\cdot, 0) \asymp \varrho$ on $\mathbb{R}^d \setminus B$. Since b is compactly supported, $G_1(\cdot, 0) \asymp G(\cdot, 0)$ on $\mathbb{R}^d \setminus B$, and the estimate for ϕ is established. The same argument yields the same estimate for $\tilde{\phi}$, and Assumption 2 holds. Moreover, $\phi\alpha$ is bounded.

To check Assumption 4 we use Theorem 2.13. An h -transform of the $(L, \psi; \mathbb{R}^d)$ -superprocess with $h = \phi$ gives an $(L_0^\phi, \psi^\phi; \mathbb{R}^d)$ -superprocess, where L_0^ϕ corresponds to a conservative, positive recurrent motion and $\psi^\phi(x, z) = -\lambda_c z + \phi(x)\alpha(x)z^2$. Since $\phi\alpha \asymp \mathbf{1}$, $w^\phi \asymp \mathbf{1}$ by the same argument as in Example 4.7. Hence (4.5) implies $w/\phi \asymp \mathbf{1}$ and conditions (i) and (ii) of Theorem 2.13 have been verified in Examples 12 and 13 of [17] with $a(t) = \sqrt{2\|\beta\|_\infty}t$, $K > 1/\sqrt{2\lambda_c}$. Since w is bounded and the underlying diffusion is recurrent, Theorem 4.4 in [19] shows that the support of X is recurrent.

4.3. *Bounded domains.* In the situation that D is a bounded Lipschitz domain, and L is a uniformly elliptic operator with smooth coefficients, Liu et al. [46] prove the SLLN for a general branching mechanism with β any bounded, measurable function on D and α and Π as in Section 1.1.

However, the Wright–Fisher diffusion on domain $D = (0, 1)$ is a diffusion process whose diffusion matrix $a(x) = x(1 - x)$ is not uniformly elliptic. The process has attracted a wide interest in the literature; see, for example, [4, 30, 32]. Fleischmann and Swart [30] studied the large-time behavior of the corresponding superprocess with spatially independent, quadratic branching mechanism

on $[0, 1]$. They conjecture a SLLN for the process restricted to $D = (0, 1)$ (see above (23) in [30]) but prove only convergence in L^2 . The Wright–Fisher diffusion is not conservative, so the arguments in Section 4.1 are not applicable. However, Theorem 1.2 applies, and the following theorem proves the conjecture for all Lebesgue-almost everywhere continuous test functions $f \in p(D)$ with f/ϕ bounded. (Fleischmann and Swart do not assume any continuity.)

THEOREM 4.9 (Super-Wright–Fisher diffusion). *Let $D = (0, 1)$, $\beta \in (1, \infty)$, $\alpha > 0$, $\Pi \equiv 0$ and*

$$L = \frac{1}{2}x(1-x)\frac{d^2}{dx^2} = \frac{1}{2}\frac{d}{dx}x(1-x)\frac{d}{dx} + \frac{2x-1}{2}\frac{d}{dx}.$$

Then Theorem 1.2 applies with $\phi(x) = 6x(1-x)$, $\tilde{\phi} = \mathbf{1}$ and $\lambda_c = \beta - 1$.

PROOF. Let $h(x) = 6x(1-x)$. Fleischmann and Swart proved in Lemma 20 of [30] that the generator

$$L_0^h = \frac{1}{2}\frac{d}{dx}x(1-x)\frac{d}{dx} + \frac{1-2x}{2}\frac{d}{dx}$$

corresponds to an ergodic diffusion with invariant law $h(x)\ell(dx)$ on D . Using $\beta^h = \beta - 1$, we deduce that $\lambda_c(L_0^h + \beta^h) = \beta - 1$, $\phi^h = \mathbf{1}$, $\check{\phi}^h = h$ and, using Lemma 2.10, Assumption 2 for the $(L, \psi; D)$ superprocess as well as the stated identities for ϕ , $\check{\phi}$ and λ_c are established. Assumption 1 holds by Lemma 4.3; the boundedness of α and ϕ implies that Assumption 3 is satisfied. To verify Assumption 4, we notice that condition (i) of Theorem 2.13 is trivially satisfied for $D_t = D$, and (2.33) for $D_t = D$ and $K = 1$ has been proved in Lemma 20 of [30]. Hence Assumption 4 follows from Theorems 2.13 and Theorem 1.2 applies. \square

APPENDIX

A.1. Feynman–Kac arguments. In this section, we prove an integral identity that is used several times in this article. Versions of this result appeared in Lemma A.I.1.5 of [11] and Lemma 4.1.2 of [12], but the format and assumptions are different. Like in the remainder of the article, $(\xi = (\xi_t)_{t \geq 0} : (\mathbb{P}_x)_{x \in D})$ is a diffusion as described in Section 1.1.

LEMMA A.1. *Let $T > 0$ and either $B = D$ or $B \subset\subset D$ open. Write $\tau = \inf\{t \geq 0 : \xi_t \notin B\}$ and $A = D$ if $B = D$; $A = \overline{B}$ if $B \subset\subset D$.*

(i) *Let $f_1 \in b(A)$, $g_1 : A \times [0, T] \rightarrow \mathbb{R}$ measurable and bounded from above and $f_2, g_2 \in b(A \times [0, T])$. If for all $(x, t) \in A \times [0, T]$,*

$$(A.1) \quad \begin{aligned} v(x, t) = & \mathbb{P}_x \left[e^{\int_0^{t \wedge \tau} (g_1 + g_2)(\xi_r, t-r) dr} f_1(\xi_{t \wedge \tau}) \right] \\ & + \mathbb{P}_x \left[\int_0^{t \wedge \tau} e^{\int_0^s (g_1 + g_2)(\xi_r, t-r) dr} f_2(\xi_s, t-s) ds \right], \end{aligned}$$

then, for all $(x, t) \in A \times [0, T]$,

$$v(x, t) = \mathbb{P}_x \left[e^{\int_0^{t \wedge \tau} g_1(\xi_r, t-r) dr} f_1(\xi_{t \wedge \tau}) \right] + \mathbb{P}_x \left[\int_0^{t \wedge \tau} e^{\int_0^s g_1(\xi_r, t-r) dr} (f_2(\xi_s, t-s) + g_2(\xi_s, t-s)v(\xi_s, t-s)) ds \right].$$

(ii) The statement of (i) remains valid when $f_1 \in bp(A)$, $f_2, g_1, g_2 : A \times [0, T] \rightarrow \mathbb{R}$ measurable with g_1 bounded from above, g_2 nonnegative, f_2 non-positive and $g_1 + g_2$ bounded from above. Notice that in this case, v might attain the value $-\infty$.

PROOF. For all $t \geq 0$, write

$$\mathcal{Y}_t = e^{\int_0^{t \wedge \tau} (g_1+g_2)(\xi_r, t-r) dr} f_1(\xi_{t \wedge \tau})$$

and

$$\mathcal{Z}_t = \int_0^{t \wedge \tau} e^{\int_0^s (g_1+g_2)(\xi_r, t-r) dr} f_2(\xi_s, t-s) ds.$$

By assumption, $v(x, t) = \mathbb{P}_x[\mathcal{Y}_t + \mathcal{Z}_t]$ for all $(x, t) \in A \times [0, T]$. The Markov property implies

$$\int_0^t \mathbb{P}_x[\mathbb{1}_{\{s < \tau\}} e^{\int_0^s g_1(\xi_r, t-r) dr} g_2(\xi_s, t-s) \mathbb{P}_{\xi_s}[\mathcal{Y}_{t-s}]] ds = \int_0^t \mathbb{P}_x[\mathbb{1}_{\{s < \tau\}} e^{\int_0^s g_1(\xi_r, t-r) dr} g_2(\xi_s, t-s) e^{\int_s^{t \wedge \tau} (g_1+g_2)(\xi_r, t-r) dr} f_1(\xi_{t \wedge \tau})] ds.$$

If g_2 is bounded, then Fubini's theorem and the fundamental theorem of calculus (FTC) for Lebesgue integrals imply that the right-hand side equals

$$\mathbb{P}_x \left[f_1(\xi_{t \wedge \tau}) e^{\int_0^{t \wedge \tau} g_1(\xi_r, t-r) dr} \int_0^{t \wedge \tau} g_2(\xi_s, t-s) e^{\int_s^{t \wedge \tau} g_2(\xi_r, t-r) dr} ds \right] = \mathbb{P}_x [f_1(\xi_{t \wedge \tau}) e^{\int_0^{t \wedge \tau} g_1(\xi_r, t-r) dr} (e^{\int_0^{t \wedge \tau} g_2(\xi_r, t-r) dr} - 1)].$$

In the situation of (ii), the same identity can be obtained by truncating g_2 before the application of FTC and using the monotone convergence theorem afterwards. The Markov property and Fubini's theorem [in case (ii) its application is justified by the nonpositivity of the integrand] yield

$$\int_0^t \mathbb{P}_x[\mathbb{1}_{\{s < \tau\}} e^{\int_0^s g_1(\xi_r, t-r) dr} g_2(\xi_s, t-s) \mathbb{P}_{\xi_s}[\mathcal{Z}_{t-s}]] ds = \int_0^t \mathbb{P}_x \left[\mathbb{1}_{\{s < \tau\}} e^{\int_0^s g_1(\xi_r, t-r) dr} g_2(\xi_s, t-s) \times \int_s^{t \wedge \tau} e^{\int_s^u (g_1+g_2)(\xi_r, t-r) dr} f_2(\xi_u, t-u) du \right] ds$$

$$= \mathbb{P}_x \left[\int_0^{t \wedge \tau} e^{\int_0^u g_1(\xi_r, t-r) dr} f_2(\xi_u, t-u) \times \int_0^u g_2(\xi_s, t-s) e^{\int_s^u g_2(\xi_r, t-r) dr} ds du \right].$$

As above, the FTC implies that the right-hand side equals

$$\mathbb{P}_x \left[\int_0^{t \wedge \tau} e^{\int_0^u g_1(\xi_r, t-r) dr} f_2(\xi_u, t-u) (e^{\int_0^u g_2(\xi_r, t-r) dr} - 1) du \right].$$

Since $v(x, t) = \mathbb{P}_x[\mathcal{Y}_t + \mathcal{Z}_t]$ for all $(x, t) \in A \times [0, T]$, we conclude that in the situation of (i),

$$\begin{aligned} & \mathbb{P}_x \left[\int_0^{t \wedge \tau} e^{\int_0^s g_1(\xi_r, t-r) dr} (f_2(\xi_s, t-s) + g_2(\xi_s, t-s)v(\xi_s, t-s)) ds \right] \\ &= \mathbb{P}_x \left[f_1(\xi_{t \wedge \tau}) e^{\int_0^{t \wedge \tau} g_1(\xi_r, t-r) dr} (e^{\int_0^{t \wedge \tau} g_2(\xi_r, t-r) dr} - 1) \right. \\ & \quad \left. + \int_0^{t \wedge \tau} e^{\int_0^s (g_1+g_2)(\xi_r, t-r) dr} f_2(\xi_s, t-s) ds \right]. \end{aligned}$$

In the situation of (ii), this use of linearity is justified since none of the summed integrals can take the value $+\infty$. Since f_1 is bounded and $g_1, g_1 + g_2$ are bounded from above, the first term on the right can be written as the difference of two finite integrals and (A.1) yields the claim. \square

A.2. Monotonicity. In this section, a generalized version of the mild equation (1.3) is studied. For the exit measures \tilde{X}_t^β from Section 2.1.1, we establish the monotonicity in the domain, and we prove Lemma 4.5. As a by-product, we re-prove the existence and uniqueness of the solutions to (1.3) and (2.7).

To this end, we assume only that β is bounded above, not necessarily from below. More specifically, the setup is as follows. Let $\beta : D \rightarrow \mathbb{R}$ be measurable with $\bar{\beta} = \max\{\sup_{x \in D} \beta(x), 0\} < \infty$, $\alpha \in bp(D)$, Π a kernel from D to $(0, \infty)$ such that $x \mapsto \int_{(0, \infty)} (y \wedge y^2) \Pi(x, dy)$ belongs to $bp(D)$, and let $(\xi; \mathbb{P})$ be a diffusion as described in Section 1.1. We denote by $lbp(D \times [0, \infty))$ the space of all functions $f \in p(D \times [0, \infty))$ with $\|f\|_{\infty, T} := \sup_{t \in [0, T]} \|f(\cdot, t)\|_{\infty} < \infty$ for all $T > 0$.

For $f \in bp(D)$ and $g \in lbp(D \times [0, \infty))$, we are interested in solutions to the integral equation

$$(A.2) \quad u(x, t) + \int_0^t S_s[\psi_0(\cdot, u(\cdot, t-s))](x) ds = S_t f(x) + \int_0^t S_s[g(\cdot, t-s)](x) ds, \quad (x, t) \in D \times [0, \infty),$$

where $S_t f(x) = \mathbb{P}_x[e^{\int_0^t \beta(\xi_s) ds} f(\xi_t)]$ for $f \in p(D)$. A similar analysis to ours has been carried out by Dynkin (Chapter 4, Sections 1 and 3 in [11]) for the case $\beta = 0$

and $g = \mathbf{0}$. The greater generality of (A.2) allows us to handle the general setup of the present article and to prove Lemma 4.5.

Note that $z \mapsto \psi_0(x, z)$, defined in (1.1) is increasing, convex and nonnegative. In particular, any nonnegative solution u to (A.2), satisfies

$$0 \leq u(x, t) \leq e^{\bar{\beta}t} \|f\|_\infty + \int_0^t e^{\bar{\beta}s} \|g(\cdot, t-s)\|_\infty ds \quad \text{for all } (x, t) \in D \times [0, \infty).$$

Hence any nonnegative solution to (A.2) is an element of $lbp(D \times [0, \infty))$. Moreover, ψ_0 is locally Lipschitz continuous in the sense that for every fixed $c > 0$, there exists $\mathcal{L}(c) \in [0, \infty)$ such that

$$(A.3) \quad |\psi_0(x, z_1) - \psi_0(x, z_2)| \leq \mathcal{L}(c)|z_1 - z_2| \quad \text{for all } z_1, z_2 \in [0, c], x \in D.$$

We use the following version of Gronwall’s lemma. For a proof see Theorem A.5.1 in [25].

LEMMA A.2 (Gronwall’s lemma). *Let $T > 0, C, \rho \geq 0$ and $h \in b([0, T])$. If*

$$h(t) \leq C + \rho \int_0^t h(s) ds \quad \text{for all } t \in [0, T],$$

then $h(t) \leq Ce^{\rho t}$ for all $t \in [0, T]$.

Lemmas A.3 and A.4 in the case $\beta = \mathbf{0}$ and $g = \mathbf{0}$ are Theorems 4.1.1 and 4.3.1 in [12].

LEMMA A.3 (Uniqueness). *Let $f, \hat{f} \in bp(D)$, $g, \hat{g} \in lbp(D \times [0, \infty))$, and suppose that u and \hat{u} are nonnegative solutions to (A.2) for (f, g) and (\hat{f}, \hat{g}) , respectively. Then for every $T > 0$, there exists a constant $C > 0$ such that*

$$\|u - \hat{u}\|_{\infty, T} \leq C(\|f - \hat{f}\|_\infty + \|g - \hat{g}\|_{\infty, T}).$$

In particular, the solution to (A.2) is unique.

PROOF. Fix $T > 0$, and let $c \geq \max\{\|u\|_{\infty, T}, \|\hat{u}\|_{\infty, T}\}$. Then (A.3) yields

$$|\psi_0(x, \hat{u}(x, t)) - \psi_0(x, u(x, t))| \leq \mathcal{L}(c)|\hat{u}(x, t) - u(x, t)|$$

for all $(x, t) \in D \times [0, T]$.

Writing $h(x, t) = |u(x, t) - \hat{u}(x, t)|$ and $M = e^{\bar{\beta}T}$, (A.2) implies that for all $(x, t) \in D \times [0, T]$,

$$h(x, t) \leq M\|f - \hat{f}\|_\infty + MT\|g - \hat{g}\|_{\infty, T} + \int_0^t M\mathcal{L}(c)\|h(\cdot, s)\|_\infty ds.$$

Lemma A.2 yields the claim. \square

LEMMA A.4 (Existence). *Let $f \in bp(D)$ and $g \in lbp(D \times [0, \infty))$. There exists a nonnegative solution $u \in lbp(D \times [0, \infty))$ to (A.2).*

PROOF. Fix $T > 0$ and let $M = e^{\bar{\beta}T}$. For $k \in [0, \infty)$ and $u \in lbp(D \times [0, T])$, that is, $u \in p(D \times [0, T])$ with $\|u\|_{\infty, T} < \infty$, we define for all $(x, t) \in D \times [0, T]$,

$$F_k u(x, t) = e^{-kt} S_t f(x) + \int_0^t e^{-ks} S_s [g(\cdot, t - s)](x) ds + \int_0^t e^{-ks} S_s [ku(\cdot, t - s) - \psi_0(\cdot, u(\cdot, t - s))](x) ds.$$

Let $c \geq M\|f\|_{\infty} + MT\|g\|_{\infty, T}$ and $k \geq \mathcal{L}(c)$. Write $v(x, t) = e^{\bar{\beta}t}\|f\|_{\infty} + \int_0^t e^{\bar{\beta}s} ds \|g\|_{\infty, T}$ for all $x \in D, t \in [0, T]$. We show the following:

- (i) $\mathbf{0} \leq F_k \mathbf{0} \leq v$ on $D \times [0, T]$.
- (ii) If $\mathbf{0} \leq u_1 \leq u_2 \leq v$ on $D \times [0, T]$, then $F_k u_1 \leq F_k u_2$ on $D \times [0, T]$.
- (iii) $F_k v \leq v$ on $D \times [0, T]$.

Indeed, $F_k \mathbf{0}(x, t) = e^{-kt} S_t f(x) + \int_0^t e^{-ks} S_s [g(\cdot, t - s)](x) ds \in [0, v(x, t)]$ since f and g are nonnegative and $k \geq 0$. For (ii), we use that $v(x, t) \leq c$ for all $(x, t) \in D \times [0, T]$ and (A.3) to obtain

$$\begin{aligned} &F_k u_2(x, t) - F_k u_1(x, t) \\ &= \int_0^t e^{-ks} S_s [k(u_2 - u_1)(\cdot, t - s) - (\psi_0(\cdot, u_2(\cdot, t - s)) - \psi_0(\cdot, u_1(\cdot, t - s)))](x) ds \\ &\geq \int_0^t e^{-ks} S_s [(k - \mathcal{L}(c))(u_2 - u_1)(\cdot, t - s)](x) ds \geq 0. \end{aligned}$$

To show (iii), we use that ψ_0 is nonnegative and the definition of v and Fubini's theorem to obtain

$$\begin{aligned} F_k v(x, t) &\leq e^{-kt} e^{\bar{\beta}t} \|f\|_{\infty} + \int_0^t e^{-ks} e^{\bar{\beta}s} \|g\|_{\infty, T} ds \\ &\quad + \int_0^t e^{-ks} e^{\bar{\beta}s} k \left(e^{\bar{\beta}(t-s)} \|f\|_{\infty} + \int_0^{t-s} e^{\bar{\beta}r} dr \|g\|_{\infty, T} \right) ds \\ &= \left(e^{-kt} + \int_0^t k e^{-ks} ds \right) e^{\bar{\beta}t} \|f\|_{\infty} \\ &\quad + \left(\int_0^t e^{(\bar{\beta}-k)s} ds + \int_0^t k e^{-ks} \int_s^t e^{\bar{\beta}r} dr ds \right) \|g\|_{\infty, T} \\ &= v(x, t). \end{aligned}$$

In the next step, we construct a solution to (A.2) via a Picard iteration. Let $u_0 = \mathbf{0}$ and $u_n = F_k u_{n-1}$ for all $n \in \mathbb{N}$. We show by induction that $\mathbf{0} \leq u_{n-1} \leq u_n \leq v$ on

$D \times [0, T]$ for all $n \in \mathbb{N}$. For $n = 1$, this is statement (i). The induction step follows from (ii)–(iii). In particular, $(u_n)_{n \in \mathbb{N}_0}$ has a pointwise limit u which is a fixed point of F_k by the dominated convergence theorem. Lemma A.1(i) applied to $g_1 = \beta$, which is bounded from above, and the bounded functions $g_2 = -k$, $f_1 = f$ and $f_2(x, t) = g(x, t) + ku(x, t) - \psi_0(x, u(x, t))$, shows that u solves (A.2). \square

Choosing $g = \mathbf{0}$, Lemmas A.3 and A.4 imply the existence of a unique solution to (1.3) and (2.7). The following lemma will be used in the proof of Lemma 4.5.

LEMMA A.5 [Monotonicity in (f, g)]. *Let $f, \hat{f} \in bp(D)$, $g, \hat{g} \in lbp(D \times [0, \infty))$ with $f \leq \hat{f}$ and $g \leq \hat{g}$ and denote by u and \hat{u} the unique solutions to (A.2) corresponding to (f, g) and (\hat{f}, \hat{g}) , respectively. Then $u \leq \hat{u}$.*

PROOF. Since the solution is unique according to Lemma A.3, the claim follows immediately from the construction of the solution via Picard iteration in the proof of Lemma A.4. \square

PROOF OF LEMMA 4.5(i). According to (1.3), u_f is the unique solution to (A.2) with $g = \mathbf{0}$. Moreover, (1.3) for \hat{u}_f and Lemma A.1(i) applied to $g_1 = \beta$, $g_2 = \hat{\beta} - \beta$, $f_1 = f$ and $f_2(x, t) = -\hat{\psi}_0(x, \hat{u}_f(x, t))$ implies that \hat{u}_f satisfies

$$\begin{aligned} \hat{u}_f(x, t) &= S_t f(x) + \int_0^t S_s [-\hat{\psi}_0(\cdot, \hat{u}_f(\cdot, t-s)) + (\hat{\beta} - \beta)\hat{u}_f(\cdot, t-s)](x) ds \\ &= S_t f(x) - \int_0^t S_s [\psi_0(\cdot, \hat{u}_f(\cdot, t-s))](x) ds \\ &\quad + \int_0^t S_s [\psi_\beta(\cdot, \hat{u}_f(\cdot, t-s)) - \hat{\psi}_{\hat{\beta}}(\cdot, \hat{u}_f(\cdot, t-s))](x) ds. \end{aligned}$$

In particular, \hat{u}_f solves (A.2) with $g(x, t) = \psi_\beta(x, \hat{u}_f(x, t)) - \hat{\psi}_{\hat{\beta}}(x, \hat{u}_f(x, t)) \geq 0$. Now Lemma A.5 yields the claim. \square

The following lemma is used in the proofs of Theorem 2.3 and Lemma 4.3.

LEMMA A.6 (Monotonicity in B). *Let $B \subset\subset D$ and $f \in bp(D)$ such that the support of f , $\text{supp}(f)$, is compactly embedded in B . There exists a unique nonnegative solution $u_f^B \in lbp(D \times [0, \infty))$ to*

$$\begin{aligned} (A.4) \quad u(x, t) &= \mathbb{P}_x \left[e^{\int_0^{t \wedge \tau_B} \beta(\xi_s) ds} f(\xi_{t \wedge \tau_B}) \right] \\ &\quad - \mathbb{P}_x \left[\int_0^{t \wedge \tau_B} e^{\int_0^s \beta(\xi_r) dr} \psi_0(\xi_s, u(\xi_s, t-s)) ds \right]. \end{aligned}$$

Moreover, if B_1 and B_2 are domains with $\text{supp}(f) \subseteq B_1 \subseteq B_2$, then $u_f^{B_1} \leq u_f^{B_2}$.

PROOF. Let $T > 0$, $c \geq e^{\beta T} \|f\|_\infty$, $k \geq \mathcal{L}(c)$ and for $u \in lbp(D \times [0, T])$, define

$$\begin{aligned}
 F_k u(x, t) &= \mathbb{P}_x \left[e^{\int_0^t [\beta(\xi_s) - k] ds} f(\xi_t) \mathbb{1}_{\{t < \tau_B\}} \right] \\
 &\quad + \int_0^t \mathbb{P}_x \left[e^{\int_0^s [\beta(\xi_r) - k] dr} [ku(\xi_s, t - s) - \psi_0(\xi_s, u(\xi_s, t - s))] \mathbb{1}_{\{s \leq \tau_B\}} \right] ds.
 \end{aligned}$$

Since $kz - \psi_0(x, z) \geq kz - \mathcal{L}(c)z \geq 0$ for all $z \in [0, c]$, $F_k u$ is increasing in B for all u with $u(x, t) \leq e^{\beta t} \|f\|_\infty =: v(x, t)$. As in Lemmas A.3 and A.4, the unique solution to (A.4) can be obtained as a pointwise limit of the increasing sequence $u_0 = \mathbf{0}$, $u_{n+1} = F_k u_n$ with $u_n \leq v$ for all n . Denote by $u_n^{(1)}$ and $u_n^{(2)}$ the iterates for the operators $F_k^{(1)}$, $F_k^{(2)}$ corresponding to B_1 and B_2 , respectively. We show by induction that $u_n^{(1)} \leq u_n^{(2)}$ for every n . For $n = 0$ this is trivial. For the induction step we first use the induction hypothesis and monotonicity of $F_k^{(1)}$ [see (ii) in the proof of Lemma A.4] and then the monotonicity of F_k in B to deduce

$$u_{n+1}^{(1)} = F_k^{(1)} u_n^{(1)} \leq F_k^{(1)} u_n^{(2)} \leq F_k^{(2)} u_n^{(2)} = u_{n+1}^{(2)}. \quad \square$$

For $B \subset\subset D$, the tuple $(L, \psi_{\beta^*}^*; B)$ satisfies the assumptions of Section 1.1, where the motion is killed at the boundary of B . Hence, Lemma A.6 implies that the $(L, \psi_{\beta^*}^*; D)$ -superprocess can be obtained as a distributional limit of $(L, \psi_{\beta^*}^*; B)$ -superprocesses using an increasing sequence of compactly embedded domains to approximate D ; cf. the argument before Corollary 6.2 in [43] or Lemma A2 and Theorem A1 in [19].

Acknowledgments. The research has been initiated at the workshop ‘‘Branching random walks and searching in trees (10w5085)’’ at Banff International Research Station (BIRS). Further parts of this research have been completed at the Institute for Mathematical Research (FIM)—ETH Zurich and at the Mathematisches Forschungsinstitut Oberwolfach (MFO). Andreas E. Kyprianou and Matthias Winkel thank BIRS and Maren Eckhoff and Andreas E. Kyprianou thank FIM and MFO for their hospitality.

REFERENCES

[1] ASMUSSEN, S. and HERING, H. (1976). Strong limit theorems for general supercritical branching processes with applications to branching diffusions. *Z. Wahrsch. Verw. Gebiete* **36** 195–212. MR0420889

[2] ATHREYA, K. B. (1968). Some results on multitype continuous time Markov branching processes. *Ann. Math. Statist.* **39** 347–357. MR0221600

[3] BERESTYCKI, J., KYPRIANOU, A. E. and MURILLO-SALAS, A. (2011). The prolific backbone for supercritical superprocesses. *Stochastic Process. Appl.* **121** 1315–1331. MR2794978

- [4] BERESTYCKI, N. (2009). *Recent Progress in Coalescent Theory. Ensaios Matemáticos [Mathematical Surveys]* **16**. Sociedade Brasileira de Matemática, Rio de Janeiro. MR2574323
- [5] BERTOIN, J., FONTBONA, J. and MARTÍNEZ, S. (2008). On prolific individuals in a supercritical continuous-state branching process. *J. Appl. Probab.* **45** 714–726. MR2455180
- [6] BIGGINS, J. D. (1992). Uniform convergence of martingales in the branching random walk. *Ann. Probab.* **20** 137–151. MR1143415
- [7] CHEN, Z.-Q., REN, Y.-X. and WANG, H. (2008). An almost sure scaling limit theorem for Dawson–Watanabe superprocesses. *J. Funct. Anal.* **254** 1988–2019. MR2397881
- [8] CHEN, Z.-Q. and SHIOZAWA, Y. (2007). Limit theorems for branching Markov processes. *J. Funct. Anal.* **250** 374–399. MR2352485
- [9] DUQUESNE, T. and WINKEL, M. (2007). Growth of Lévy trees. *Probab. Theory Related Fields* **139** 313–371. MR2322700
- [10] DYNKIN, E. B. (1993). On regularity of superprocesses. *Probab. Theory Related Fields* **95** 263–281. MR1214090
- [11] DYNKIN, E. B. (1993). Superprocesses and partial differential equations. *Ann. Probab.* **21** 1185–1262. MR1235414
- [12] DYNKIN, E. B. (2002). *Diffusions, Superdiffusions and Partial Differential Equations. American Mathematical Society Colloquium Publications* **50**. Amer. Math. Soc., Providence, RI. MR1883198
- [13] DYNKIN, E. B. and KUZNETSOV, S. E. (2004). \mathbb{N} -measures for branching exit Markov systems and their applications to differential equations. *Probab. Theory Related Fields* **130** 135–150. MR2092876
- [14] ENGLÄNDER, J. (2004). An example and a conjecture concerning scaling limits of superdiffusions. *Statist. Probab. Lett.* **66** 363–368. MR2045481
- [15] ENGLÄNDER, J. (2007). Branching diffusions, superdiffusions and random media. *Probab. Surv.* **4** 303–364. MR2368953
- [16] ENGLÄNDER, J. (2009). Law of large numbers for superdiffusions: The non-ergodic case. *Ann. Inst. Henri Poincaré Probab. Stat.* **45** 1–6. MR2500226
- [17] ENGLÄNDER, J., HARRIS, S. C. and KYPRIANOU, A. E. (2010). Strong law of large numbers for branching diffusions. *Ann. Inst. Henri Poincaré Probab. Stat.* **46** 279–298. MR2641779
- [18] ENGLÄNDER, J. and KYPRIANOU, A. E. (2004). Local extinction versus local exponential growth for spatial branching processes. *Ann. Probab.* **32** 78–99. MR2040776
- [19] ENGLÄNDER, J. and PINSKY, R. G. (1999). On the construction and support properties of measure-valued diffusions on $D \subseteq \mathbf{R}^d$ with spatially dependent branching. *Ann. Probab.* **27** 684–730. MR1698955
- [20] ENGLÄNDER, J., REN, Y.-X. and SONG, R. (2013). Weak extinction versus global exponential growth of total mass for superdiffusions. Preprint. Available at arXiv:1301.6842.
- [21] ENGLÄNDER, J. and TURAEV, D. (2002). A scaling limit theorem for a class of superdiffusions. *Ann. Probab.* **30** 683–722. MR1905855
- [22] ENGLÄNDER, J. and WINTER, A. (2006). Law of large numbers for a class of superdiffusions. *Ann. Inst. Henri Poincaré Probab. Stat.* **42** 171–185. MR2199796
- [23] ETHERIDGE, A. M. (2000). *An Introduction to Superprocesses. University Lecture Series* **20**. Amer. Math. Soc., Providence, RI. MR1779100
- [24] ETHERIDGE, A. M. and WILLIAMS, D. R. E. (2003). A decomposition of the $(1 + \beta)$ -superprocess conditioned on survival. *Proc. Roy. Soc. Edinburgh Sect. A* **133** 829–847. MR2006204
- [25] ETHIER, S. N. and KURTZ, T. G. (2005). *Markov Processes: Characterization and Convergence*, 2nd ed. Wiley, New York.
- [26] EVANS, S. N. (1993). Two representations of a conditioned superprocess. *Proc. Roy. Soc. Edinburgh Sect. A* **123** 959–971. MR1249698

- [27] EVANS, S. N. and O'CONNELL, N. (1994). Weighted occupation time for branching particle systems and a representation for the supercritical superprocess. *Canad. Math. Bull.* **37** 187–196. MR1275703
- [28] FITZSIMMONS, P. J. (1988). Construction and regularity of measure-valued Markov branching processes. *Israel J. Math.* **64** 337–361. MR0995575
- [29] FLEISCHMANN, K. and MUELLER, C. (2004). Super-Brownian motion with extra birth at one point. *SIAM J. Math. Anal.* **36** 740–772 (electronic). MR2111914
- [30] FLEISCHMANN, K. and SWART, J. M. (2003). Extinction versus exponential growth in a supercritical super-Wright–Fisher diffusion. *Stochastic Process. Appl.* **106** 141–165. MR1983047
- [31] FLEISCHMANN, K. and SWART, J. M. (2004). Trimmed trees and embedded particle systems. *Ann. Probab.* **32** 2179–2221. MR2073189
- [32] GREVEN, A., KLENKE, A. and WAKOLBINGER, A. (2001). Interacting Fisher–Wright diffusions in a catalytic medium. *Probab. Theory Related Fields* **120** 85–117. MR1856196
- [33] GREY, D. R. (1974). Asymptotic behaviour of continuous time, continuous state-space branching processes. *J. Appl. Probab.* **11** 669–677. MR0408016
- [34] GRUMMT, R. and KOLB, M. (2013). Law of large numbers for super-Brownian motions with a single point source. *Stochastic Process. Appl.* **123** 1183–1212. MR3016220
- [35] HARDY, R. and HARRIS, S. C. (2009). A spine approach to branching diffusions with applications to L^p -convergence of martingales. In *Séminaire de Probabilités XLII. Lecture Notes in Math.* **1979** 281–330. Springer, Berlin. MR2599214
- [36] HARRIS, S. C. and ROBERTS, M. I. (2014). A strong law of large numbers for branching processes: Almost sure spine events. *Electron. Commun. Probab.* **19** 1–6. MR3208326
- [37] KAPLAN, N. and ASMUSSEN, S. (1976). Branching random walks. II. *Stochastic Process. Appl.* **4** 15–31. MR0400430
- [38] KESTEN, H. and STIGUM, B. P. (1966). A limit theorem for multidimensional Galton–Watson processes. *Ann. Math. Statist.* **37** 1211–1223. MR0198552
- [39] KLENKE, A. (2008). *Probability Theory. A Comprehensive Course*. Springer, London. Translated from the 2006 German original. MR2372119
- [40] KOURITZIN, M. A. and REN, Y.-X. (2014). A strong law of large numbers for super-stable processes. *Stochastic Process. Appl.* **124** 505–521. MR3131303
- [41] KYPRIANOU, A. E., LIU, R.-L., MURILLO-SALAS, A. and REN, Y.-X. (2012). Supercritical super-Brownian motion with a general branching mechanism and travelling waves. *Ann. Inst. Henri Poincaré Probab. Stat.* **48** 661–687. MR2976558
- [42] KYPRIANOU, A. E. and MURILLO-SALAS, A. (2013). Super-Brownian motion: L^p -convergence of martingales through the pathwise spine decomposition. In *Advances in Superprocesses and Nonlinear PDEs. Springer Proc. Math. Stat.* **38** 113–121. Springer, New York. MR3111226
- [43] KYPRIANOU, A. E., PEREZ, J.-L. and REN, Y.-X. (2013). The backbone decomposition for spatially dependent supercritical superprocesses. Preprint. Available at arXiv:1304.2019v1.
- [44] KYPRIANOU, A. E. and REN, Y.-X. (2012). Backbone decomposition for continuous-state branching processes with immigration. *Statist. Probab. Lett.* **82** 139–144. MR2863035
- [45] LIU, R.-L., REN, Y.-X. and SONG, R. (2009). $L \log L$ criterion for a class of superdiffusions. *J. Appl. Probab.* **46** 479–496. MR2535827
- [46] LIU, R.-L., REN, Y.-X. and SONG, R. (2013). Strong law of large numbers for a class of superdiffusions. *Acta Appl. Math.* **123** 73–97. MR3010225
- [47] MIŁOŚ, P. (2012). Spatial CLT for the supercritical Ornstein–Uhlenbeck superprocess. Preprint. Available at arXiv:1203.6661.

- [48] PERKINS, E. (2002). Dawson–Watanabe superprocesses and measure-valued diffusions. In *Lectures on Probability Theory and Statistics (Saint-Flour, 1999)*. *Lecture Notes in Math.* **1781** 125–324. Springer, Berlin. MR1915445
- [49] PINCHOVER, Y. (2013). Some aspects of large time behavior of the heat kernel: An overview with perspectives. In *Mathematical Physics, Spectral Theory and Stochastic Analysis* 299–339. Birkhäuser/Springer, Basel. MR3077281
- [50] PINSKY, R. G. (1995). *Positive Harmonic Functions and Diffusion*. *Cambridge Studies in Advanced Mathematics* **45**. Cambridge Univ. Press, Cambridge. MR1326606
- [51] PINSKY, R. G. (1996). Transience, recurrence and local extinction properties of the support for supercritical finite measure-valued diffusions. *Ann. Probab.* **24** 237–267. MR1387634
- [52] REN, Y.-X., SONG, R. and ZHANG, R. (2014). Central limit theorems for super Ornstein–Uhlenbeck processes. *Acta Appl. Math.* **130** 9–49. MR3180938
- [53] REVUZ, D. and YOR, M. (1999). *Continuous Martingales and Brownian Motion*, 3rd ed. *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]* **293**. Springer, Berlin. MR1725357
- [54] SALISBURY, T. S. and SEZER, A. D. (2013). Conditioning super-Brownian motion on its boundary statistics, and fragmentation. *Ann. Probab.* **41** 3617–3657. MR3127894
- [55] SALISBURY, T. S. and VERZANI, J. (1999). On the conditioned exit measures of super Brownian motion. *Probab. Theory Related Fields* **115** 237–285. MR1720367
- [56] SALISBURY, T. S. and VERZANI, J. (2000). Non-degenerate conditionings of the exit measures of super Brownian motion. *Stochastic Process. Appl.* **87** 25–52. MR1751163
- [57] SHEU, Y.-C. (1997). Lifetime and compactness of range for super-Brownian motion with a general branching mechanism. *Stochastic Process. Appl.* **70** 129–141. MR1472962
- [58] STROOCK, D. W. and VARADHAN, S. R. S. (1979). *Multidimensional Diffusion Processes*. *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]* **233**. Springer, New York. MR0532498
- [59] WANG, L. (2010). An almost sure limit theorem for super-Brownian motion. *J. Theoret. Probab.* **23** 401–416. MR2644866
- [60] WATANABE, S. (1967). Limit theorem for a class of branching processes. In *Markov Processes and Potential Theory* 205–232. Wiley, New York. MR0237007

M. ECKHOFF
 A. E. KYPRIANOU
 DEPARTMENT OF MATHEMATICAL SCIENCES
 UNIVERSITY OF BATH
 BATH BA2 7AY
 UNITED KINGDOM
 E-MAIL: m.eckhoff@bath.ac.uk
 a.kyprianou@bath.ac.uk

M. WINKEL
 DEPARTMENT OF STATISTICS
 UNIVERSITY OF OXFORD
 OXFORD OX1 3TG
 UNITED KINGDOM
 E-MAIL: winkel@stats.ox.ac.uk