

Analyticity of Entropy Rates of Continuous-State Hidden Markov Models

Vladislav Z. B. Tadić¹ and Arnaud Doucet

Abstract—The analyticity of the entropy and relative entropy rates of continuous-state hidden Markov models is studied here. Using the analytic continuation principle and the stability properties of the optimal filter, the analyticity of these rates is established for analytically parameterized models. The obtained results hold under relatively mild conditions and cover several useful classes of hidden Markov models. These results are relevant for several theoretically and practically important problems arising in statistical inference, system identification and information theory.

Index Terms—Hidden Markov models, entropy rate, relative entropy rate, log-likelihood, optimal filter, analytic continuation.

I. INTRODUCTION

HIDDEN Markov models are a powerful and versatile tool for statistical modeling of complex time-series data and stochastic dynamic systems. They can be described as a discrete-time Markov chain observed through imperfect, noisy observations of its states. Proposed in the seminal paper [1] over five decades ago, hidden Markov models have found many applications in very diverse areas such as acoustics and signal processing, image analysis and computer vision, automatic control, economics and finance, computational biology, genetics and bioinformatics. Owing to their theoretical and practical importance, various aspects of hidden Markov models have been thoroughly studied in a number of papers and books — see, e.g., [2], [6], [8] and references therein.

The entropy and relative entropy rates of hidden Markov models can be considered as an information-theoretic characterization of the asymptotic properties of these models. The entropy rate of a hidden Markov model can be interpreted as a measure of the average information revealed by the model through noisy observations of the states. The relative entropy rate between two hidden Markov models can be viewed as a measure of discrepancy between these models. The entropy rates of hidden Markov models and their analytical properties have recently gained significant attention in the information theory community. These properties and

their links with statistical inference, system identification, stochastic optimization and information theory have been studied extensively in several papers [10] – [13], [14], [19], [20], [22], [23]. However, to the best of our knowledge, the existing results on the analytical properties of the entropy rates of hidden Markov models apply exclusively to scenarios where the hidden Markov chain takes values in a finite state-space. We establish here analytical properties of the entropy rates of continuous-state hidden Markov models. As indicated in [26], such results can be very useful when analyzing algorithms for statistical inference in hidden Markov models.

In many applications, a hidden Markov model depends on an unknown parameter whose value needs to be inferred from a set of state-observations. In online settings, the unknown parameter is typically estimated using the recursive maximum likelihood method [21], [23]. In [23], it has been shown that the convergence and convergence rate of recursive maximum likelihood estimation in finite-state hidden Markov models is closely linked to the analyticity of the underlying (average) log-likelihood, i.e. of the underlying relative entropy rate. In view of recent results on stochastic gradient search [25], a similar link is expected to hold for continuous-state hidden Markov models. However, to apply the results of [25] to recursive maximum likelihood estimation in continuous-state hidden Markov models, it is necessary to establish the analyticity of the average log-likelihood for these models. Hence, one of the first and most important steps to carry out the asymptotic analysis of recursive maximum likelihood estimation in continuous-state hidden Markov models is to show the analyticity of the entropy rates of such models. The results presented here should provide a theoretical basis for this step.

In this paper, we study analytically parameterized continuous-state hidden Markov models (i.e., the models whose state transition kernel and the observation conditional distribution are analytic in the model parameters). Using mixing conditions on the model dynamics, we construct a geometrically ergodic analytic continuation of the state transition kernel and an exponentially stable analytic continuation of the optimal filter. Relying on these continuations and their asymptotic properties, we demonstrate that the entropy and relative entropy rates are analytic in the model parameters. The obtained results hold under relatively mild conditions and cover a broad and common class of state-space and continuous-state hidden Markov models. Moreover, these results generalize the existing results on the analyticity of entropy rates of finite-state hidden Markov models. Additionally, the results presented here are relevant for several

Manuscript received September 5, 2018; revised July 17, 2019; accepted July 29, 2019. Date of publication August 16, 2019; date of current version November 20, 2019.

V. Z. B. Tadić is with the School of Mathematics, University of Bristol, Bristol BS8 1TW, U.K. (e-mail: v.b.tadic@bristol.ac.uk).

A. Doucet is with the Department of Statistics, University of Oxford, Oxford OX1 3LB, U.K. (e-mail: doucet@stats.ox.ac.uk).

Communicated by P. Harremoës, Associate Editor for Probability and Statistics.

Color versions of one or more of the figures in this article are available online at <http://ieeexplore.ieee.org>.

Digital Object Identifier 10.1109/TIT.2019.2935777

0018-9448 © 2019 IEEE. Personal use is permitted, but republication/redistribution requires IEEE permission.

See http://www.ieee.org/publications_standards/publications/rights/index.html for more information.

important problems related to statistical inference, system identification and information theory.

The rest of this paper is organized as follows. In Section II, the entropy rates of hidden Markov models are specified and the main results are presented. Examples illustrating the main results are provided in Sections III and IV. In Sections V – VII, the main results are proved.

II. MAIN RESULTS

To define hidden Markov models and their entropy rates, we use the following notations. (Ω, \mathcal{F}, P) is a probability space. $d_x \geq 1$ and $d_y \geq 1$ are integers, while $\mathcal{X} \subseteq \mathbb{R}^{d_x}$ and $\mathcal{Y} \subseteq \mathbb{R}^{d_y}$ are Borel sets. $P(x, dx')$ is a transition kernel on \mathcal{X} , while $Q(x, dy)$ is a conditional probability measure on \mathcal{Y} given $x \in \mathcal{X}$. A hidden Markov model can be defined as the $\mathcal{X} \times \mathcal{Y}$ -valued stochastic process $\{(X_n, Y_n)\}_{n \geq 0}$ which is defined on (Ω, \mathcal{F}, P) and satisfies

$$\begin{aligned} P((X_{n+1}, Y_{n+1}) \in B | X_{0:n}, Y_{0:n}) \\ = \int I_B(x, y) Q(x, dy) P(X_n, dx) \end{aligned}$$

almost surely for $n \geq 0$ and any Borel set $B \subseteq \mathcal{X} \times \mathcal{Y}$. $\{X_n\}_{n \geq 0}$ are the unobservable states, while $\{Y_n\}_{n \geq 0}$ are the observations. Y_n can be interpreted as a noisy measurement of state X_n . States $\{X_n\}_{n \geq 0}$ form a Markov chain, while $P(x, dx')$ is their transition kernel. Conditionally on $\{X_n\}_{n \geq 0}$, state-observations $\{Y_n\}_{n \geq 0}$ are mutually independent, while $Q(X_n, dy)$ is the conditional distribution of Y_n given $X_{0:n}$. For more details on hidden Markov models, see [2], [6] and references therein.

In addition to the model $\{(X_n, Y_n)\}_{n \geq 0}$, we also consider a parameterized family of hidden Markov models. To specify such a family, we rely on the following notations. Let $d \geq 1$ be an integer, while $\Theta \subset \mathbb{R}^d$ is an open set. $\mathcal{P}(\mathcal{X})$ is the set of probability measures on \mathcal{X} . $\mu(dx)$ and $\nu(dy)$ are measures on \mathcal{X} and \mathcal{Y} (respectively), while $p_\theta(x'|x)$ and $q_\theta(y|x)$ are functions which map $\theta \in \Theta$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$ to $[0, \infty)$ and satisfy

$$\int_{\mathcal{X}} p_\theta(x'|x) \mu(dx') = \int_{\mathcal{Y}} q_\theta(y|x) \nu(dy) = 1$$

for all $\theta \in \Theta$, $x \in \mathcal{X}$. A family of hidden Markov models can then be defined as a collection of $\mathcal{X} \times \mathcal{Y}$ -valued stochastic processes $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ on (Ω, \mathcal{F}, P) , parameterized by $\theta \in \Theta$, $\lambda \in \mathcal{P}(\mathcal{X})$ and satisfying

$$\begin{aligned} P((X_0^{\theta, \lambda}, Y_0^{\theta, \lambda}) \in B) &= \int \int I_B(x, y) q_\theta(y|x) \lambda(dx), \\ P((X_{n+1}^{\theta, \lambda}, Y_{n+1}^{\theta, \lambda}) \in B | X_{0:n}^{\theta, \lambda}, Y_{0:n}^{\theta, \lambda}) \\ &= \int \int I_B(x, y) q_\theta(y|x) p_\theta(x | X_n^{\theta, \lambda}) \mu(dx) \nu(dy) \end{aligned}$$

almost surely for $n \geq 0$ and any Borel set $B \subseteq \mathcal{X} \times \mathcal{Y}$. $\{X_n^{\theta, \lambda}\}_{n \geq 0}$ are the hidden states of this model, while $\{Y_n^{\theta, \lambda}\}_{n \geq 0}$ are the corresponding observations. $p_\theta(x'|x)$ is the transition density of the Markov chain $\{X_n^{\theta, \lambda}\}_{n \geq 0}$, while $q_\theta(y|x)$ is the conditional density of $Y_n^{\theta, \lambda}$ given $X_{0:n}^{\theta, \lambda}$. In the context of the identification of stochastic dynamical systems and parameter

estimation in time-series models, $\{(X_n, Y_n)\}_{n \geq 0}$ is interpreted as the true system (or true model), while $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ is viewed as a candidate model for $\{(X_n, Y_n)\}_{n \geq 0}$.

To define the entropy rates of hidden Markov models, we introduce further notations. $r_\theta(y, x'|x)$ is the transition density of $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$, i.e.,

$$r_\theta(y, x'|x) = q_\theta(y|x') p_\theta(x'|x)$$

for $\theta \in \Theta$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$. $q_\theta^n(y_{1:n}|\lambda)$ is the density of $Y_{1:n}^{\theta, \lambda}$, i.e.,

$$\begin{aligned} q_\theta^n(y_{1:n}|\lambda) &= \int \cdots \int \int \left(\prod_{k=1}^n r_\theta(y_k, x_k | x_{k-1}) \right) \\ &\quad \cdot \mu(dx_n) \cdots \mu(dx_1) \lambda(dx_0), \end{aligned}$$

where $\lambda \in \mathcal{P}(\mathcal{X})$, $y_{1:n} = (y_1, \dots, y_n) \in \mathcal{Y}^n$, $n \geq 1$. The (average) entropy $h_n(\theta, \lambda)$ of $Y_{1:n}^{\theta, \lambda}$ is given by

$$h_n(\theta, \lambda) = -E \left(\frac{1}{n} \log q_\theta^n(Y_{1:n}^{\theta, \lambda} | \lambda) \right). \quad (1)$$

The expected (average) log-likelihood $l_n(\theta, \lambda)$ of $Y_{1:n}$ given the model $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ is specified as

$$l_n(\theta, \lambda) = E \left(\frac{1}{n} \log q_\theta^n(Y_{1:n} | \lambda) \right). \quad (2)$$

The entropy rate of model $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ (i.e., the entropy rate of stochastic process $\{Y_n^{\theta, \lambda}\}_{n \geq 0}$) can then be defined as the limit

$$\lim_{n \rightarrow \infty} h_n(\theta, \lambda).$$

Similarly, the relative entropy rate between models $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ and $\{(X_n, Y_n)\}_{n \geq 0}$ (i.e., the relative entropy rate between stochastic processes $\{Y_n^{\theta, \lambda}\}_{n \geq 0}$ and $\{Y_n\}_{n \geq 0}$) can be defined as the limit

$$- \lim_{n \rightarrow \infty} (l_n(\theta, \lambda) + h),$$

where h is the entropy rate of $\{Y_n\}_{n \geq 0}$ (provided h exists). In this context, the limit

$$\lim_{n \rightarrow \infty} l_n(\theta, \lambda)$$

can be viewed/referred to as the log-likelihood rate of $\{Y_n\}_{n \geq 0}$ given the model $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$. Entropy rate $\lim_{n \rightarrow \infty} h_n(\theta, \lambda)$ can be considered as a measure of the information revealed by the model $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ through its state-observations $\{Y_n^{\theta, \lambda}\}_{n \geq 0}$. Relative entropy rate $-\lim_{n \rightarrow \infty} (l_n(\theta, \lambda) + h)$ can be interpreted as a measure of discrepancy between the models $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ and $\{(X_n, Y_n)\}_{n \geq 0}$. The entropy rates of hidden Markov models are closely related to a number of important problems arising in engineering and statistics such as system identification, parameter estimation, model reduction and data compression. For example, in the recursive maximum likelihood approach to the identification of stochastic dynamical systems and parameter estimation in time-series models, the candidate model $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ providing the best approximation to the true model $\{(X_n, Y_n)\}_{n \geq 0}$ is selected through the minimization of

– $\lim_{n \rightarrow \infty} (l_n(\theta, \lambda) + h)$ (i.e., through the maximization of $\lim_{n \rightarrow \infty} l_n(\theta, \lambda)$). For more details on the entropy rates and their applications, see [7], [9] and references therein.

We study here the rates $\lim_{n \rightarrow \infty} h_n(\theta, \lambda)$, $\lim_{n \rightarrow \infty} l_n(\theta, \lambda)$ and their analytical properties. To formulate the assumptions under which these rates are analyzed, we rely on the following notations. For $\eta \in \mathbb{C}^d$, $\|\eta\|$ denotes the Euclidean norm of η . For $\gamma \in (0, 1)$, $V_\gamma(\Theta)$ is the open γ -vicinity of Θ in \mathbb{C}^d , i.e.,

$$V_\gamma(\Theta) = \{\eta \in \mathbb{C}^d : \exists \theta \in \Theta, \|\eta - \theta\| < \gamma\}.$$

Our analysis is based on the following assumptions.

Assumption 2.1. *There exists a real number $\varepsilon \in (0, 1)$ and for each $\theta \in \Theta$, $y \in \mathcal{Y}$, there exists a finite measure $\lambda_\theta(dx|y)$ on \mathcal{X} such that*

$$\varepsilon \lambda_\theta(B|y) \leq \int_B r_\theta(y, x'|x) \mu(dx') \leq \frac{\lambda_\theta(B|y)}{\varepsilon}$$

for all $x \in \mathcal{X}$ and any Borel set $B \subseteq \mathcal{X}$.

Assumption 2.2. *$r_\theta(y, x'|x)$ is real-analytic in θ for each $\theta \in \Theta$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$. Moreover, $r_\theta(y, x'|x)$ has a complex-valued continuation $\hat{r}_\theta(y, x'|x)$ with the following properties:*

- (i) $\hat{r}_\theta(y, x'|x)$ maps $\eta \in \mathbb{C}^d$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$ to \mathbb{C} .
- (ii) $\hat{r}_\theta(y, x'|x) = r_\theta(y, x'|x)$ for all $\theta \in \Theta$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$.
- (iii) There exists a real number $\delta \in (0, 1)$ such that $\hat{r}_\eta(y, x'|x)$ is analytic in η for each $\eta \in V_\delta(\Theta)$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$.
- (iv) There exists a function $\varphi_\eta(y)$ which maps $\eta \in \mathbb{C}^d$, $y \in \mathcal{Y}$ to \mathbb{C} , is analytic in η for each $\eta \in V_\delta(\Theta)$, $y \in \mathcal{Y}$ and satisfies

$$\varphi_\eta(y) \neq 0, \quad |\hat{r}_\eta(y, x'|x)| \leq |\varphi_\eta(y)|$$

for all $\eta \in V_\delta(\Theta)$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$.

(v) There exist functions $\phi, \psi : \mathcal{Y} \rightarrow (0, \infty)$ such that $\int \phi(y) \nu(dy) < \infty$ and

$$|\varphi_\eta(y)| \leq \phi(y), \quad |\log |\varphi_\eta(y)|| \leq \psi(y)$$

for all $\eta \in V_\delta(\Theta)$, $y \in \mathcal{Y}$.

Assumption 2.3. *There exists a real number $\gamma \in (0, 1)$ such that*

$$\int r_\theta(y, x'|x) \mu(dx') \geq \gamma |\varphi_\theta(y)|$$

for all $\theta \in \Theta$, $x \in \mathcal{X}$, $y \in \mathcal{Y}$.

Assumption 2.4. $\int \psi(y) \phi(y) \nu(dy) < \infty$.

Assumption 2.5. *There exists a real number $K \in [1, \infty)$ such that*

$$\int \psi(y) \mathcal{Q}(x, dy) \leq K$$

for all $x \in \mathcal{X}$. Moreover, there exist a probability measure $\pi(dx)$ on \mathcal{X} and a real number $\rho \in (0, 1)$ such that

$$|P^n(x, B) - \pi(B)| \leq K \rho^n \quad (3)$$

for all $x \in \mathcal{X}$, $n \geq 0$ and any Borel-set $B \subseteq \mathcal{X}$.

Assumption 2.1 is related to the stability of the hidden Markov model $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ and its optimal filter. This assumption ensures that the Markov chain $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ is geometrically ergodic (see Lemma 5.4) and that the optimal filter for the model $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ forgets initial conditions at an exponential rate (see Lemma 6.2). In this or similar form, Assumption 2.1 is an ingredient of a number of asymptotic results on optimal filtering and maximum likelihood estimation in hidden Markov models (see [4], [5], [16], [17]).

Assumption 2.2 is a condition on the parameterization of the model $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$. It requires the transition kernel and density of the chain $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ to be real-analytic in parameter θ . Together with Assumption 2.1, Assumption 2.2 ensures that an analytic continuation of this kernel exists and is geometrically ergodic (see Lemma 5.4).

Assumption 2.3 is also related to the parameterization of the model $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$. This assumption ensures that the ratio

$$\frac{r_\theta(y, x'|x)}{\int r_\theta(y, x''|x) \mu(dx'')}$$

is uniformly bounded in θ, x, x' . Together with Assumptions 2.1 and 2.2, Assumption 2.3 ensures that an analytic continuation of the optimal filter for the model $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ exists and forgets initial conditions at an exponential rate (see Lemma 6.6).

Assumption 2.4 requires the product of the bounding functions $\phi(y)$, $\psi(y)$ to be integrable with respect to the measure $\nu(dy)$. Together with Assumption 2.2, Assumption 2.4 ensures that the entropy $h_n(\theta, \lambda)$ exists and has an analytic continuation in θ (see Lemma 7.2).

Assumption 2.5 ensures that the Markov chain $\{(X_n, Y_n)\}_{n \geq 0}$ is geometrically ergodic (see Lemma 5.1). Together with Assumption 2.5, Assumption 2.2 also ensures that the log-likelihood $l_n(\theta, \lambda)$ defined in (2) exists and admits an analytic continuation.

The following two theorems are the main results of the paper.

Theorem 2.1. *Let Assumptions 2.1 – 2.3 and 2.5 hold. Then, there exists a function $l : \Theta \rightarrow \mathbb{R}$ such that $l(\theta)$ is real-analytic for each $\theta \in \Theta$ and $l(\theta) = \lim_{n \rightarrow \infty} l_n(\theta, \lambda)$ for all $\theta \in \Theta$, $\lambda \in \mathcal{P}(\mathcal{X})$.*

Theorem 2.2. *Let Assumptions 2.1 – 2.4 hold. Then, there exists a function $h : \Theta \rightarrow \mathbb{R}$ such that $h(\theta)$ is real-analytic for each $\theta \in \Theta$ and $h(\theta) = \lim_{n \rightarrow \infty} h_n(\theta, \lambda)$ for all $\theta \in \Theta$, $\lambda \in \mathcal{P}(\mathcal{X})$.*

Remark. *As Θ can be represented as a union of open balls, it is sufficient to show Theorems 2.1 and 2.2 for the case where Θ is convex and bounded. Therefore, throughout the analysis carried out in Sections V – VIII, we assume that Θ is a bounded open convex set.*

Theorems 2.1 and 2.2 are proved in Section VII. According to these theorems, for all $\theta \in \Theta$, $\lambda \in \mathcal{P}(\mathcal{X})$, rates $\lim_{n \rightarrow \infty} h_n(\theta, \lambda)$ and $\lim_{n \rightarrow \infty} l_n(\theta, \lambda)$ are well-defined.

Moreover, for each $\theta \in \Theta$, the rates $\lim_{n \rightarrow \infty} h_n(\theta, \lambda)$ and $\lim_{n \rightarrow \infty} l_n(\theta, \lambda)$ are independent of λ and real-analytic in θ .

The analytical properties of the entropy rates of hidden Markov models have already been extensively studied in several papers [10] – [14], [19], [20], [22], [23]. However, the results presented therein apply exclusively to models with finite state-spaces. To the best of our knowledge, Theorems 2.1 and 2.2 are the first results on the analyticity of the entropy rates of continuous-state hidden Markov models. These theorems also generalize the existing results on the analyticity of the entropy rates of finite-state hidden Markov models. More specifically, [12] can be considered as the strongest existing result of this kind. Theorem 2.2 includes, as a particular case, the results of [12] and simplifies the conditions under which these results hold (see Appendix 2 for details). Theorems 2.1 and 2.2 are relevant for several theoretically and practically important problems arising in statistical inference and system identification. In [26], we rely on these theorems to analyze recursive maximum likelihood estimation in non-linear state-space models. The same theorems can also be used to study the higher-order statistical asymptotics for maximum likelihood estimation in time-series models (for details on such asymptotics, see [27]).

III. EXAMPLE: MIXTURE OF DENSITIES

In this section, the main results are applied to the case when $p_\theta(x'|x)$ and $q_\theta(y|x)$ are mixtures of probability densities, i.e.,

$$p_\theta(x'|x) = \sum_{i=1}^{N_x} a_\theta^i(x) v_i(x'), \quad (4)$$

$$q_\theta(y|x) = \sum_{j=1}^{N_y} b_\theta^j(x) w_j(y) \quad (5)$$

for $\theta \in \Theta$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$ and integers $N_x > 1$ and $N_y > 1$. Here Θ , \mathcal{X} , \mathcal{Y} have the same meaning as in the previous section. $\{v_i(x)\}_{1 \leq i \leq N_x}$ and $\{w_j(y)\}_{1 \leq j \leq N_y}$ are functions which map $x \in \mathcal{X}$, $y \in \mathcal{Y}$ to $[0, \infty)$ and satisfy

$$\int v_i(x) \mu(dx) = \int w_j(y) \nu(dy) = 1$$

for each $1 \leq i \leq N_x$, $1 \leq j \leq N_y$ ($\mu(dx)$, $\nu(dy)$ have the same meaning as in the previous section). $\{a_\theta^i(x)\}_{1 \leq i \leq N_x}$ and $\{b_\theta^j(x)\}_{1 \leq j \leq N_y}$ are functions which map $\theta \in \Theta$, $x \in \mathcal{X}$ to $[0, \infty)$ and satisfy

$$\sum_{i=1}^{N_x} a_\theta^i(x) = \sum_{j=1}^{N_y} b_\theta^j(x) = 1.$$

Under these conditions, $v_i(x)$ and $w_j(y)$ are probability densities on \mathcal{X} and \mathcal{Y} (respectively), while $a_\theta^i(x)$ and $b_\theta^j(x)$ are probability masses in i and j (respectively). Hence, in x' , y , $p_\theta(x'|x)$ and $q_\theta(y|x)$ are mixtures of probability densities. $v_i(x)$ and $w_j(y)$ are the components of these mixtures, while $a_\theta^i(x)$ and $b_\theta^j(x)$ are the corresponding weights.

The entropy rates of hidden Markov model specified in (4), (5) are studied under the following assumptions.

Assumption 3.1. \mathcal{X} is a compact set.

Assumption 3.2. $a_\theta^i(x) > 0$ and $b_\theta^j(x) > 0$ for all $\theta \in \Theta$, $x \in \mathcal{X}$, $1 \leq i \leq N_x$, $1 \leq j \leq N_y$. Moreover, $a_\theta^i(x)$ and $b_\theta^j(x)$ are real-analytic in (θ, x) for each $\theta \in \Theta$, $x \in \mathcal{X}$, $1 \leq i \leq N_x$, $1 \leq j \leq N_y$.

Assumption 3.3. There exists a real number $\varepsilon \in (0, 1)$ such that $\varepsilon \leq v_i(x) \leq 1/\varepsilon$ for all $x \in \mathcal{X}$, $1 \leq i \leq N_x$.

Assumption 3.4. $\int |\log w_k(y)| w_j(y) \nu(dy) < \infty$ for each $1 \leq j, k \leq N_y$.

Assumption 3.5. There exists a real number $K \in [1, \infty)$ such that

$$\int |\log w_k(y)| Q(x, dy) \leq K$$

for all $x \in \mathcal{X}$, $1 \leq k \leq N_y$. Moreover, there exist a probability measure $\pi(dx)$ on \mathcal{X} and a real number $\rho \in (0, 1)$ such that (3) holds for all $x \in \mathcal{X}$, $n \geq 0$ and any Borel-measurable set $B \subseteq \mathcal{X}$.

Assumptions 3.1 – 3.5 cover several classes of hidden Markov models met in practice. These assumptions indeed hold if $q_\theta(y|x)$ is a mixture of Gamma, Gaussian, Pareto and logistic distributions, and if $p_\theta(x'|x)$ is a mixture of the same distributions truncated to a compact domain.

Using Theorem 2.1 and Theorem 2.2, we obtain the following results.

Corollary 3.1. Let Assumptions 3.1 – 3.3 and 3.5 hold. Then, all conclusions of Theorem 2.1 are true.

Corollary 3.2. Let Assumptions 3.1 – 3.4 hold. Then, all conclusions of Theorem 2.2 are true.

Corollaries 3.1 and 3.2 are proved in Section VIII.

IV. EXAMPLE: NON-LINEAR STATE-SPACE MODELS

In this section, the main results are used to study the entropy rates of non-linear state-space models. We consider the following parameterized state-space model:

$$X_{n+1}^{\theta, \lambda} = A_\theta(X_n^{\theta, \lambda}) + B_\theta(X_n^{\theta, \lambda}) V_n, \quad (6)$$

$$Y_n^{\theta, \lambda} = C_\theta(X_n^{\theta, \lambda}) + D_\theta(X_n^{\theta, \lambda}) W_n, \quad n \geq 0. \quad (7)$$

Here $\theta \in \Theta$, $\lambda \in \mathcal{P}(\mathcal{X})$ are the parameters indexing the state-space model (6), (7) (Θ , $\mathcal{P}(\mathcal{X})$ have the same meaning as in Section II). $A_\theta(x)$ and $B_\theta(x)$ are functions which map $\theta \in \Theta$, $x \in \mathbb{R}^{d_x}$ (respectively) to \mathbb{R}^{d_x} and $\mathbb{R}^{d_x \times d_x}$ (d_x has the same meaning as in Section II). $C_\theta(x)$ and $D_\theta(x)$ are functions which map $\theta \in \Theta$, $x \in \mathbb{R}^{d_x}$ (respectively) to \mathbb{R}^{d_y} and $\mathbb{R}^{d_y \times d_y}$ (d_y has the same meaning as in Section II). $X_0^{\theta, \lambda}$ is an \mathbb{R}^{d_x} -valued random variable defined on a probability space (Ω, \mathcal{F}, P) and distributed according to λ . $\{V_n\}_{n \geq 0}$ are \mathbb{R}^{d_x} -valued i.i.d. random variables which are defined on (Ω, \mathcal{F}, P) and have (marginal) probability density $v(x)$ with respect to the Lebesgue measure. $\{W_n\}_{n \geq 0}$ are \mathbb{R}^{d_y} -valued i.i.d. random variables which are defined on (Ω, \mathcal{F}, P) and have (marginal) probability density $w(y)$ with respect to the Lebesgue measure. We also assume that $X_0^{\theta, \lambda}$, $\{V_n\}_{n \geq 0}$ and $\{W_n\}_{n \geq 0}$ are (jointly) independent.

We use here the following notations. For $\theta \in \Theta$, $x, x' \in \mathbb{R}^{d_x}$, $y \in \mathbb{R}^{d_y}$, $\tilde{p}_\theta(x'|x)$ and $\tilde{q}_\theta(y|x)$ are the functions defined by

$$\tilde{p}_\theta(x'|x) = \frac{v \left(B_\theta^{-1}(x)(x' - A_\theta(x)) \right)}{|\det B_\theta(x)|},$$

$$\tilde{q}_\theta(y|x) = \frac{w \left(D_\theta^{-1}(x)(y - C_\theta(x)) \right)}{|\det D_\theta(x)|}$$

(provided that $B_\theta(x)$, $D_\theta(x)$ are invertible), while $p_\theta(x'|x)$ and $q_\theta(y|x)$ are defined by

$$p_\theta(x'|x) = \frac{v \left(B_\theta^{-1}(x)(x' - A_\theta(x)) \right) 1_{\mathcal{X}}(x')}{\int_{\mathcal{X}} v \left(B_\theta^{-1}(x)(x'' - A_\theta(x)) \right) dx''}, \quad (8)$$

$$q_\theta(y|x) = \frac{w \left(D_\theta^{-1}(x)(y - C_\theta(x)) \right) 1_{\mathcal{Y}}(y)}{\int_{\mathcal{Y}} w \left(D_\theta^{-1}(x)(y' - C_\theta(x)) \right) dy'}.$$

It is straightforward to show that $\tilde{p}_\theta(x'|x)$ and $\tilde{q}_\theta(y|x)$ are the conditional densities of $X_{n+1}^{\theta, \lambda}$ and $Y_n^{\theta, \lambda}$ (respectively) given $X_n^{\theta, \lambda} = x$. $p_\theta(x'|x)$ and $q_\theta(y|x)$ can be interpreted as truncations of $\tilde{p}_\theta(x'|x)$ and $\tilde{q}_\theta(y|x)$ to domains \mathcal{X} and \mathcal{Y} (i.e., the hidden Markov model specified in (8), (9) can be viewed as a truncated version of the original model (6), (7)). $p_\theta(x'|x)$ and $q_\theta(y|x)$ accurately approximate $\tilde{p}_\theta(x'|x)$ and $\tilde{q}_\theta(y|x)$ when domains \mathcal{X} and \mathcal{Y} are sufficiently large (i.e., when \mathcal{X} , \mathcal{Y} contain balls of sufficiently large radius). This kind of approximation is involved (implicitly or explicitly) in any numerical implementation of the optimal filter for state-space model (6), (7) (for details see e.g., [2], [3], [6]).

The entropy rates of the hidden Markov model (8), (9) are studied under the following assumptions.

Assumption 4.1. \mathcal{X} and \mathcal{Y} are compact sets with non-empty interiors.

Assumption 4.2. $v(x) > 0$ and $w(y) > 0$ for all $x \in \mathbb{R}^{d_x}$, $y \in \mathbb{R}^{d_y}$. Moreover, $v(x)$ and $w(y)$ are real-analytic for each $x \in \mathbb{R}^{d_x}$, $y \in \mathbb{R}^{d_y}$.

Assumption 4.3. $B_\theta(x)$ and $D_\theta(x)$ are invertible for all $\theta \in \Theta$, $x \in \mathbb{R}^{d_x}$. Moreover, $A_\theta(x)$, $B_\theta(x)$, $C_\theta(x)$ and $D_\theta(x)$ are real-analytic in (θ, x) for each $\theta \in \Theta$, $x \in \mathbb{R}^{d_x}$.

Assumption 4.4. There exist a probability measure $\pi(dx)$ on \mathcal{X} and real numbers $\rho \in (0, 1)$, $K \in [1, \infty)$ such that (3) holds for all $x \in \mathcal{X}$, $n \geq 0$ and any Borel-measurable set $B \subseteq \mathcal{X}$.

Assumptions 4.1 – 4.3 are relevant for several practically important classes of non-linear state-space models. E.g., these assumptions cover stochastic volatility and dynamic probit models and their truncated versions. For other models satisfying (6), (7) and Assumptions 4.1 – 4.3, see [2], [3], [6] and references cited therein.

Using Theorems 2.1 and 2.2, we get the following results.

Corollary 4.1. Let Assumptions 4.1 – 4.4 hold. Then, all conclusions of Theorem 2.1 are true.

Corollary 4.2. Let Assumptions 4.1 – 4.3 hold. Then, all conclusions of Theorem 2.2 are true.

Corollaries 4.1 and 4.2 are proved in Section VIII.

V. RESULTS RELATED TO KERNELS OF $\{(X_n, Y_n)\}_{n \geq 0}$ AND $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$

In this section, an analytical (complex-valued) continuation of the transition kernel of $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ is constructed, and its asymptotic properties (geometric ergodicity) are studied. The same properties of the transition kernel of $\{(X_n, Y_n)\}_{n \geq 0}$ are studied, too. Throughout this and later sections, the following notations is used. Let \mathcal{W} be any Borel set in \mathbb{R}^{d_w} , where d_w is any positive integer. Then, $\mathcal{B}(\mathcal{W})$ denotes the collection of Borel sets in \mathcal{W} . $\mathcal{P}(\mathcal{W})$ is the collection of probability measures on \mathcal{W} , while $\mathcal{M}_p(\mathcal{W})$ is the set of positive measures on \mathcal{W} . $\mathcal{M}_c(\mathcal{W})$ is the collection of complex measures on \mathcal{W} , while $\mathcal{P}_c(\mathcal{W})$ is the set defined by

$$\mathcal{P}_c(\mathcal{W}) = \{\zeta \in \mathcal{M}_c(\mathcal{W}) : \zeta(\mathcal{W}) = 1\}.$$

For $\zeta \in \mathcal{M}_c(\mathcal{W})$, $\|\zeta\|$ denotes the total variation norm of ζ , while $|\zeta|(dw)$ is the total variation of $\zeta(dw)$. For $w \in \mathcal{W}$, $\delta_w(dw')$ is the Dirac measure centered at w (i.e., $\delta_w(B) = I_B(w)$ for $B \in \mathcal{B}(\mathcal{W})$).

We rely here on the following notations, too. \mathcal{Z} is the set defined by $\mathcal{Z} = \mathcal{Y} \times \mathcal{X}$. $\hat{s}_\eta(x)$ and $\tilde{r}_\eta(y, x'|x)$ are the functions defined by

$$\hat{s}_\eta(x) = \int \int \hat{r}_\eta(y', x'|x) v(dy') \mu(dx''), \quad (10)$$

$$\tilde{r}_\eta(y, x'|x) = \begin{cases} \hat{r}_\eta(y, x'|x) / \hat{s}_\eta(x), & \text{if } \hat{s}_\eta(x) \neq 0 \\ 0, & \text{otherwise} \end{cases} \quad (11)$$

for $\eta \in \mathbb{C}^d$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$. $\tilde{\psi}(z)$ is the function defined by

$$\tilde{\psi}(z) = 1 + \psi(y), \quad (12)$$

where $z = (y, x)$. $u_\eta^n(x_{0:n}, y_{1:n})$ is the function defined by

$$u_\eta^n(x_{0:n}, y_{1:n}) = \prod_{k=1}^n \tilde{r}_\eta(y_k, x_k | x_{k-1}), \quad (13)$$

where $x_0, \dots, x_n \in \mathcal{X}$, $y_1, \dots, y_n \in \mathcal{Y}$, $n \geq 1$. $\sigma(dz)$ is the measure defined by

$$\sigma(B) = \int \int I_B(y, x) Q(x, dy) \pi(dx)$$

for $B \in \mathcal{B}(\mathcal{Z})$. $S(z, dz')$, $S_\eta(z, dz')$ are the kernels defined by

$$S(z, B) = \int \int I_B(y', x') Q(x', dy') P(x, dx'), \quad (14)$$

$$S_\eta(z, B) = \int \int I_B(y', x') \tilde{r}_\eta(y', x'|x) v(dy') \mu(dx') \quad (15)$$

(as in (12), z denotes (y, z)). $\{S^n(z, dz')\}_{n \geq 0}$, $\{S_\eta^n(z, dz')\}_{n \geq 0}$ are the kernels recursively defined by $S^0(z, B) = S_\eta^0(z, B) = \delta_z(B)$ and

$$S^{n+1}(z, B) = \int S^n(z', B) S(z, dz'),$$

$$S_\eta^{n+1}(z, B) = \int S_\eta^n(z', B) S_\eta(z, dz').$$

$\{(S_\eta^n \zeta)(dz)\}_{n \geq 0}$ are the measures defined by

$$(S_\eta^n \zeta)(B) = \int S_\eta^n(z, B) \zeta(dz),$$

where $\zeta \in \mathcal{M}_c(\mathcal{Z})$.

Remark. $S(z, dz')$ and $\sigma(dz)$ are the transition kernel and the invariant distribution of $\{(X_n, Y_n)\}_{n \geq 0}$. When $\theta \in \Theta$, $S_\theta(z, dz')$ boils down to the transition kernel of $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$. Hence, for $\eta \in \mathbb{C}^d$, $S_\eta(z, dz')$ can be considered as a complex-valued continuation of the transition kernel of $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$. The kernel $S_\eta^n(z, dz')$ admits the representation

$$\begin{aligned} (S_\eta^n \zeta)(B) &= \int \cdots \int I_B(y_n, x_n) u_\eta^n(x_{0:n}, y_{1:n}) \\ &\quad \cdot (v \times \mu)(dy_n, dx_n) \cdots (v \times \mu)(dy_1, dx_1) \\ &\quad \cdot \zeta(dy_0, dx_0). \end{aligned} \quad (16)$$

This representation is used to show that $S_\eta(z, dz')$ is geometrically ergodic (see Lemma 5.4 and its proof). It is also used to show the analyticity of integral (112) (see Lemma 7.2 and its proof).

Remark. Throughout this section and later sections, the following convention is applied. Diacritic $\tilde{\cdot}$ is used to denote a locally defined quantity, i.e., a quantity whose definition holds only within the proof where the quantity appears.

Lemma 5.1. Let Assumption 2.5 hold. Then, there exists a real number $C_1 \in [1, \infty)$ such that

$$\begin{aligned} \int \tilde{\psi}(z') S(z, dz') &\leq C_1, \\ |S^n - \sigma|(z, B) &\leq C_1 \rho^n \end{aligned}$$

for all $z \in \mathcal{Z}$, $B \in \mathcal{B}(\mathcal{Z})$, $n \geq 0$ (here, $|S^n - \sigma|(z, dz')$ denotes the total variation of $S^n(z, dz') - \sigma(dz')$, while ρ is specified in Assumption 2.5).

Proof. Let $C_1 = 2K$ (K is specified in Assumption 2.5). Moreover, let x, y be any elements of \mathcal{X}, \mathcal{Y} (respectively), while $z = (y, x)$ (notice that z can be any element of \mathcal{Z}). Then, we have

$$\begin{aligned} \int \tilde{\psi}(z') S(z, dz') &= \int \int (1 + \psi(y')) Q(x', dy') P(x, dx') \\ &\leq 1 + K \leq C_1. \end{aligned}$$

We also have

$$\begin{aligned} |S^n(z, B) - \sigma(B)| &= \left| \int \int I_B(y', x') Q(x', dy') (P^n - \pi)(x, dx') \right| \\ &\leq \int \int I_B(y', x') Q(x', dy') |P^n - \pi|(x, dx') \\ &\leq 2K \rho^n \leq C_1 \rho^n \end{aligned}$$

for $B \in \mathcal{B}(\mathcal{Z})$, $n \geq 0$. \square

Lemma 5.2. Let Assumption 2.2 hold. Then, the following is true:

(i) $\tilde{r}_\theta(y, x'|x) = r_\theta(y, x'|x)$ for all $\theta \in \Theta$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$.

(ii) There exists a real number $\delta_1 \in (0, \delta]$ such that $\tilde{r}_\eta(y, x'|x)$ is analytic in η and satisfies

$$\begin{aligned} |\tilde{r}_\eta(y, x'|x)| &\leq 2|\varphi_\eta(y)|, \\ \int \int \tilde{r}_\eta(y', x''|x) v(dy') \mu(dx'') &= 1 \end{aligned}$$

for all $\eta \in V_{\delta_1}(\Theta)$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$ (δ is specified in Assumption 2.2).

Remark. As a direct consequence of Lemma 5.2 (Part (ii)), we have $S_\eta^n \zeta \in \mathcal{P}_c(\mathcal{Z})$ (i.e., $(S_\eta^n \zeta)(\mathcal{Z}) = 1$) for $\eta \in V_{\delta_1}(\Theta)$, $\zeta \in \mathcal{P}_c(\mathcal{Z})$, $n \geq 1$.

Proof. Due to Assumption 2.2, we have

$$\int \int \phi(y) v(dy) \mu(dx) = \|\mu\| \int \phi(y) v(dy) < \infty.$$

Then, using Assumption 2.2 and Lemma A1.1 (see Appendix 1), we conclude that $\hat{s}_\eta(x)$ is analytic in η for each $\eta \in V_\delta(\Theta)$, $x \in \mathcal{X}$. Relying on the same arguments, we deduce

$$|\hat{r}_{\eta'}(y, x'|x) - \hat{r}_{\eta''}(y, x'|x)| \leq \frac{d \phi(y) \|\eta' - \eta''\|}{\delta} \quad (17)$$

for $\eta', \eta'' \in V_\delta(\Theta)$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$ (here, d denotes the dimension of vectors in Θ , $V_\delta(\Theta)$).

Throughout the rest of the proof, the following notations is used. \tilde{C}, δ_1 are the real numbers defined by

$$\begin{aligned} \tilde{C} &= \frac{d \|\mu\|}{\delta} \int \phi(y) v(dy), \\ \delta_1 &= \min \left\{ \delta, \frac{1}{2\tilde{C}} \right\}. \end{aligned}$$

η, η', η'' are any elements in $V_{\delta_1}(\Theta)$, while θ is any element of Θ satisfying $\|\eta - \theta\| < \delta_1$. x, x' are any elements of \mathcal{X} , while y is any element in \mathcal{Y} .

Using (17), we conclude

$$\begin{aligned} &\left| \int \int (\hat{r}_{\eta'}(y, x'|x) - \hat{r}_{\eta''}(y, x'|x)) v(dy) \mu(dx') \right| \\ &\leq \int \int |\hat{r}_{\eta'}(y, x'|x) - \hat{r}_{\eta''}(y, x'|x)| v(dy) \mu(dx') \\ &\leq \frac{d \|\mu\| \|\eta' - \eta''\|}{\delta} \int \phi(y) v(dy) = \tilde{C} \|\eta' - \eta''\|. \end{aligned}$$

Consequently, we have

$$\begin{aligned} |\hat{s}_\eta(x)| &= \left| \int \int \hat{r}_\eta(y, x'|x) v(dy) \mu(dx') \right| \\ &\geq \int \int \hat{r}_\theta(y, x'|x) v(dy) \mu(dx') \\ &\quad - \left| \int \int (\hat{r}_\eta(y, x'|x) - \hat{r}_\theta(y, x'|x)) v(dy) \mu(dx') \right| \\ &\geq 1 - \tilde{C} \|\eta - \theta\| \geq \frac{1}{2}. \end{aligned}$$

Hence, we get

$$|\hat{s}_\eta(x)| = \left| \int \int \hat{r}_\eta(y, x'|x) \nu(dy) \mu(dx') \right| \geq \frac{1}{2}. \quad (18)$$

Therefore, we have

$$\tilde{r}_\eta(y, x'|x) = \frac{\hat{r}_\eta(y, x'|x)}{\hat{s}_\eta(x)}. \quad (19)$$

As $\hat{s}_\eta(x)$ is analytic in η for each $\eta \in V_{\delta_1}(\Theta)$, we conclude from Assumption 2.2 and (18), (19) that (i), (ii) are true. \square

Lemma 5.3. *Let Assumption 2.2 hold. Then, the following is true:*

(i) $u_\eta^n(x_{0:n}, y_{1:n})$ is analytic in η for all $\eta \in V_{\delta_1}(\Theta)$, $x_0, \dots, x_n \in \mathcal{X}$, $y_1, \dots, y_n \in \mathcal{Y}$, $n \geq 1$ (δ_1 is specified in Lemma 5.2).

(ii) There exists a non-decreasing sequence $\{K_n\}_{n \geq 1}$ in $[1, \infty)$ such that

$$\begin{aligned} |u_\eta^n(x_{0:n}, y_{1:n})| &\leq K_n \left(\prod_{k=1}^n \phi(y_k) \right), \\ |u_{\eta'}^n(x_{0:n}, y_{1:n}) - u_{\eta''}^n(x_{0:n}, y_{1:n})| \\ &\leq K_n \|\eta' - \eta''\| \left(\prod_{k=1}^n \phi(y_k) \right) \end{aligned}$$

for all $\eta, \eta', \eta'' \in V_{\delta_1}(\Theta)$, $x_0, \dots, x_n \in \mathcal{X}$, $y_1, \dots, y_n \in \mathcal{Y}$, $n \geq 1$.

Proof. Throughout the proof, the following notations is used. $\{K_n\}_{n \geq 1}$ are the real numbers defined by $K_n = 2^n d / \delta_1$ for $n \geq 1$ (here, d denotes the dimension of vectors in Θ , $V_\delta(\Theta)$). η, η', η'' are any elements of $V_{\delta_1}(\Theta)$. $\{x_n\}_{n \geq 0}$, $\{y_n\}_{n \geq 1}$ are any sequences in \mathcal{X} , \mathcal{Y} (respectively).

Owing to Lemma 5.2, $u_\eta^n(x_{0:n}, y_{1:n})$ is analytic in η for each $\eta \in V_{\delta_1}(\Theta)$. Due to Assumption 2.2 and the same lemma, we have

$$|u_\eta^n(x_{0:n}, y_{1:n})| \leq 2^n \left(\prod_{k=1}^n |\phi_\eta(y_k)| \right) \leq K_n \left(\prod_{k=1}^n \phi(y_k) \right)$$

for $n \geq 1$. Consequently, Lemma A1.1 (see Appendix 1) yields

$$\begin{aligned} &|u_{\eta'}^n(x_{0:n}, y_{1:n}) - u_{\eta''}^n(x_{0:n}, y_{1:n})| \\ &\leq \frac{2^n d \|\eta' - \eta''\|}{\delta_1} \left(\prod_{k=1}^n \phi(y_k) \right) = K_n \|\eta' - \eta''\| \left(\prod_{k=1}^n \phi(y_k) \right) \end{aligned}$$

for $n \geq 1$. \square

Lemma 5.4. *Let Assumptions 2.1, 2.2 and 2.4 hold. Then, the following is true:*

(i) There exist real numbers $\delta_2 \in (0, \delta_1]$, $C_2 \in [1, \infty)$ such that

$$\begin{aligned} |S_{\eta'} - S_{\eta''}|(z, B) &\leq C_2 \|\eta' - \eta''\|, \\ \int \tilde{\psi}(z') |S_\eta|(z, dz') &\leq C_2 \end{aligned}$$

for all $\eta, \eta', \eta'' \in V_{\delta_2}(\Theta)$, $z \in \mathcal{Z}$, $B \in \mathcal{B}(\mathcal{Z})$ (here, $|S_{\eta'} - S_{\eta''}|(z, dz')$ denotes the total variation of $S_{\eta'}(z, dz') - S_{\eta''}(z, dz')$, while δ_1 is specified in Lemma 5.2).

(ii) For each $\eta \in V_{\delta_2}(\Theta)$, there exists a complex measure $\sigma_\eta(dz)$ on \mathcal{Z} such that $\sigma_\eta(B) = \lim_{n \rightarrow \infty} S_\eta^n(z, B)$ for all $z \in \mathcal{Z}$, $B \in \mathcal{B}(\mathcal{Z})$.

(iii) There exists a real number $\gamma_1 \in (0, 1)$, such that

$$|S_\eta^n - \sigma_\eta|(z, B) \leq C_2 \gamma_1^n$$

for all $\eta \in V_{\delta_2}(\Theta)$, $z \in \mathcal{Z}$, $B \in \mathcal{B}(\mathcal{Z})$, $n \geq 0$ (here, $|S_\eta^n - \sigma_\eta|(z, dz')$ stands for the total variation of $S_\eta^n(z, dz') - \sigma_\eta(dz')$).

Proof. Throughout the proof, the following notations is used. \tilde{C}_1, \tilde{C}_2 are the real numbers defined by

$$\tilde{C}_1 = 2\|\mu\| \int \psi(y) \phi(y) \nu(dy),$$

$$\tilde{C}_2 = 2\|\mu\| \int \phi(y) \nu(dy),$$

while n_0 is the integer defined as

$$n_0 = \left\lceil \frac{\log 4}{|\log(1 - \varepsilon^2)|} \right\rceil$$

($\varepsilon, \phi(y), \psi(y)$, are specified in Assumptions 2.1, 2.3). $\{\tilde{K}_n\}_{n \geq 1}$ are the real numbers defined by $\tilde{K}_n = (1 + \tilde{C}_2)^n K_n$ for $n \geq 1$, while $\delta_2, \gamma_1, \tilde{C}_3, C_2$ are the real numbers defined as $\delta_2 = \delta_1 / (4\tilde{K}_{n_0})$, $\gamma_1 = 2^{-1/n_0}$, $\tilde{C}_3 = \tilde{K}_1 + \tilde{C}_1 + \tilde{C}_2$, $C_2 = 16\tilde{C}_3 \gamma_1^{-n_0}$ (δ_1, K_n are specified in Lemmas 5.2, 5.3). η, η', η'' are any elements in $V_{\delta_2}(\Theta)$, while θ is any element of Θ satisfying $\|\eta - \theta\| < \delta_2$. x, y are any elements of \mathcal{X} , \mathcal{Y} (respectively), while $z = (y, x)$. ζ, ζ', ζ'' are any elements of $\mathcal{P}_c(\mathcal{Z})$, while B is any element of $\mathcal{B}(\mathcal{Z})$. $n \geq 1, k \geq 0$ are any integers.

Relying on Assumption 2.4 and Lemma 5.2, we deduce

$$\begin{aligned} \int \tilde{\psi}(z') |S_\eta|(z, dz') &\leq \int \int (1 + \psi(y')) |\tilde{r}_\eta(y', x'|x)| \\ &\quad \cdot \nu(dy') \mu(dx') \\ &\leq 2\|\mu\| \int (1 + \psi(y')) \phi(y') \nu(dy') \\ &= \tilde{C}_1 + \tilde{C}_2 \leq C_2 \end{aligned}$$

as $\tilde{C}_1 + \tilde{C}_2 \leq \tilde{C}_3 \leq C_2$. Moreover, using Lemma 5.3, we conclude

$$\begin{aligned} &|(S_{\eta'}^n \zeta)(B) - (S_{\eta''}^n \zeta)(B)| \\ &\leq \int \cdots \int \int I_B(y_n, x_n) |u_{\eta'}^n(x_{0:n}, y_{1:n}) - u_{\eta''}^n(x_{0:n}, y_{1:n})| \\ &\quad \cdot (\nu \times \mu)(dy_n, dx_n) \cdots (\nu \times \mu)(dy_1, dx_1) |\zeta|(dy_0, dx_0) \\ &\leq K_n \|\mu\|^n \|\zeta\| \|\eta' - \eta''\| \left(\prod_{k=1}^n \int \phi(y_k) \nu(dy_k) \right) \\ &\leq \tilde{K}_n \|\zeta\| \|\eta' - \eta''\|. \end{aligned}$$

Therefore, we get

$$\left\| S_{\eta'}^n \zeta - S_{\eta''}^n \zeta \right\| \leq \tilde{K}_n \|\zeta\| \|\eta' - \eta''\|. \quad (20)$$

Hence, we have

$$\begin{aligned} |S_{\eta'} - S_{\eta''}|(z, B) &= |S_{\eta'} \delta_z - S_{\eta''} \delta_z|(B) \\ &\leq \tilde{K}_1 \|\delta_z\| \|\eta' - \eta''\| \\ &\leq C_2 \|\eta' - \eta''\| \end{aligned}$$

as $\tilde{K}_1 \leq \tilde{C}_3 \leq C_2$.

Let $\tau_\theta(dz)$ be the measure defined by

$$\tau_\theta(B) = \int \int I_B(y, x) \lambda_\theta(dx|y) \nu(dy).$$

Owing to Assumption 2.1, we have

$$1 = \int \int r_\theta(y, x'|x) \nu(dy) \mu(dx') \leq \frac{1}{\varepsilon} \int \lambda_\theta(\mathcal{X}|y) \nu(dy).$$

Hence, we get

$$\tau_\theta(\mathcal{Z}) = \int \lambda_\theta(\mathcal{X}|y) \nu(dy) \geq \varepsilon.$$

Moreover, due to Assumption 2.1 and Lemma 5.2, we have

$$\begin{aligned} S_\theta(z, B) &= \int \int I_B(y', x') r_\theta(y', x'|x) \nu(dy') \mu(dx') \\ &\geq \varepsilon \int \int I_B(y', x') \lambda_\theta(dx'|y') \nu(dy') = \varepsilon \tau_\theta(B). \end{aligned}$$

Then, standard results in Markov chain theory (see e.g., [18, Theorem 16.0.2]) imply that there exists a probability measure $\sigma_\theta(dz)$ on \mathcal{Z} such that

$$|S_\theta^n(z, B) - \sigma_\theta(B)| \leq (1 - \varepsilon \tau_\theta(\mathcal{Z}))^n \leq (1 - \varepsilon^2)^n.$$

As $\sigma_\theta(B)(\zeta'(\mathcal{Z}) - \zeta''(\mathcal{Z})) = 0$, we get

$$\begin{aligned} &|(S_\theta^n \zeta')(B) - (S_\theta^n \zeta'')(B)| \\ &= \left| \int (S_\theta^n - \sigma_\theta)(z, B)(\zeta' - \zeta'')(dz) \right| \\ &\leq \int |S_\theta^n - \sigma_\theta|(z, B) |\zeta' - \zeta''|(dz) \\ &\leq (1 - \varepsilon^2)^n \|\zeta' - \zeta''\|. \end{aligned}$$

Hence, we have

$$\|S_\theta^n \zeta' - S_\theta^n \zeta''\| \leq (1 - \varepsilon^2)^n \|\zeta' - \zeta''\|. \quad (21)$$

Since $S_\theta^n(z, dz')$ is an element of $\mathcal{P}(\mathcal{Z})$, we conclude $\|S_\theta^n \zeta\| \leq \|\zeta\|$. Then, owing to (20), we have

$$\begin{aligned} \|S_\eta^n \zeta\| &\leq \|S_\theta^n \zeta\| + \|(S_\eta^n - S_\theta^n) \zeta\| \\ &\leq (1 + \tilde{K}_n \|\eta - \theta\|) \|\zeta\| \\ &\leq (1 + \tilde{K}_n \delta_2) \|\zeta\| \leq 2 \|\zeta\| \end{aligned} \quad (22)$$

when $n \leq n_0$, as $\|\eta - \theta\| < \delta_2$, $\tilde{K}_n \delta_2 \leq \tilde{K}_{n_0} \delta_2 = \delta_1/4 \leq 1/4$. Moreover, due to (20), (21), we have

$$\begin{aligned} \|S_\eta^n \zeta' - S_\eta^n \zeta''\| &\leq \|S_\theta^n \zeta' - S_\theta^n \zeta''\| + \|(S_\eta^n - S_\theta^n)(\zeta' - \zeta'')\| \\ &\leq \left((1 - \varepsilon^2)^n + \tilde{K}_n \|\eta - \theta\| \right) \|\zeta' - \zeta''\| \end{aligned}$$

$$\leq \left((1 - \varepsilon^2)^n + \frac{1}{4} \right) \|\zeta' - \zeta''\|$$

when $n \leq n_0$. Setting $n = n_0$, we conclude

$$\|S_\eta^{n_0} \zeta' - S_\eta^{n_0} \zeta''\| \leq \frac{\|\zeta' - \zeta''\|}{2}$$

as $(1 - \varepsilon^2)^{n_0} \leq 1/4$. Since $S_\eta^n \zeta \in \mathcal{P}_c(\mathcal{Z})$ (see Lemma 5.2 and the remark immediately after its statement), we have

$$\begin{aligned} \|S_\eta^{(k+1)n_0}(\zeta' - \zeta'')\| &= \|S_\eta^{n_0} (S_\eta^{kn_0} \zeta' - S_\eta^{kn_0} \zeta'')\| \\ &\leq \frac{1}{2} \|S_\eta^{kn_0}(\zeta' - \zeta'')\|. \end{aligned} \quad (23)$$

Iterating (23), we get

$$\|S_\eta^{kn_0}(\zeta' - \zeta'')\| \leq \frac{1}{2^k} \|\zeta' - \zeta''\|. \quad (24)$$

Using (22), (24), we conclude

$$\begin{aligned} \|S_\eta^{(k+1)n_0} \zeta - S_\eta^{kn_0} \zeta\| &= \|S_\eta^{kn_0} (S_\eta^{n_0} \zeta - \zeta)\| \\ &\leq \frac{1}{2^k} \|S_\eta^{n_0} \zeta - \zeta\| \\ &\leq \frac{1}{2^k} (\|S_\eta^{n_0} \zeta\| + \|\zeta\|) \\ &\leq \frac{\|\zeta\|}{2^{k-2}}. \end{aligned} \quad (25)$$

Hence, we get

$$\sum_{k=0}^{\infty} \|S_\eta^{(k+1)n_0} \zeta - S_\eta^{kn_0} \zeta\| \leq \sum_{k=0}^{\infty} \frac{\|\zeta\|}{2^{k-2}} = 8 \|\zeta\| < \infty. \quad (26)$$

Let $(S_\eta^\infty \zeta)(dz)$ be the measure defined by

$$(S_\eta^\infty \zeta)(B) = \zeta(B) + \sum_{k=0}^{\infty} \left((S_\eta^{(k+1)n_0} \zeta)(B) - (S_\eta^{kn_0} \zeta)(B) \right).$$

Then, due to (26), $(S_\eta^\infty \zeta)(dz)$ is well-defined and satisfies $S_\eta^\infty \zeta \in \mathcal{P}_c(\mathcal{Z})$. Moreover, owing to (25), (26), we have

$$\begin{aligned} \|S_\eta^{kn_0} \zeta - S_\eta^\infty \zeta\| &= \left\| \sum_{j=k}^{\infty} (S_\eta^{(j+1)n_0} \zeta - S_\eta^{jn_0} \zeta) \right\| \\ &\leq \sum_{j=k}^{\infty} \|S_\eta^{(j+1)n_0} \zeta - S_\eta^{jn_0} \zeta\| \\ &\leq \sum_{j=k}^{\infty} \frac{\|\zeta\|}{2^{j-2}} = \frac{\|\zeta\|}{2^{k-3}}. \end{aligned} \quad (27)$$

Combining this with (24), we get

$$\begin{aligned} \|S_\eta^\infty \zeta' - S_\eta^\infty \zeta''\| &\leq \|S_\eta^{kn_0} \zeta' - S_\eta^\infty \zeta'\| + \|S_\eta^{kn_0} \zeta'' - S_\eta^\infty \zeta''\| \\ &\quad + \|S_\eta^{kn_0} \zeta' - S_\eta^{kn_0} \zeta''\| \\ &\leq \frac{\|\zeta'\| + \|\zeta''\| + \|\zeta' - \zeta''\|}{2^{k-3}}. \end{aligned}$$

Therefore, $S_\eta^\infty \zeta' = S_\eta^\infty \zeta''$ for any $\zeta', \zeta'' \in \mathcal{P}_c(\mathcal{Z})$. Consequently, there exists $\sigma_\eta \in \mathcal{P}_c(\mathcal{Z})$ such that $S_\eta^\infty \zeta = \sigma_\eta$ for any

$\zeta \in \mathcal{P}_c(\mathcal{Z})$. Hence, $S_\eta^\infty(S_\eta^n \zeta) = \sigma_\eta$, as $S_\eta^n \zeta \in \mathcal{P}_c(\mathcal{Z})$. Then, (22), (27) imply

$$\begin{aligned} \left\| S_\eta^n \zeta - \sigma_\eta \right\| &= \left\| S_\eta^{kn_0}(S_\eta^{n-kn_0} \zeta) - S_\eta^\infty(S_\eta^{n-kn_0} \zeta) \right\| \\ &\leq \frac{1}{2^{k-3}} \left\| S_\eta^{n-kn_0} \zeta \right\| \\ &\leq \frac{\|\zeta\|}{2^{k-4}} \leq C_2 \gamma_1^n \|\zeta\| \end{aligned} \quad (28)$$

when $(k+1)n_0 \geq n > kn_0$, as $2^{-(k-4)} = 16\gamma_1^{kn_0} \leq (16\gamma_1^{-n_0})\gamma_1^n \leq C_2\gamma_1^n$. Thus, we get

$$\begin{aligned} \left| S_\eta^n - \sigma_\eta \right|(z, B) &= \left| S_\eta^n \delta_z - \sigma_\eta \right|(B) \\ &\leq \left\| S_\eta^n \delta_z - \sigma_\eta \right\| \\ &\leq C_2 \gamma_1^n \|\delta_z\| = C_2 \gamma_1^n \end{aligned}$$

by setting $k = \lfloor (n-m)/n_0 \rfloor$ in (28).

VI. RESULTS RELATED TO OPTIMAL FILTER

In this section, an analytic (complex-valued) continuation of the optimal filter is constructed, and its asymptotic properties (exponential forgetting) are studied. Here, we rely on the following notations. $\mathcal{B}(\mathcal{X})$, $\mathcal{P}(\mathcal{X})$, $\mathcal{M}_p(\mathcal{X})$ and $\mathcal{M}_c(\mathcal{X})$ have been defined at the beginning of Section V. For $x \in \mathcal{X}$, $\xi \in \mathcal{M}_c(\mathcal{X})$, $\|\xi\|$, $|\xi|(\mathrm{d}x')$ and $\delta_x(\mathrm{d}x')$ are the norm and measures specified at the beginning of Section V. For $\gamma \in (0, 1)$, $V_\gamma(\mathcal{P}(\mathcal{X}))$ is the open γ -vicinity of $\mathcal{P}(\mathcal{X})$, i.e.,

$$V_\gamma(\mathcal{P}(\mathcal{X})) = \{\zeta \in \mathcal{M}_c(\mathcal{X}) : \exists \lambda \in \mathcal{P}(\mathcal{X}), \|\zeta - \lambda\| < \gamma\}.$$

$R_{\eta,y}(\mathrm{d}x|\zeta)$ is the measure defined by

$$R_{\eta,y}(B|\zeta) = \int \int I_B(x') \tilde{r}_\eta(y, x'|x) \mu(\mathrm{d}x') \zeta(\mathrm{d}x) \quad (29)$$

for $\eta \in \mathbb{C}^d$, $\zeta \in \mathcal{M}_c(\mathcal{X})$, $B \in \mathcal{B}(\mathcal{X})$, $y \in \mathcal{Y}$ ($\tilde{r}_\eta(y, x'|x)$ is specified in Lemma 5.2). $\Phi_{\eta,y}(\zeta)$ is the function defined by

$$\Phi_{\eta,y}(\zeta) = \begin{cases} \log R_{\eta,y}(\mathcal{X}|\zeta), & \text{if } R_{\eta,y}(\mathcal{X}|\zeta) \neq 0 \\ 0, & \text{otherwise} \end{cases} \quad (30)$$

$v_{\eta,y}^{m:n}(x_{m:n})$ and $\varphi_{\eta,y}^{m:n}$ are the functions defined by

$$v_{\eta,y}^{m:n}(x_{m:n}) = \prod_{k=m+1}^n \tilde{r}_\eta(y_k, x_k|x_{k-1}), \quad (31)$$

$$\varphi_{\eta,y}^{m:n} = \prod_{k=m+1}^n \varphi_\eta(y_k), \quad (32)$$

where $x_m, \dots, x_n \in \mathcal{X}$, $n > m \geq 0$ and $\mathbf{y} = \{y_n\}_{n \geq 1}$ is any sequence in \mathcal{Y} . $r_{\eta,y}^{m:n}(x'|x)$ is the function defined by

$$\begin{aligned} r_{\eta,y}^{m:n}(x'|x) &= \int \int \dots \int \int v_{\eta,y}^{m:n}(x_{m:n}) \\ &\cdot \delta_{x'}(\mathrm{d}x_n) \mu(\mathrm{d}x_{n-1}) \dots \mu(\mathrm{d}x_{m+1}) \delta_x(\mathrm{d}x_m), \end{aligned} \quad (33)$$

where $x, x' \in \mathcal{X}$. $R_{\eta,y}^{m:n}(\mathrm{d}x|\zeta)$ and $R_{\eta,y}^{m:n}(\mathrm{d}x|\zeta)$ are the measures defined by $R_{\eta,y}^{m:n}(B|\zeta) = \zeta(B)$ and

$$R_{\eta,y}^{m:n}(B|\zeta) = \int \int I_B(x') r_{\eta,y}^{m:n}(x'|x) \mu(\mathrm{d}x') \zeta(\mathrm{d}x). \quad (34)$$

$f_{\eta,y}^{m:n}(x|\zeta)$, $g_{\eta,y}^{m:n}(x'|x, \zeta)$, $h_{\eta,y}^{m:n}(x|x', \zeta)$ are the functions defined by

$$g_{\eta,y}^{m:n}(x'|x, \zeta) = \begin{cases} r_{\eta,y}^{m:n}(x'|x) / R_{\eta,y}^{m:n}(\mathcal{X}|\zeta), & \text{if } R_{\eta,y}^{m:n}(\mathcal{X}|\zeta) \neq 0 \\ 0, & \text{otherwise} \end{cases}, \quad (35)$$

$$f_{\eta,y}^{m:n}(x|\zeta) = \int g_{\eta,y}^{m:n}(x|x'', \zeta) \xi(\mathrm{d}x''), \quad (36)$$

$$\begin{aligned} h_{\eta,y}^{m:n}(x'|x, \zeta) &= -f_{\eta,y}^{m:n}(x'|\zeta) \int g_{\eta,y}^{m:n}(x''|x, \zeta) \mu(\mathrm{d}x'') \\ &\quad + g_{\eta,y}^{m:n}(x'|x, \zeta). \end{aligned} \quad (37)$$

$F_{\eta,y}^{m:n}(\mathrm{d}x|\zeta)$ and $F_{\eta,y}^{m:n}(\mathrm{d}x|\zeta)$ are the measures defined by $F_{\eta,y}^{m:n}(B|\zeta) = \zeta(B)$ and

$$F_{\eta,y}^{m:n}(B|\zeta) = \int I_B(x) f_{\eta,y}^{m:n}(x|\zeta) \mu(\mathrm{d}x). \quad (38)$$

Throughout this and later sections, measures $R_{\eta,y}^{m:n}(\mathrm{d}x|\zeta)$, $F_{\eta,y}^{m:n}(\mathrm{d}x|\zeta)$ are also denoted by $R_{\eta,y}^{m:n}(\zeta)$, $F_{\eta,y}^{m:n}(\zeta)$ (short-hand notations), while $\langle R_{\eta,y}^{m:n}(\zeta) \rangle$, $\langle F_{\eta,y}^{m:n}(\zeta) \rangle$ are defined by

$$\langle R_{\eta,y}^{m:n}(\zeta) \rangle = R_{\eta,y}^{m:n}(\mathcal{X}|\zeta), \quad (39)$$

$$\langle F_{\eta,y}^{m:n}(\zeta) \rangle = F_{\eta,y}^{m:n}(\mathcal{X}|\zeta).$$

Remark. When $\theta \in \Theta$, $\lambda \in \mathcal{P}(\mathcal{X})$, $F_{\theta,y}^{m:n}(\lambda)$ is the optimal filter for the model $\{(X_n^{\theta,\lambda}, Y_n^{\theta,\lambda})\}_{n \geq 0}$, i.e.,

$$F_{\theta,y}^{1:n}(B|\lambda) = P\left(X_n^{\theta,\lambda} \in B \mid Y_{1:n}^{\theta,\lambda} = y_{1:n}\right).$$

Hence, for $\eta \in \mathbb{C}^d$, $\zeta \in \mathcal{M}_c(\mathcal{X})$, $F_{\theta,y}^{m:n}(\zeta)$ can be considered as a complex-valued continuation of the optimal filter. Consequently, $f_{\theta,y}^{m:n}(x|\zeta)$ can be viewed as a complex-valued continuation of the optimal filtering density. $h_{\theta,y}^{m:n}(x'|x, \zeta)$ can be described as the Gateaux derivative of $f_{\theta,y}^{m:n}(x|\zeta)$ with respect to ζ (see (75) – (77)). $h_{\theta,y}^{m:n}(x'|x, \zeta)$ is used to show that $F_{\eta,y}^{m:n}(\zeta)$ forgets initial condition ξ at an exponential rate (see Lemmas 6.5, 6.6 and their proofs).

Lemma 6.1. Let η, ζ be any elements of \mathbb{C}^d , $\mathcal{M}_c(\mathcal{X})$ (respectively), while $\mathbf{y} = \{y_n\}_{n \geq 1}$ is any sequence in \mathcal{Y} . Moreover, let n, m, k be any integers satisfying $n \geq k \geq m$. Then, the following is true:

- (i) $R_{\eta,y}^{m:n}(\zeta) = R_{\eta,y}^{k:n}(R_{\eta,y}^{m:k}(\zeta))$.
- (ii) $\langle R_{\eta,y}^{m:n}(\zeta) \rangle = \langle R_{\eta,y}^{k:n}(F_{\eta,y}^{m:k}(\zeta)) \rangle \langle R_{\eta,y}^{m:k}(\zeta) \rangle$ when $\langle R_{\eta,y}^{m:k}(\zeta) \rangle \neq 0$.
- (iii) $F_{\eta,y}^{m:n}(\zeta) = F_{\eta,y}^{k:n}(F_{\eta,y}^{m:k}(\zeta))$ when $\langle R_{\eta,y}^{m:k}(\zeta) \rangle \neq 0$ and $\langle R_{\eta,y}^{m:n}(\zeta) \rangle \neq 0$.

Proof. (i) When $k = m$ or $k = n$, (i) is trivially satisfied. In what follows in this part of the proof, we assume $n > k > m$.

Owing to (29), we have

$$v_{\eta,y}^{m:n}(x_{m:n}) = v_{\eta,y}^{k:n}(x_{k:n}) v_{\eta,y}^{m:k}(x_{m:k})$$

for $x_m, \dots, x_n \in \mathcal{X}$. Combining this with (33), it is easy to show

$$r_{\eta,y}^{m:n}(x'|x) = \int r_{\eta,y}^{k:n}(x'|x'') r_{\eta,y}^{m:k}(x''|x) \mu(\mathrm{d}x'')$$

for $x, x' \in \mathcal{X}$. Then, using (34), we conclude

$$\begin{aligned} R_{\eta,y}^{m:n}(B|\xi) &= \int \int \int I_B(x') r_{\eta,y}^{k:n}(x'|x'') r_{\eta,y}^{m:k}(x''|x) \\ &\quad \cdot \mu(dx'') \mu(dx') \xi(dx) \\ &= \int \int I_B(x') r_{\eta,y}^{k:n}(x'|x'') R_{\eta,y}^{m:k}(dx''|\xi) \mu(dx') \\ &= R_{\eta,y}^{k:n}(B|R_{\eta,y}^{m:k}(\xi)) \end{aligned}$$

for $B \in \mathcal{B}(\mathcal{X})$. Hence, (i) holds when $n > k > m$.

(ii) We assume $\langle R_{\eta,y}^{m:k}(\xi) \rangle \neq 0$ (i.e., $R_{\eta,y}^{m:k}(\mathcal{X}|\xi) \neq 0$). Then, using (35), (36), (38), we conclude

$$F_{\eta,y}^{m:k}(\xi) = \frac{R_{\eta,y}^{m:k}(\xi)}{\langle R_{\eta,y}^{m:k}(\xi) \rangle}. \quad (40)$$

Since $\langle R_{\eta,y}^{k:n}(\xi) \rangle$ is linear in ξ , we deduce

$$\left\langle R_{\eta,y}^{k:n}(F_{\eta,y}^{m:k}(\xi)) \right\rangle = \frac{\left\langle R_{\eta,y}^{k:n}(R_{\eta,y}^{m:k}(\xi)) \right\rangle}{\left\langle R_{\eta,y}^{m:k}(\xi) \right\rangle}.$$

Combining this with (i), we get

$$\begin{aligned} \left\langle R_{\eta,y}^{m:n}(\xi) \right\rangle &= \left\langle R_{\eta,y}^{k:n}(R_{\eta,y}^{m:k}(\xi)) \right\rangle \\ &= \left\langle R_{\eta,y}^{k:n}(F_{\eta,y}^{m:k}(\xi)) \right\rangle \left\langle R_{\eta,y}^{m:k}(\xi) \right\rangle. \end{aligned} \quad (41)$$

Thus, (ii) is true.

(iii) We assume $\langle R_{\eta,y}^{m:k}(\xi) \rangle \neq 0$, $\langle R_{\eta,y}^{m:n}(\xi) \rangle \neq 0$. Therefore, (ii) implies $\langle R_{\eta,y}^{k:n}(F_{\eta,y}^{m:k}(\xi)) \rangle \neq 0$. Then, using the same arguments as in (ii), we deduce

$$\begin{aligned} F_{\eta,y}^{m:n}(\xi) &= \frac{R_{\eta,y}^{m:n}(\xi)}{\langle R_{\eta,y}^{m:n}(\xi) \rangle}, \\ F_{\eta,y}^{k:n}(F_{\eta,y}^{m:k}(\xi)) &= \frac{R_{\eta,y}^{k:n}(F_{\eta,y}^{m:k}(\xi))}{\langle R_{\eta,y}^{k:n}(F_{\eta,y}^{m:k}(\xi)) \rangle}. \end{aligned}$$

Combining this with (i) and (40), (41), we get

$$\begin{aligned} F_{\eta,y}^{m:n}(\xi) &= \frac{R_{\eta,y}^{k:n}(R_{\eta,y}^{m:k}(\xi))}{\langle R_{\eta,y}^{k:n}(F_{\eta,y}^{m:k}(\xi)) \rangle \langle R_{\eta,y}^{m:k}(\xi) \rangle} \\ &= \frac{R_{\eta,y}^{k:n}(F_{\eta,y}^{m:k}(\xi))}{\langle R_{\eta,y}^{k:n}(F_{\eta,y}^{m:k}(\xi)) \rangle} = F_{\eta,y}^{k:n}(F_{\eta,y}^{m:k}(\xi)) \end{aligned}$$

by using again the fact that $R_{\eta,y}^{k:n}(\xi)$ is linear in ξ . Hence, (iii) holds. \square

Lemma 6.2. *Let Assumption 2.1 hold. Then, there exist real numbers $\delta_3 \in (0, \delta_1]$, $\gamma_2 \in (0, 1)$, $C_3 \in [1, \infty)$ such that*

$$\left\| F_{\theta,y}^{m:n}(\lambda') - F_{\theta,y}^{m:n}(\lambda'') \right\| \leq C_3 \gamma_2^{n-m} \|\lambda' - \lambda''\|$$

for all $\theta \in \Theta$, $\lambda', \lambda'' \in V_{\delta_3}(\mathcal{P}(\mathcal{X})) \cap \mathcal{M}_p(\mathcal{X})$, $n \geq m \geq 0$ and any sequence $\mathbf{y} = \{y_n\}_{n \geq 1}$ in \mathcal{Y} (δ_1 is specified in Lemma 5.2).

Proof. Due to [17, Proposition 4.1, Corollary 4.2] (or [24, Theorem 3.1]) and Lemma 5.2, there exist real numbers $\gamma_2 \in (0, 1)$, $C_3 \in [1, \infty)$ such that

$$\left\| F_{\theta,y}^{m:n}(\lambda') - F_{\theta,y}^{m:n}(\lambda'') \right\| \leq \frac{C_3 \gamma_2^{n-m}}{4} \left\| \frac{\lambda'}{\|\lambda'\|} - \frac{\lambda''}{\|\lambda''\|} \right\| \quad (42)$$

for all $\theta \in \Theta$, $\lambda', \lambda'' \in \mathcal{M}_p(\mathcal{X})$, $n \geq m \geq 0$ and any sequence $\mathbf{y} = \{y_n\}_{n \geq 1}$ in \mathcal{Y} . We have used here the identity $F_{\theta,y}^{m:n}(\lambda) = F_{\theta,y}^{m:n}(\lambda/\|\lambda\|)$ for $\lambda \in \mathcal{M}_p(\mathcal{X})$.

Let $\delta_3 = \min\{1/2, \delta_1\}$, while $\mathbf{y} = \{y_n\}_{n \geq 1}$ is any sequence in \mathcal{Y} . Then, we have $\|\lambda\| \geq 1 - \delta_3 \geq 1/2$ for $\lambda \in V_{\delta_3}(\mathcal{P}(\mathcal{X})) \cap \mathcal{M}_p(\mathcal{X})$. Consequently, (42) implies

$$\begin{aligned} &\left\| F_{\theta,y}^{m:n}(\lambda') - F_{\theta,y}^{m:n}(\lambda'') \right\| \\ &\leq \frac{C_3 \gamma_2^{n-m}}{4} \left\| \frac{\lambda' - \lambda''}{\|\lambda'\|} - \frac{\lambda''(\|\lambda'\| - \|\lambda''\|)}{\|\lambda'\| \|\lambda''\|} \right\| \\ &\leq \frac{C_3 \gamma_2^{n-m} \|\lambda' - \lambda''\|}{2 \|\lambda'\|} \\ &\leq C_3 \gamma_2^{n-m} \|\lambda' - \lambda''\| \end{aligned}$$

for $\theta \in \Theta$, $\lambda', \lambda'' \in V_{\delta_3}(\mathcal{P}(\mathcal{X})) \cap \mathcal{M}_p(\mathcal{X})$, $n \geq m \geq 0$, as $2\|\lambda'\| \geq 1$, $\|\lambda''/\|\lambda''\|\| = 1$, $\|\|\lambda'\| - \|\lambda''\|\| \leq \|\lambda' - \lambda''\|$. \square

Lemma 6.3. *Let Assumptions 2.2 and 2.3 hold. Then, the following is true:*

(i) $\langle R_{\eta,y}^{m:n}(\xi) \rangle$ is analytic in η for all $\eta \in V_{\delta_1}(\Theta)$, $\xi \in V_{\delta_1}(\mathcal{P}(\mathcal{X}))$, $n \geq m \geq 0$ and any sequence $\mathbf{y} = \{y_n\}_{n \geq 1}$ in \mathcal{Y} (δ_1 is specified in Lemma 5.2).

(ii) There exists a non-decreasing sequence $\{L_n\}_{n \geq 1}$ in $[1, \infty)$ such that

$$\begin{aligned} &\left| \frac{\langle R_{\eta,y}^{m:n}(\xi) \rangle}{\varphi_{\eta,y}^{m:n}} \right| \leq L_{n-m}, \\ &\left| \frac{\langle R_{\eta',y}^{m:n}(\xi') \rangle}{\varphi_{\eta',y}^{m:n}} - \frac{\langle R_{\eta'',y}^{m:n}(\xi'') \rangle}{\varphi_{\eta'',y}^{m:n}} \right| \\ &\leq L_{n-m} (\|\eta' - \eta''\| + \|\xi' - \xi''\|) \end{aligned}$$

for all $\eta, \eta', \eta'' \in V_{\delta_1}(\Theta)$, $\xi, \xi', \xi'' \in V_{\delta_1}(\mathcal{P}(\mathcal{X}))$, $n > m \geq 0$ and any sequence $\mathbf{y} = \{y_n\}_{n \geq 1}$ in \mathcal{Y} .

(iii) There exists a non-increasing sequence $\{a_n\}_{n \geq 1}$ in $(0, \delta_1]$ such that

$$\frac{\operatorname{Re} \left\{ \langle R_{\eta,y}^{m:n}(\xi) \rangle \right\}}{|\varphi_{\theta,y}^{m:n}|} \geq \frac{1}{L_{n-m}}$$

for all $\eta \in V_{a_{n-m}}(\Theta)$, $\xi \in V_{a_{n-m}}(\mathcal{P}(\mathcal{X}))$, $n > m \geq 0$ and any sequence $\mathbf{y} = \{y_n\}_{n \geq 1}$ in \mathcal{Y} .

(iv) There exists a non-decreasing sequence $\{M_n\}_{n \geq 1}$ in $[1, \infty)$ such that

$$\begin{aligned} \max & \left\{ \left| f_{\eta, \mathbf{y}}^{m:n}(x|\xi) \right|, \left| h_{\eta, \mathbf{y}}^{m:n}(x'|x, \xi) \right| \right\} \leq M_{n-m}, \\ & \left| f_{\eta', \mathbf{y}}^{m:n}(x|\xi') - f_{\eta'', \mathbf{y}}^{m:n}(x|\xi'') \right| \\ & \leq M_{n-m} (\|\eta' - \eta''\| + \|\xi' - \xi''\|), \\ & \left| h_{\eta', \mathbf{y}}^{m:n}(x'|x, \xi') - h_{\eta'', \mathbf{y}}^{m:n}(x'|x, \xi'') \right| \\ & \leq M_{n-m} (\|\eta' - \eta''\| + \|\xi' - \xi''\|) \end{aligned}$$

for all $\eta, \eta', \eta'' \in V_{\alpha_{n-m}}(\Theta)$, $\xi, \xi', \xi'' \in V_{\alpha_{n-m}}(\mathcal{P}(\mathcal{X}))$, $x, x' \in \mathcal{X}$, $n > m \geq 0$ and any sequence $\mathbf{y} = \{y_n\}_{n \geq 1}$ in \mathcal{Y} .

Proof. (i) and (ii) Throughout these parts of the proof, the following notations is used. $\{\tilde{L}_l\}_{l \geq 1}$, $\{L_l\}_{l \geq 1}$ are the real numbers defined by

$$\tilde{L}_l = \frac{2^{l+1}d}{\delta_1} \left(\|\mu\| + \frac{1}{\gamma} \right)^l, \quad L_l = 2\tilde{L}_l^2 \quad (43)$$

for $l \geq 1$, where γ , K_l are specified in Assumption 2.3 and Lemma 5.3. m, n are any integers satisfying $n > m \geq 0$. In what follows in the proof of (i), (ii), both m, n are kept fixed. η, η', η'' are any elements in $V_{\delta_1}(\Theta)$. ξ, ξ', ξ'' are any elements of $V_{\delta_1}(\mathcal{P}(\mathcal{X}))$. x, x' are any elements of \mathcal{X} , while $\mathbf{y} = \{y_n\}_{n \geq 0}$ is any sequence in \mathcal{Y} .

Using (33), (34), (39), it is straightforward to verify

$$\begin{aligned} \frac{\langle R_{\eta, \mathbf{y}}^{m:n}(\xi) \rangle}{\varphi_{\eta, \mathbf{y}}^{m:n}} &= \int \dots \int \frac{v_{\eta, \mathbf{y}}^{m:n}(x_{m:n})}{\varphi_{\eta, \mathbf{y}}^{m:n}} \\ &\quad \cdot \mu(dx_n) \dots \mu(dx_{m+1}) \xi(dx_m). \end{aligned} \quad (44)$$

Moreover, Lemma 5.2 yields

$$\left| \frac{v_{\eta, \mathbf{y}}^{m:n}(x_{m:n})}{\varphi_{\eta, \mathbf{y}}^{m:n}} \right| = \prod_{k=m+1}^n \left| \frac{\tilde{r}_\eta(y_k, x_k | x_{k-1})}{\varphi_\eta(y_k)} \right| \leq 2^{n-m}. \quad (45)$$

Since $v_{\eta, \mathbf{y}}^{m:n}(x_{m:n})/\varphi_{\eta, \mathbf{y}}^{m:n}$ is analytic in η for each $\eta \in V_{\delta_1}(\Theta)$, $x_m, \dots, x_n \in \mathcal{X}$ (due to Assumption 2.2), Lemma A1.1 (see Appendix 1) and (44), (45) imply that $\langle R_{\eta, \mathbf{y}}^{m:n}(\xi) \rangle/\varphi_{\eta, \mathbf{y}}^{m:n}$ is analytic in η for all $\eta \in V_{\delta_1}(\Theta)$. Consequently, $\langle R_{\eta, \mathbf{y}}^{m:n}(\xi) \rangle$ is analytic in η for each $\eta \in V_{\delta_1}(\Theta)$. Hence, (i) holds.

Owing to (44), (45), we have

$$\begin{aligned} \left| \frac{\langle R_{\eta, \mathbf{y}}^{m:n}(\xi) \rangle}{\varphi_{\eta, \mathbf{y}}^{m:n}} \right| &\leq \int \dots \int \left| \frac{v_{\eta, \mathbf{y}}^{m:n}(x_{m:n})}{\varphi_{\eta, \mathbf{y}}^{m:n}} \right| \\ &\quad \cdot \mu(dx_n) \dots \mu(dx_{m+1}) |\xi|(dx_m) \\ &\leq 2^{n-m} \|\mu\|^{n-m} \|\xi\| \\ &\leq \tilde{L}_{n-m} \leq L_{n-m} \end{aligned} \quad (46)$$

as $\xi \in V_{\delta_1}(\mathcal{P}(\mathcal{X}))$ results in $\|\xi\| \leq 1 + \delta_1 \leq 2$. Using similar arguments, we get

$$\begin{aligned} \left| \frac{\langle R_{\eta, \mathbf{y}}^{m:n}(\xi') \rangle}{\varphi_{\eta, \mathbf{y}}^{m:n}} - \frac{\langle R_{\eta, \mathbf{y}}^{m:n}(\xi'') \rangle}{\varphi_{\eta, \mathbf{y}}^{m:n}} \right| &= \left| \frac{\langle R_{\eta, \mathbf{y}}^{m:n}(\xi' - \xi'') \rangle}{\varphi_{\eta, \mathbf{y}}^{m:n}} \right| \\ &\leq 2^{n-m} \|\mu\|^{n-m} \|\xi' - \xi''\| \\ &\leq \tilde{L}_{n-m} \|\xi' - \xi''\|. \end{aligned} \quad (47)$$

Since $\langle R_{\eta, \mathbf{y}}^{m:n}(\xi) \rangle/\varphi_{\eta, \mathbf{y}}^{m:n}$ is analytic in η for each $\eta \in V_{\delta_1}(\Theta)$, Lemma A1.1 and (46) imply

$$\begin{aligned} \left| \frac{\langle R_{\eta', \mathbf{y}}^{m:n}(\xi) \rangle}{\varphi_{\eta', \mathbf{y}}^{m:n}} - \frac{\langle R_{\eta'', \mathbf{y}}^{m:n}(\xi) \rangle}{\varphi_{\eta'', \mathbf{y}}^{m:n}} \right| &\leq \frac{2^{n-m} d \|\mu\|^{n-m} \|\xi\| \|\eta' - \eta''\|}{\delta_1} \\ &\leq \tilde{L}_{n-m} \|\eta' - \eta''\|. \end{aligned}$$

Then, we have

$$\begin{aligned} &\left| \frac{\langle R_{\eta', \mathbf{y}}^{m:n}(\xi') \rangle}{\varphi_{\eta', \mathbf{y}}^{m:n}} - \frac{\langle R_{\eta'', \mathbf{y}}^{m:n}(\xi'') \rangle}{\varphi_{\eta'', \mathbf{y}}^{m:n}} \right| \\ &\leq \left| \frac{\langle R_{\eta', \mathbf{y}}^{m:n}(\xi') \rangle}{\varphi_{\eta', \mathbf{y}}^{m:n}} - \frac{\langle R_{\eta'', \mathbf{y}}^{m:n}(\xi') \rangle}{\varphi_{\eta'', \mathbf{y}}^{m:n}} \right| \\ &\quad + \left| \frac{\langle R_{\eta'', \mathbf{y}}^{m:n}(\xi') \rangle}{\varphi_{\eta'', \mathbf{y}}^{m:n}} - \frac{\langle R_{\eta'', \mathbf{y}}^{m:n}(\xi'') \rangle}{\varphi_{\eta'', \mathbf{y}}^{m:n}} \right| \\ &\leq \tilde{L}_{n-m} (\|\eta' - \eta''\| + \|\xi' - \xi''\|). \end{aligned} \quad (48)$$

Using (46), (48), we conclude that (ii) is true.

(iii) and (iv) Throughout these parts of the proof, we use the following notations. $\{\tilde{L}_l\}_{l \geq 1}$ has the same meaning as in (43), while $\{\alpha_l\}_{l \geq 1}$, $\{\tilde{M}_l\}_{l \geq 1}$, $\{M_l\}_{l \geq 1}$ are the numbers defined by

$$\alpha_l = \frac{\delta_1}{4\tilde{L}_l^2}, \quad \tilde{M}_l = 10\tilde{L}_l^4, \quad M_l = 5\tilde{M}_l^2 (\|\mu\| + 1).$$

m, n are any integers satisfying $n > m \geq 0$. In what follows in the proof of (iii), (iv), both m, n are kept fixed. η, η', η'' are any elements of $V_{\alpha_{n-m}}(\Theta)$, while θ is any element of Θ satisfying $\|\eta - \theta\| < \alpha_{n-m}$. ξ, ξ', ξ'' are any elements of $V_{\alpha_{n-m}}(\mathcal{P}(\mathcal{X}))$, while λ is any element of $\mathcal{P}(\mathcal{X})$ satisfying $\|\xi - \lambda\| < \alpha_{n-m}$. x, x' are any elements of \mathcal{X} , while $\mathbf{y} = \{y_n\}_{n \geq 1}$ is any sequence in \mathcal{Y} .

Using Lemma 5.2 and (33), it is straightforward to verify

$$\begin{aligned} \langle R_{\theta, \mathbf{y}}^{m:k+1}(\lambda) \rangle &= \int \dots \int \left(\int r_\theta(y_{k+1}, x_{k+1} | x_k) \mu(dx_{k+1}) \right) \\ &\quad \cdot v_{\theta, \mathbf{y}}^{m:k}(x_{m:k}) \mu(dx_k) \dots \mu(dx_{m+1}) \lambda(dx_m) \end{aligned}$$

for $k > m$. Consequently, Assumption 2.3 yields

$$\begin{aligned} \langle R_{\theta, \mathbf{y}}^{m:k+1}(\lambda) \rangle &\geq \gamma |\varphi_\theta(y_{k+1})| \int \dots \int \int v_{\theta, \mathbf{y}}^{m:k}(x_{m:k}) \\ &\quad \cdot \mu(dx_k) \dots \mu(dx_{m+1}) \lambda(dx_m) \\ &= \gamma |\varphi_\theta(y_{k+1})| \langle R_{\theta, \mathbf{y}}^{m:k}(\lambda) \rangle. \end{aligned} \quad (49)$$

The same arguments also imply

$$\begin{aligned} \langle R_{\theta, \mathbf{y}}^{m:m+1}(\lambda) \rangle &= \int \left(\int r_\theta(y_{m+1}, x_{m+1} | x_m) \mu(dx_{m+1}) \right) \lambda(dx_m) \\ &\geq \gamma |\varphi_\theta(y_{m+1})| \|\lambda\| = \gamma |\varphi_\theta(y_{m+1})|. \end{aligned}$$

Then, iterating (49), we get

$$\begin{aligned} \langle R_{\theta,y}^{m:k+1}(\lambda) \rangle &\geq \gamma^{k-m-1} \left(\prod_{l=m+2}^{k+1} |\varphi_{\theta}(y_l)| \right) \langle R_{\theta,y}^{m:m+1}(\lambda) \rangle \\ &\geq \gamma^{k-m} \left(\prod_{l=m+1}^{k+1} |\varphi_{\theta}(y_l)| \right) = \gamma^{k-m} |\varphi_{\theta,y}^{m:n}|. \end{aligned}$$

Hence, we have

$$\frac{\langle R_{\theta,y}^{m:n}(\lambda) \rangle}{|\varphi_{\theta,y}^{m:n}|} \geq \frac{1}{\tilde{L}_{n-m}}$$

as $\tilde{L}_{n-m} \geq \gamma^{-(n-m)}$. Combining this with (48), we get

$$\begin{aligned} \frac{\operatorname{Re} \left\{ \left\langle R_{\eta,y}^{m:n}(\xi) \right\rangle \right\}}{|\varphi_{\eta,y}^{m:n}|} &\geq \frac{\langle R_{\theta,y}^{m:n}(\lambda) \rangle}{|\varphi_{\theta,y}^{m:n}|} - \left| \frac{\langle R_{\eta,y}^{m:n}(\xi) \rangle}{\varphi_{\eta,y}^{m:n}} - \frac{\langle R_{\theta,y}^{m:n}(\lambda) \rangle}{\varphi_{\theta,y}^{m:n}} \right| \\ &\geq \frac{1}{\tilde{L}_{n-m}} - \tilde{L}_{n-m} (\|\eta - \theta\| + \|\xi - \lambda\|) \\ &\geq \frac{1}{\tilde{L}_{n-m}} - 2\tilde{L}_{n-m}\alpha_{n-m} \\ &\geq \frac{1}{2\tilde{L}_{n-m}} \geq \frac{1}{2\tilde{L}_{n-m}} \end{aligned} \quad (50)$$

as $\|\eta - \theta\| < \alpha_{n-m}$, $\|\xi - \lambda\| < \alpha_{n-m}$.

Using (33), it is straightforward to verify

$$\begin{aligned} \frac{r_{\eta,y}^{m:n}(x'|x)}{\varphi_{\eta,y}^{m:n}} &= \int \int \dots \int \int \frac{v_{\eta,y}^{m:n}(x_{m:n})}{\varphi_{\eta,y}^{m:n}} \\ &\quad \cdot \delta_{x'}(dx_n) \mu(dx_{n-1}) \dots \mu(dx_{m+1}) \delta_x(dx_m). \end{aligned} \quad (51)$$

Consequently, Assumption 2.2, Lemma A1.1 and (45) imply that $r_{\eta,y}^{m:n}(x'|x)/\varphi_{\eta,y}^{m:n}$ is analytic in η for each $\eta \in V_{\delta_1}(\Theta)$. Therefore, $r_{\eta,y}^{m:n}(x'|x)$ is analytic in η for all $\eta \in V_{\delta_1}(\Theta)$. Since $\langle R_{\eta,y}^{m:n}(\xi) \rangle$ is non-zero and analytic in η for all $\eta \in V_{\delta_1}(\Theta)$, we then conclude from (35), (50) that $g_{\eta,y}^{m:n}(x'|x, \xi)$ is analytic in η for all $\eta \in V_{\delta_1}(\Theta)$.

Owing to (45), (51), we have

$$\begin{aligned} \left| \frac{r_{\eta,y}^{m:n}(x'|x)}{\varphi_{\eta,y}^{m:n}} \right| &\leq \int \int \dots \int \int \left| \frac{v_{\eta,y}^{m:n}(x_{m:n})}{\varphi_{\eta,y}^{m:n}} \right| \\ &\quad \cdot \delta_{x'}(dx_n) \mu(dx_{n-1}) \dots \mu(dx_{m+1}) \delta_x(dx_m) \\ &\leq 2^{n-m} \|\delta_x\| \|\delta_{x'}\| \|\mu\|^{n-m-1} \leq \tilde{L}_{n-m}. \end{aligned} \quad (52)$$

Then, (35), (50) imply

$$\left| g_{\eta,y}^{m:n}(x'|x, \xi) \right| = \left| \frac{r_{\eta,y}^{m:n}(x'|x)}{\langle R_{\eta,y}^{m:n}(\xi) \rangle} \right| \leq 2\tilde{L}_{n-m}^2 \leq \tilde{M}_{n-m} \quad (53)$$

as $|\langle R_{\eta,y}^{m:n}(\xi) \rangle| \geq \operatorname{Re}(\langle R_{\eta,y}^{m:n}(\xi) \rangle) > 0$. Consequently, (36) yields

$$\begin{aligned} \left| f_{\eta,y}^{m:n}(x|\xi) \right| &\leq \int \left| g_{\eta,y}^{m:n}(x|x', \xi) \right| |\xi|(dx') \\ &\leq \tilde{M}_{n-m} \|\xi\| \leq 2\tilde{M}_{n-m} \leq M_{n-m} \end{aligned} \quad (54)$$

as $\xi \in V_{\alpha_{n-m}}(\mathcal{P}(\mathcal{X}))$ results in $\|\xi\| \leq 1 + \alpha_{n-m} \leq 2$. Similarly, we have

$$\int \left| g_{\eta,y}^{m:n}(x'|x, \xi) \right| \mu(dx') \leq \tilde{M}_{n-m} \|\mu\|. \quad (55)$$

Combining this with (37), (53), (54), we get

$$\begin{aligned} \left| h_{\eta,y}^{m:n}(x'|x, \xi) \right| &\leq \left| f_{\eta,y}^{m:n}(x|\xi) \right| \int \left| g_{\eta,y}^{m:n}(x''|x, \xi) \right| \mu(dx'') \\ &\quad + \left| g_{\eta,y}^{m:n}(x'|x, \xi) \right| \\ &\leq \tilde{M}_{n-m} + 2\tilde{M}_{n-m}^2 \|\mu\| \leq M_{n-m}. \end{aligned} \quad (56)$$

Since $g_{\eta,y}^{m:n}(x'|x, \xi)$ is analytic in η for each $\eta \in V_{\alpha_{n-m}}(\Theta)$, Lemma A1.1 and (53) imply

$$\begin{aligned} \left| g_{\eta',y}^{m:n}(x'|x, \xi) - g_{\eta'',y}^{m:n}(x'|x, \xi) \right| &\leq \frac{2d\tilde{L}_{n-m}^2 \|\eta' - \eta''\|}{\alpha_{n-m}} \\ &\leq \tilde{M}_{n-m} \|\eta' - \eta''\|. \end{aligned} \quad (57)$$

Moreover, (47), (50), (53) yield

$$\begin{aligned} \left| g_{\eta,y}^{m:n}(x'|x, \xi') - g_{\eta,y}^{m:n}(x'|x, \xi'') \right| &= \left| g_{\eta,y}^{m:n}(x'|x, \xi') \right| \left| \frac{\langle R_{\eta,y}^{m:n}(\xi') \rangle - \langle R_{\eta,y}^{m:n}(\xi'') \rangle}{\langle R_{\eta,y}^{m:n}(\xi'') \rangle} \right| \\ &\leq 2\tilde{L}_{n-m}^4 \|\xi' - \xi''\| \leq \tilde{M}_{n-m} \|\xi' - \xi''\|. \end{aligned} \quad (58)$$

Combining (57), (58), we get

$$\begin{aligned} \left| g_{\eta',y}^{m:n}(x'|x, \xi') - g_{\eta'',y}^{m:n}(x'|x, \xi'') \right| &\leq \left| g_{\eta',y}^{m:n}(x'|x, \xi') - g_{\eta'',y}^{m:n}(x'|x, \xi') \right| \\ &\quad + \left| g_{\eta'',y}^{m:n}(x'|x, \xi') - g_{\eta'',y}^{m:n}(x'|x, \xi'') \right| \\ &\leq \tilde{M}_{n-m} (\|\eta' - \eta''\| + \|\xi' - \xi''\|). \end{aligned} \quad (59)$$

Consequently, (36), (53) imply

$$\begin{aligned} \left| f_{\eta',y}^{m:n}(x|\xi') - f_{\eta'',y}^{m:n}(x|\xi'') \right| &\leq \int \left| g_{\eta',y}^{m:n}(x|x', \xi') - g_{\eta'',y}^{m:n}(x|x', \xi'') \right| |\xi'| (dx') \\ &\quad + \int \left| g_{\eta'',y}^{m:n}(x|x', \xi'') \right| |\xi' - \xi''| (dx') \\ &\leq \tilde{M}_{n-m} \|\xi'\| (\|\eta' - \eta''\| + \|\xi' - \xi''\|) + \tilde{M}_{n-m} \|\xi' - \xi''\| \\ &\leq 3\tilde{M}_{n-m} (\|\eta' - \eta''\| + \|\xi' - \xi''\|) \\ &\leq M_{n-m} (\|\eta' - \eta''\| + \|\xi' - \xi''\|) \end{aligned} \quad (60)$$

as $\|\xi'\| \leq 1 + \alpha_{n-m} \leq 2$. Similarly, we get

$$\begin{aligned} \int \left| g_{\eta',y}^{m:n}(x'|x, \xi') - g_{\eta'',y}^{m:n}(x'|x, \xi'') \right| \mu(dx') &\leq \tilde{M}_{n-m} \|\mu\| (\|\eta' - \eta''\| + \|\xi' - \xi''\|). \end{aligned}$$

Combining this with (37), (54), (55), (59), (60), we get

$$\begin{aligned} \left| h_{\eta',y}^{m:n}(x'|x, \xi') - h_{\eta'',y}^{m:n}(x'|x, \xi'') \right| &\leq \left| g_{\eta',y}^{m:n}(x'|x, \xi') - g_{\eta'',y}^{m:n}(x'|x, \xi'') \right| \\ &\quad + \left| f_{\eta',y}^{m:n}(x|\xi') - f_{\eta'',y}^{m:n}(x|\xi'') \right| \int \left| g_{\eta',y}^{m:n}(x''|x, \xi') \right| \mu(dx'') \\ &\quad + \left| f_{\eta'',y}^{m:n}(x|\xi'') \right| \int \left| g_{\eta'',y}^{m:n}(x''|x, \xi') - g_{\eta'',y}^{m:n}(x''|x, \xi'') \right| \mu(dx'') \end{aligned}$$

$$\begin{aligned} &\leq (\tilde{M}_{n-m} + 5\tilde{M}_{n-m}^2 \|\mu\|)(\|\eta' - \eta''\| + \|\xi' - \xi''\|) \\ &\leq M_{n-m}(\|\eta' - \eta''\| + \|\xi' - \xi''\|). \end{aligned} \quad (61)$$

Using (50), (54), (56) – (61), we conclude that (iii), (iv) hold. \square

Lemma 6.4. *Let Assumptions 2.2 and 2.3 hold. Then, the following is true:*

(i) *There exists a real number $\delta_4 \in (0, \delta_1]$ such that $\text{Re}\{R_{\eta,y}(\mathcal{X}|\xi)\} > 0$ for all $\eta \in V_{\delta_4}(\Theta)$, $\xi \in V_{\delta_4}(\mathcal{P}(\mathcal{X}))$, $y \in \mathcal{Y}$ (δ_1 is specified in Lemma 5.2).*

(ii) *There exists a real number $C_4 \in [1, \infty)$ such that*

$$\begin{aligned} |\Phi_{\eta,y}(\xi)| &\leq C_4(1 + \psi(y)), \\ |\Phi_{\eta',y}(\xi') - \Phi_{\eta'',y}(\xi'')| \\ &\leq C_4(1 + \psi(y))(\|\eta' - \eta''\| + \|\xi' - \xi''\|) \end{aligned}$$

for all $\eta, \eta', \eta'' \in V_{\delta_4}(\Theta)$, $\xi, \xi', \xi'' \in V_{\delta_4}(\mathcal{P}(\mathcal{X}))$, $y \in \mathcal{Y}$.

Proof. Throughout the proof, the following notations is used. δ_4, C_4 are the real numbers defined by $\delta_4 = \alpha_1$, $C_4 = 4L_1^2$ (α_1, L_1 are specified in Lemma 6.3). η, η', η'' are any elements of $V_{\delta_4}(\Theta)$, while ξ, ξ', ξ'' are any elements in $V_{\delta_4}(\mathcal{P}(\mathcal{X}))$. y is any element of \mathcal{Y} .

Since $R_{\eta,y}(\mathcal{X}|y) = \langle R_{\eta,y}^{0:1}(\xi) \rangle$ for any sequence $\mathbf{y} = \{y_n\}_{n \geq 1}$ in \mathcal{Y} satisfying $y = y_1$, Lemma 6.3 yields

$$\text{Re}\{R_{\eta,y}(\mathcal{X}|\xi)\} \geq \frac{|\varphi_{\eta}(y)|}{L_1}, \quad (62)$$

$$|R_{\eta,y}(\mathcal{X}|\xi)| \leq L_1|\varphi_{\eta}(y)|, \quad (63)$$

$$|R_{\eta,y}(\mathcal{X}|\xi') - R_{\eta,y}(\mathcal{X}|\xi'')| \leq L_1|\varphi_{\eta}(y)|\|\xi' - \xi''\|. \quad (64)$$

Due to the same arguments, $R_{\eta,y}(\mathcal{X}|\xi)$ is analytic in η for each $\eta \in V_{\delta_4}(\Theta)$. As $\varphi_{\eta}(y) \neq 0$ (owing to Assumption 2.2), (62) implies that (i) holds. Consequently, (30) yields that $\Phi_{\eta,y}(\xi)$ is analytic in η for all $\eta \in V_{\delta_4}(\Theta)$. Moreover, due to (62), (63), we have

$$\begin{aligned} \log |R_{\eta,y}(\mathcal{X}|\xi)| &\leq \log L_1 + \log |\varphi_{\eta}(y)| \leq L_1(1 + \psi(y)), \\ \log |R_{\eta,y}(\mathcal{X}|\xi)| &\geq -\log L_1 + \log |\varphi_{\eta}(y)| \geq -L_1(1 + \psi(y)). \end{aligned}$$

Therefore, we get

$$\begin{aligned} |\Phi_{\eta,y}(\xi)| &= |\log R_{\eta,y}(\mathcal{X}|\xi)| \leq |\log |R_{\eta,y}(\mathcal{X}|\xi)|| + \pi \\ &\leq 4L_1(1 + \psi(y)) \\ &\leq C_4(1 + \psi(y)). \end{aligned} \quad (65)$$

Then, Lemma A1.1 implies

$$\begin{aligned} |\Phi_{\eta',y}(\xi) - \Phi_{\eta'',y}(\xi)| &\leq \frac{4dL_1(1 + \psi(y))\|\eta' - \eta''\|}{\delta_4} \\ &\leq C_4(1 + \psi(y))\|\eta' - \eta''\|. \end{aligned} \quad (66)$$

Let $\phi_{\eta,y}(t|\xi', \xi'')$ be the function defined by

$$\phi_{\eta,y}(t|\xi', \xi'') = \log(tR_{\eta,y}(\mathcal{X}|\xi') + (1-t)R_{\eta,y}(\mathcal{X}|\xi''))$$

for $t \in [0, 1]$. Due to Assumption 2.2 and (62), we have

$$\begin{aligned} &|tR_{\eta,y}(\mathcal{X}|\xi') + (1-t)R_{\eta,y}(\mathcal{X}|\xi'')| \\ &\geq t\text{Re}\{R_{\eta,y}(\mathcal{X}|\xi')\} + (1-t)\text{Re}\{R_{\eta,y}(\mathcal{X}|\xi'')\} \\ &\geq \frac{|\varphi_{\eta}(y)|}{L_1} > 0 \end{aligned} \quad (67)$$

for $t \in [0, 1]$. Hence, $\phi_{\eta,y}(t|\xi', \xi'')$ is well-defined and differentiable in t for each $t \in [0, 1]$. We also have

$$\begin{aligned} \phi'_{\eta,y}(t|\xi', \xi'') &= \frac{\partial}{\partial t} \phi_{\eta,y}(t|\xi', \xi'') \\ &= \frac{\text{Re}\{R_{\eta,y}(\mathcal{X}|\xi')\} - \text{Re}\{R_{\eta,y}(\mathcal{X}|\xi'')\}}{t\text{Re}\{R_{\eta,y}(\mathcal{X}|\xi')\} + (1-t)\text{Re}\{R_{\eta,y}(\mathcal{X}|\xi'')\}}. \end{aligned}$$

Consequently, (64), (67) yield

$$|\phi'_{\eta,y}(t|\xi', \xi'')| \leq L_1^2\|\xi' - \xi''\|.$$

Thus, we get

$$\begin{aligned} |\Phi_{\eta,y}(\xi') - \Phi_{\eta,y}(\xi'')| &= |\phi_{\eta,y}(1|\xi', \xi'') - \phi_{\eta,y}(0|\xi', \xi'')| \\ &= \left| \int_0^1 \phi'_{\eta,y}(t|\xi', \xi'') dt \right| \\ &\leq L_1^2\|\xi' - \xi''\| \leq C_4\|\xi' - \xi''\|. \end{aligned}$$

Consequently, (66) implies

$$\begin{aligned} |\Phi_{\eta',y}(\xi') - \Phi_{\eta'',y}(\xi'')| &\leq |\Phi_{\eta',y}(\xi') - \Phi_{\eta'',y}(\xi')| \\ &\quad + |\Phi_{\eta'',y}(\xi') - \Phi_{\eta'',y}(\xi'')| \\ &\leq C_4(1 + \psi(y))(\|\eta' - \eta''\| + \|\xi' - \xi''\|). \end{aligned} \quad (68)$$

Using (65), (68), we deduce that (ii) is true. \square

Lemma 6.5. *Let Assumptions 2.1 – 2.3 hold. Then, the following is true:*

(i) *There exist real numbers $\delta_5, \delta_6 \in (0, \delta_4]$, $C_5 \in [1, \infty)$ and an integer $n_0 \geq 1$ such that $\text{Re}\{R_{\eta,y}^{m:n}(\xi)\} > 0$, $F_{\eta,y}^{m:n}(\xi) \in V_{\delta_4}(\mathcal{P}(\mathcal{X}))$, $F_{\eta,y}^{m:m+n_0}(\xi) \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$ and*

$$\|F_{\eta,y}^{m:n}(\xi') - F_{\eta,y}^{m:n}(\xi'')\| \leq C_5\|\xi' - \xi''\|, \quad (69)$$

$$\|F_{\eta,y}^{m:m+n_0}(\xi') - F_{\eta,y}^{m:m+n_0}(\xi'')\| \leq \frac{\|\xi' - \xi''\|}{2} \quad (70)$$

for all $\eta \in V_{\delta_5}(\Theta)$, $\xi, \xi', \xi'' \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$, $m + n_0 \geq n \geq m \geq 0$ and any sequence $\mathbf{y} = \{y_n\}_{n \geq 1}$ in \mathcal{Y} (δ_4 is specified in Lemma 6.4).

(ii) *There exist real numbers $\delta_7 \in (0, \delta_5]$, $\delta_8 \in (0, \delta_6]$ such that $F_{\eta,y}^{m:n}(\xi) \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$ for all $\eta \in V_{\delta_7}(\Theta)$, $\xi \in V_{\delta_8}(\mathcal{P}(\mathcal{X}))$, $m + n_0 \geq n \geq m \geq 0$ and any sequence $\mathbf{y} = \{y_n\}_{n \geq 1}$ in \mathcal{Y} .*

Proof. (i) Throughout this part of the proof, the following notations is used. n_0 is the integer defined by

$$n_0 = \left\lceil \frac{\log(4C_3)}{|\log \gamma_2|} \right\rceil,$$

while C_5, δ_5, δ_6 are the real numbers defined by $C_5 = M_{n_0}(1 + \|\mu\|)$ and

$$\delta_5 = \frac{\min\{\alpha_{n_0}, \delta_4\}}{16C_5^2}, \quad \delta_6 = 2C_5\delta_5 \quad (71)$$

($\gamma_2, C_3, \alpha_n, M_n$ are specified in Lemmas 6.2, 6.3). η, η', η'' are any elements in $V_{\delta_5}(\Theta)$, while θ is any element of Θ satisfying $\|\eta - \theta\| < \delta_5$. ξ, ξ', ξ'' are any elements in $V_{\delta_6}(\mathcal{P}(\mathcal{X}))$, while λ is any element of $\mathcal{P}(\mathcal{X})$ satisfying $\|\xi - \lambda\| < \delta_6$. x is any element of \mathcal{X} , while $\mathbf{y} = \{y_n\}_{n \geq 1}$ is any sequence in \mathcal{Y} . B is any element of $\mathcal{B}(\mathcal{X})$. m, n are any integers satisfying $m + n_0 \geq n > m > 0$.

Owing to Lemma 6.3, we have $\text{Re}\{R_{\eta,y}^{m:n}(\xi)\} > 0$, as $\delta_5 \leq \delta_6 \leq \alpha_{n_0} \leq \alpha_{n-m}$ results from $n-m \leq n_0$. Hence, we get

$$\begin{aligned} & \text{Re} \left\{ \left\langle R_{\eta,y}^{m:n}(t\xi' + (1-t)\xi'') \right\rangle \right\} \\ &= t \text{Re} \left\{ \left\langle R_{\eta,y}^{m:n}(\xi') \right\rangle \right\} + (1-t) \text{Re} \left\{ \left\langle R_{\eta,y}^{m:n}(\xi'') \right\rangle \right\} > 0 \end{aligned} \quad (72)$$

for $t \in [0, 1]$. Moreover, using Lemma 6.3, we conclude

$$\begin{aligned} & \left| \int I_B(x') \left(h_{\eta',y}^{m:n}(x'|x, \xi') - h_{\eta'',y}^{m:n}(x'|x, \xi'') \right) \mu(dx') \right| \\ & \leq \int I_B(x') \left| h_{\eta',y}^{m:n}(x'|x, \xi') - h_{\eta'',y}^{m:n}(x'|x, \xi'') \right| \mu(dx') \\ & \leq M_{n-m} \|\mu\| (\|\eta' - \eta''\| + \|\xi' - \xi''\|) \\ & \leq C_5 (\|\eta' - \eta''\| + \|\xi' - \xi''\|) \end{aligned} \quad (73)$$

as $C_5 \geq M_{n_0} \geq M_{n-m}$ results from $n-m \leq n_0$. Relying on the same lemma, we deduce

$$\begin{aligned} & \left| F_{\eta',y}^{m:n}(B|\xi') - F_{\eta'',y}^{m:n}(B|\xi'') \right| \\ & \leq \int I_B(x) \left| f_{\eta',y}^{m:n}(x|\xi') - f_{\eta'',y}^{m:n}(x|\xi'') \right| \mu(dx) \\ & \leq M_{n-m} \|\mu\| (\|\eta' - \eta''\| + \|\xi' - \xi''\|) \\ & \leq C_5 (\|\eta' - \eta''\| + \|\xi' - \xi''\|). \end{aligned}$$

Hence, we have

$$\left\| F_{\eta',y}^{m:n}(\xi') - F_{\eta'',y}^{m:n}(\xi'') \right\| \leq C_5 (\|\eta' - \eta''\| + \|\xi' - \xi''\|) \quad (74)$$

as $\|\eta - \theta\| < \delta_5$, $\|\xi - \lambda\| < \delta_6$. Setting $\eta' = \eta$, $\eta'' = \eta$ in (74), we get (69). We also get

$$\begin{aligned} \left\| F_{\eta,y}^{m:n}(\xi) - F_{\theta,y}^{m:n}(\lambda) \right\| & \leq C_5 (\|\eta - \theta\| + \|\xi - \lambda\|) \\ & < C_5(\delta_5 + \delta_6) \leq \delta_4. \end{aligned}$$

Therefore, $F_{\eta,y}^{m:n}(\xi) \in V_{\delta_4}(\mathcal{P}(\mathcal{X}))$ for $m+n_0 \geq n > m \geq 0$, as $F_{\theta,y}^{m:n}(\lambda) \in \mathcal{P}(\mathcal{X})$.

Let $\phi_{\eta,y}^{m:n}(t, x|\xi', \xi'')$ be the function defined by

$$\phi_{\eta,y}^{m:n}(t, x|\xi', \xi'') = f_{\eta,y}^{m:n}(x|t\xi' + (1-t)\xi'') \quad (75)$$

for $t \in [0, 1]$, $m+n_0 \geq n > m \geq 0$. Then, due to (72), we have

$$\begin{aligned} & \phi_{\eta,y}^{m:n}(t, x|\xi', \xi'') \\ &= \frac{\int r_{\eta,y}^{m:n}(x|x')(t\xi' + (1-t)\xi'')(dx')}{\left\langle R_{\eta,y}^{m:n}(t\xi' + (1-t)\xi'') \right\rangle} \\ &= \frac{t \int r_{\eta,y}^{m:n}(x|x')\xi'(dx') + (1-t) \int r_{\eta,y}^{m:n}(x|x')\xi''(dx')}{t \left\langle R_{\eta,y}^{m:n}(\xi') \right\rangle + (1-t) \left\langle R_{\eta,y}^{m:n}(\xi'') \right\rangle}. \end{aligned}$$

Thus, we get

$$\begin{aligned} \frac{\partial}{\partial t} \phi_{\eta,y}^{m:n}(t, x|\xi', \xi'') &= \frac{\int r_{\eta,y}^{m:n}(x|x')(\xi' - \xi'')(dx')}{\left\langle R_{\eta,y}^{m:n}(t\xi' + (1-t)\xi'') \right\rangle} \\ & \quad - f_{\eta,y}^{m:n}(x|t\xi' + (1-t)\xi'') \\ & \quad \cdot \frac{\left\langle R_{\eta,y}^{m:n}(\xi') \right\rangle - \left\langle R_{\eta,y}^{m:n}(\xi'') \right\rangle}{\left\langle R_{\eta,y}^{m:n}(t\xi' + (1-t)\xi'') \right\rangle} \\ &= \frac{\int r_{\eta,y}^{m:n}(x|x')(\xi' - \xi'')(dx')}{\left\langle R_{\eta,y}^{m:n}(t\xi' + (1-t)\xi'') \right\rangle} \\ & \quad - f_{\eta,y}^{m:n}(x|t\xi' + (1-t)\xi'') \\ & \quad \cdot \frac{\int \int r_{\eta,y}^{m:n}(x''|x')\mu(dx'')(\xi' - \xi'')(dx')}{\left\langle R_{\eta,y}^{m:n}(t\xi' + (1-t)\xi'') \right\rangle}. \end{aligned}$$

Consequently, (35) – (37) imply

$$\begin{aligned} & \frac{\partial}{\partial t} \phi_{\eta,y}^{m:n}(t, x|\xi', \xi'') \\ &= \int g_{\eta,y}^{m:n}(x|x', t\xi' + (1-t)\xi'')(\xi' - \xi'')(dx') \\ & \quad - f_{\eta,y}^{m:n}(x|t\xi' + (1-t)\xi'') \\ & \quad \cdot \int \int g_{\eta,y}^{m:n}(x''|x', t\xi' + (1-t)\xi'')\mu(dx'')(\xi' - \xi'')(dx') \\ &= \int h_{\eta,y}^{m:n}(x|x', t\xi' + (1-t)\xi'')(\xi' - \xi'')(dx'). \end{aligned} \quad (76)$$

Hence, we have

$$\begin{aligned} & f_{\eta,y}^{m:n}(x|\xi') - f_{\eta,y}^{m:n}(x|\xi'') \\ &= \phi_{\eta,y}^{m:n}(1, x|\xi', \xi'') - \phi_{\eta,y}^{m:n}(0, x|\xi', \xi'') \\ &= \int \int_0^1 h_{\eta,y}^{m:n}(x|x', t\xi' + (1-t)\xi'')(\xi' - \xi'')(dx')dt. \end{aligned} \quad (77)$$

Therefore, (38) yields

$$\begin{aligned} & F_{\eta,y}^{m:n}(B|\xi') - F_{\eta,y}^{m:n}(B|\xi'') \\ &= \int I_B(x) \left(f_{\eta,y}^{m:n}(x|\xi') - f_{\eta,y}^{m:n}(x|\xi'') \right) \mu(dx) \\ &= \int \int \int_0^1 I_B(x) h_{\eta,y}^{m:n}(x|x', t\xi' + (1-t)\xi'') \\ & \quad \cdot \mu(dx)(\xi' - \xi'')(dx')dt. \end{aligned} \quad (78)$$

Since $\lambda + at\delta_x \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$ for $a \in (0, \delta_6)$, $t \in [0, 1]$, we then get

$$\begin{aligned} & F_{\theta,y}^{m:n}(B|\lambda + a\delta_x) - F_{\theta,y}^{m:n}(B|\lambda) \\ &= \alpha \int \int_0^1 I_B(x') h_{\theta,y}^{m:n}(x'|x, \lambda + at\delta_x) \mu(dx')dt \end{aligned} \quad (79)$$

for the same α . Moreover, Lemma 6.2 yields

$$\begin{aligned} & \left| F_{\theta,y}^{m:n}(B|\lambda + a\delta_x) - F_{\theta,y}^{m:n}(B|\lambda) \right| \\ & \leq C_3 \gamma_2^{n-m} \|\alpha\delta_x\| = \alpha C_3 \gamma_2^{n-m} \end{aligned} \quad (80)$$

for $\alpha \in (0, \delta_6)$. Combining (79), (80), we get

$$\left| \int_0^1 \int_0^1 I_B(x') h_{\theta, y}^{m:n}(x'|x, \lambda + \alpha t \delta_x) \mu(dx') dt \right| \leq C_3 \gamma_2^{n-m}. \quad (81)$$

Using (73), (81), we conclude

$$\begin{aligned} & \left| \int I_B(x') h_{\theta, y}^{m:n}(x'|x, \lambda) \mu(dx') \right| \\ & \leq \left| \int_0^1 \int_0^1 I_B(x') h_{\theta, y}^{m:n}(x'|x, \lambda + \alpha t \delta_x) \mu(dx') dt \right| \\ & \quad + \int_0^1 \left| \int I_B(x') \left(h_{\theta, y}^{m:n}(x'|x, \lambda + \alpha t \delta_x) - h_{\theta, y}^{m:n}(x'|x, \lambda) \right) \mu(dx') \right| dt \\ & \leq C_3 \gamma_2^{n-m} + C_5 \alpha \end{aligned}$$

for $\alpha \in (0, \delta_6)$, as $\|\alpha t \delta_x\| \leq \alpha$. Letting $\alpha \rightarrow 0$, we deduce

$$\left| \int I_B(x') h_{\theta, y}^{m:n}(x'|x, \lambda) \mu(dx') \right| \leq C_3 \gamma_2^{n-m}.$$

Consequently, (73) yields

$$\begin{aligned} & \left| \int I_B(x') h_{\eta, y}^{m:n}(x'|x, \zeta) \mu(dx') \right| \\ & \leq \left| \int I_B(x') h_{\theta, y}^{m:n}(x'|x, \lambda) \mu(dx') \right| \\ & \quad + \left| \int I_B(x') \left(h_{\eta, y}^{m:n}(x'|x, \zeta) - h_{\theta, y}^{m:n}(x'|x, \lambda) \right) \mu(dx') \right| \\ & \leq C_3 \gamma_2^{n-m} + C_5 (\|\eta - \theta\| + \|\zeta - \lambda\|) \\ & \leq C_3 \gamma_2^{n-m} + C_5 (\delta_5 + \delta_6) \leq C_3 \gamma_2^{n-m} + \frac{1}{4} \end{aligned}$$

as $\|\eta - \theta\| < \delta_5$, $\|\zeta - \lambda\| < \delta_6$, $C_5 \delta_5 \leq C_5 \delta_6 \leq 1/8$. Combining this with (78), we get

$$\begin{aligned} & \left| F_{\eta, y}^{m:n}(B|\zeta') - F_{\eta, y}^{m:n}(B|\zeta'') \right| \\ & \leq \int_0^1 \int_0^1 \left| \int I_B(x') h_{\eta, y}^{m:n}(x'|x, t\zeta' + (1-t)\zeta'') \mu(dx') \right| \\ & \quad \cdot |\zeta' - \zeta''| (dx) dt \\ & \leq \left(C_3 \gamma_2^{n-m} + \frac{1}{4} \right) \|\zeta' - \zeta''\| \end{aligned}$$

as $t\zeta' + (1-t)\zeta'' \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$ results from $t \in [0, 1]$ and the convexity of $V_{\delta_6}(\mathcal{P}(\mathcal{X}))$. Therefore, we have

$$\left\| F_{\eta, y}^{m:n}(\zeta') - F_{\eta, y}^{m:n}(\zeta'') \right\| \leq \left(C_3 \gamma_2^{n-m} + \frac{1}{4} \right) \|\zeta' - \zeta''\|.$$

Hence, we get

$$\begin{aligned} \left\| F_{\eta, y}^{m:n_0}(\zeta') - F_{\eta, y}^{m:n_0}(\zeta'') \right\| & \leq \left(C_3 \gamma_2^{n_0} + \frac{1}{4} \right) \|\zeta' - \zeta''\| \\ & \leq \frac{\|\zeta' - \zeta''\|}{2} \end{aligned} \quad (82)$$

as $C_3 \gamma_2^{n_0} \leq 1/4$. Consequently, (70) holds. Moreover, (74) implies

$$\begin{aligned} & \left\| F_{\eta, y}^{m:n_0}(\zeta) - F_{\theta, y}^{m:n_0}(\lambda) \right\| \\ & \leq \left\| F_{\eta, y}^{m:n_0}(\zeta) - F_{\eta, y}^{m:n_0}(\lambda) \right\| \\ & \quad + \left\| F_{\eta, y}^{m:n_0}(\lambda) - F_{\theta, y}^{m:n_0}(\lambda) \right\| \\ & \leq \frac{\|\zeta - \lambda\|}{2} + C_5 \|\eta - \theta\| \\ & < \frac{\delta_6}{2} + C_5 \delta_5 = \delta_6 \end{aligned}$$

as $\|\eta - \theta\| < \delta_5$, $\|\zeta - \lambda\| < \delta_6$. Thus, $F_{\eta, y}^{m:n_0}(\zeta) \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$ for $m \geq 0$, as $F_{\theta, y}^{m:n_0}(\lambda) \in \mathcal{P}(\mathcal{X})$.

(ii) Let δ_7, δ_8 be the real numbers defined by $\delta_7 = \delta_5$, $\delta_8 = \delta_5$ (δ_5 is specified in (71)). Moreover, let θ, λ, y have the same meaning as in (i), while η, ζ are any elements of $V_{\delta_6}(\Theta)$, $V_{\delta_7}(\mathcal{P}(\mathcal{X}))$ (respectively). Consequently, when $\|\eta - \theta\| < \delta_7$, $\|\zeta - \lambda\| < \delta_8$, (74) yields

$$\begin{aligned} \left\| F_{\eta, y}^{m:n}(\zeta) - F_{\theta, y}^{m:n}(\lambda) \right\| & \leq C_5 (\|\eta - \theta\| + \|\zeta - \lambda\|) \\ & < C_5 (\delta_7 + \delta_8) \leq \delta_6 \end{aligned}$$

for $m + n_0 \geq n > m \geq 0$. Therefore, $F_{\eta, y}^{m:n}(\zeta) \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$ for $m + n_0 \geq n > m \geq 0$. \square

Lemma 6.6. *Let Assumptions 2.1 – 2.3 hold. Then, the following is true:*

(i) $\text{Re}\{R_{\eta, y}^{m:n}(\zeta)\} \neq 0$, $F_{\eta, y}^{m:n}(\zeta) \in V_{\delta_4}(\mathcal{P}(\mathcal{X}))$ for all $\eta \in V_{\delta_5}(\Theta)$, $\zeta \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$, $n \geq m \geq 0$ and any sequence $y = \{y_n\}_{n \geq 1}$ in \mathcal{Y} ($\delta_4, \delta_5, \delta_6$ are specified in Lemmas 6.4, 6.5).

(ii) There exist real numbers $\gamma_3 \in (0, 1)$, $C_6 \in [1, \infty)$ such that

$$\left\| F_{\eta, y}^{m:n}(\zeta') - F_{\eta, y}^{m:n}(\zeta'') \right\| \leq C_6 \gamma_3^{n-m} \|\zeta' - \zeta''\| \quad (83)$$

for all $\eta \in V_{\delta_5}(\Theta)$, $\zeta', \zeta'' \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$, $n \geq m \geq 0$ and any sequence $y = \{y_n\}_{n \geq 1}$ in \mathcal{Y} .

Proof. (i) Let $n_k(m)$ be the integer defined by $n_k(m) = m + kn_0$ for $m, k \geq 0$ (n_0 is specified in Lemma 6.5). Moreover, let $y = \{y_n\}_{n \geq 1}$ be any sequence in \mathcal{Y} .

First, we show

$$\text{Re}\left\{R_{\eta, y}^{m:n}(\zeta)\right\} \neq 0, \quad (84)$$

$$F_{\eta, y}^{m:n}(\zeta) \in V_{\delta_4}(\mathcal{P}(\mathcal{X})), \quad (85)$$

$$F_{\eta, y}^{m:n_k(m)}(\zeta) \in V_{\delta_6}(\mathcal{P}(\mathcal{X})) \quad (86)$$

for each $\eta \in V_{\delta_5}(\Theta)$, $\zeta \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$, $n_k(m) \geq n \geq m \geq 0$, $k \geq 0$. We prove this by induction in k .

Since $n_0(m) = n = m$ when $n_0(m) \geq n \geq m \geq 0$, we have $R_{\eta, y}^{m:n}(\zeta) = \zeta$, $F_{\eta, y}^{m:n}(\zeta) = \zeta$ for $\eta \in V_{\delta_5}(\Theta)$, $\zeta \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$, $n_0(m) \geq n \geq m \geq 0$. Hence, (84) – (86) hold for $k = 0$ and $\eta \in V_{\delta_5}(\Theta)$, $\zeta \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$, $n_k(m) \geq n \geq m \geq 0$. Now, the induction hypothesis is formulated: Suppose that (84) – (86) are true for some $k \geq 0$ and any $\eta \in V_{\delta_5}(\Theta)$, $\zeta \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$, $n_k(m) \geq n \geq m \geq 0$. Then, to show (84) – (86) for $\eta \in V_{\delta_5}(\Theta)$, $\zeta \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$, $n_{k+1}(m) \geq n \geq m \geq 0$, it is sufficient to demonstrate (84) – (86) for $\eta \in V_{\delta_5}(\Theta)$, $\zeta \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$, $n_{k+1}(m) \geq n \geq n_k(m)$, $m \geq 0$.

In the rest of the proof of (i), η, ζ are any elements of $V_{\delta_5}(\Theta)$, $V_{\delta_6}(\mathcal{P}(\mathcal{X}))$ (respectively). Owing to Lemma 6.5, we have $\text{Re}\{\langle R_{\eta,y}^{n_k(m):n}(\zeta) \rangle\} > 0$, $F_{\eta,y}^{n_k(m):n}(\zeta) \in V_{\delta_4}(\mathcal{P}(\mathcal{X}))$, $F_{\eta,y}^{n_k(m):n_{k+1}(m)}(\zeta) \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$ for $n_{k+1}(m) \geq n \geq n_k(m)$, $m \geq 0$. Since $F_{\eta,y}^{m:n_k(m)}(\zeta) \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$ (due to the induction hypothesis), we then get

$$\text{Re}\{\langle R_{\eta,y}^{n_k(m):n}(F_{\eta,y}^{m:n_k(m)}(\zeta)) \rangle\} > 0, \quad (87)$$

$$F_{\eta,y}^{n_k(m):n}(F_{\eta,y}^{m:n_k(m)}(\zeta)) \in V_{\delta_4}(\mathcal{P}(\mathcal{X})), \quad (88)$$

$$F_{\eta,y}^{n_k(m):n_{k+1}(m)}(F_{\eta,y}^{m:n_k(m)}(\zeta)) \in V_{\delta_6}(\mathcal{P}(\mathcal{X})) \quad (89)$$

for $n_{k+1}(m) \geq n \geq n_k(m)$, $m \geq 0$. As $\langle R_{\eta,y}^{m:n_k(m)}(\zeta) \rangle \neq 0$ (due to the induction hypothesis), Lemma 6.1 (Part (ii)) and (87) imply

$$\begin{aligned} \langle R_{\eta,y}^{m:n}(\zeta) \rangle &= \langle R_{\eta,y}^{m:n_k(m)}(\zeta) \rangle \langle R_{\eta,y}^{n_k(m):n}(F_{\eta,y}^{m:n_k(m)}(\zeta)) \rangle \\ &\neq 0 \end{aligned} \quad (90)$$

for $n_{k+1}(m) \geq n \geq n_k(m)$, $m \geq 0$. Since $\langle R_{\eta,y}^{m:n_k(m)}(\zeta) \rangle \neq 0$, $\langle R_{\eta,y}^{m:n_{k+1}(m)}(\zeta) \rangle \neq 0$, $F_{\eta,y}^{m:n_k(m)}(\zeta) \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$ (due to the induction hypothesis and (90)), Lemma 6.1 (Part (iii)) and (88), (89) yield

$$F_{\eta,y}^{m:n}(\zeta) = F_{\eta,y}^{n_k(m):n}(F_{\eta,y}^{m:n_k(m)}(\zeta)) \in V_{\delta_4}(\mathcal{P}(\mathcal{X})), \quad (91)$$

$$\begin{aligned} F_{\eta,y}^{m:n_{k+1}(m)}(\zeta) &= F_{\eta,y}^{n_k(m):n_{k+1}(m)}(F_{\eta,y}^{m:n_k(m)}(\zeta)) \\ &\in V_{\delta_6}(\mathcal{P}(\mathcal{X})) \end{aligned} \quad (92)$$

for $n_{k+1}(m) \geq n \geq n_k(m)$, $m \geq 0$. Combining (90) – (92) with the induction hypothesis, we deduce that (84) – (86) hold for $n_{k+1}(m) \geq n \geq m$, $m \geq 0$. Then, relying on the principle of mathematical induction, we conclude that (84) – (86) are satisfied for each $\eta \in V_{\delta_5}(\Theta)$, $\zeta \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$, $n_k(m) \geq n \geq m \geq 0$, $k \geq 0$. As a direct consequence of this, we have that (i) is true.

(ii) Let γ_3, C_6 be the real numbers defined by $\gamma_3 = 2^{-1/n_0}$, $C_6 = C_5\gamma_3^{-n_0}$ (C_5, n_0 are specified in Lemma 6.5), while $n_k(m), y$ have the same meaning as in (i). Moreover, let η be any element of $V_{\delta_6}(\Theta)$, while ζ', ζ'' are any elements in $V_{\delta_6}(\mathcal{P}(\mathcal{X}))$.

Owing to Lemmas 6.1, 6.5 and (84) – (86), we have

$$\begin{aligned} &\|F_{\eta,y}^{m:n_{k+1}(m)}(\zeta') - F_{\eta,y}^{m:n_{k+1}(m)}(\zeta'')\| \\ &= \|F_{\eta,y}^{n_k(m):n_{k+1}(m)}(F_{\eta,y}^{m:n_k(m)}(\zeta')) \\ &\quad - F_{\eta,y}^{n_k(m):n_{k+1}(m)}(F_{\eta,y}^{m:n_k(m)}(\zeta''))\| \\ &\leq \frac{1}{2} \|F_{\eta,y}^{m:n_k(m)}(\zeta') - F_{\eta,y}^{m:n_k(m)}(\zeta'')\| \end{aligned}$$

for $m, k \geq 0$. Consequently, we have

$$\begin{aligned} &\|F_{\eta,y}^{m:n_k(m)}(\zeta') - F_{\eta,y}^{m:n_k(m)}(\zeta'')\| \\ &\leq \frac{1}{2^k} \|F_{\eta,y}^m(\zeta') - F_{\eta,y}^m(\zeta'')\| = \gamma_3^{n_k(m)-m} \|\zeta' - \zeta''\|. \end{aligned}$$

Combining this with Lemma 6.5 and (84) – (86), we get

$$\begin{aligned} &\|F_{\eta,y}^{m:n}(\zeta') - F_{\eta,y}^{m:n}(\zeta'')\| \\ &= \|F_{\eta,y}^{n_k(m):n}(F_{\eta,y}^{m:n_k(m)}(\zeta')) - F_{\eta,y}^{n_k(m):n}(F_{\eta,y}^{m:n_k(m)}(\zeta''))\| \\ &\leq C_5 \|F_{\eta,y}^{m:n_k(m)}(\zeta') - F_{\eta,y}^{m:n_k(m)}(\zeta'')\| \\ &\leq C_5\gamma_3^{n_k(m)-m} \|\zeta' - \zeta''\| \leq C_6\gamma_3^{n-m} \|\zeta' - \zeta''\| \end{aligned} \quad (93)$$

for $n_{k+1}(m) > n \geq n_k(m)$, $m, k \geq 0$. Here, we also use relations $C_5\gamma_3^{n_k(m)-m} = (C_5\gamma_3^{n_k(m)-n})\gamma_3^{n-m} \leq (C_5\gamma_3^{-n_0})\gamma_3^{n-m} = C_6\gamma_3^{n-m}$. Setting $k = \lfloor (n-m)/n_0 \rfloor$ in (93), we conclude that (83) holds for each $n \geq m \geq 0$ by \square .

VII. PROOF OF MAIN RESULTS

In this section, Theorems 2.1 and 2.2 are proved. The proofs of these theorems crucially depend on the results related to the kernels $S(z, dz')$, $S_\eta(z, dz')$ and the optimal filter $F_{\eta,y}^{m:n}(\zeta)$ (i.e., on Lemmas 5.1, 5.4, 6.6). As the properties of $S(z, dz')$, $S_\eta(z, dz')$ are very similar, the proofs of Theorems 2.1 and 2.2 have many elements in common. In order not to consider these elements twice (and to prove Theorems 2.1 and 2.2 as efficiently as possible), we introduce a new kernel $T_\eta(z, dz')$, where $\eta \in \mathbb{C}^d$, $z \in \mathcal{Z}$.¹ Its purpose is to capture all common features of $S(z, dz')$, $S_\eta(z, dz')$ which are relevant for the proof of Theorems 2.1 and 2.2. Using $T_\eta(z, dz')$, we recursively define kernels $\{T_\eta^n(z, dz')\}_{n \geq 0}$ by $T_\eta^0(z, B) = \delta_z(B)$ and

$$T_\eta^{n+1}(z, B) = \int T_\eta^n(z', B) T_\eta(z, dz'),$$

where $B \in \mathcal{B}(\mathcal{Z})$.

Regarding $T_\eta(z, dz')$, we assume the following.

Assumption 7.1. For each $\theta \in \Theta$, $z \in \mathcal{Z}$, $T_\theta(z, dz')$ is a probability measure.

Assumption 7.2. (i) There exist real numbers $\alpha \in (0, \delta]$, $L \in [1, \infty)$ such that

$$\begin{aligned} |T_\eta' - T_\eta''|(z, B) &\leq L\|\eta' - \eta''\|, \\ \int \tilde{\psi}(z') |T_\eta|(z, dz') &\leq L \end{aligned}$$

for all $\eta, \eta', \eta'' \in V_\alpha(\Theta)$, $z \in \mathcal{Z}$, $B \in \mathcal{B}(\mathcal{Z})$ (here, $|T_\eta' - T_\eta''|(z, dz')$ denotes the total variation of $T_\eta'(z, dz') - T_\eta''(z, dz')$, while $\delta, \tilde{\psi}(z)$ are specified in Assumption 2.2 and (12)).

(ii) For each $\eta \in V_\alpha(\Theta)$, there exists a complex measure $\tau_\eta(dz)$ such that $\lim_{n \rightarrow \infty} T_\eta^n(z, B) = \tau_\eta(B)$ for all $z \in \mathcal{Z}$, $B \in \mathcal{B}(\mathcal{Z})$.

(iii) There exists a real number $\beta \in (0, 1)$ such that

$$|T_\eta^n - \tau_\eta|(z, B) \leq L\beta^n$$

for all $\eta \in V_\alpha(\Theta)$, $z \in \mathcal{Z}$, $B \in \mathcal{B}(\mathcal{Z})$, $n \geq 1$ (here, $|T_\eta^n - \tau_\eta|(z, dz')$ stands for the total variation of $T_\eta^n(z, dz') - \tau_\eta(dz')$).

¹ $T_\eta(z, dz')$ can be considered as a mapping with the following properties: (i) $T_\eta(z, B)$ maps $\eta \in \mathbb{C}^d$, $z \in \mathcal{Z}$, $B \in \mathcal{B}(\mathcal{Z})$ to \mathbb{C} , (ii) $T_\eta(z, B)$ is measurable in (η, z) for each $B \in \mathcal{B}(\mathcal{Z})$, and (iii) $T_\eta(z, B)$ is a complex measure in B for each $\eta \in \mathbb{C}^d$, $z \in \mathcal{Z}$.

Remark. According to Lemmas 5.1 and 5.4, both kernels $S(z, dz')$, $S_\eta(z, dz')$ satisfy Assumptions 7.1 and 7.2. These assumptions capture all common properties of $S(z, dz')$, $S_\eta(z, dz')$ relevant for the proof of Theorems 2.1 and 2.2.

Besides the notations introduced in the previous sections, we rely here on the following notations, too. $u_\eta^n(z_{0:n})$ and $F_\eta^n(\xi, z_{1:n})$ are (respectively) the function and the complex measure defined by

$$\begin{aligned} u_\eta^n(z_{0:n}) &= u_\eta^n(x_{0:n}, y_{1:n}), \\ F_\eta^n(\xi, z_{1:n}) &= F_{\eta,y}^{0:n}(\xi) \end{aligned}$$

for $\eta \in \mathbb{C}^d$, $\xi \in \mathcal{M}_c(\mathcal{X})$, $x_0, \dots, x_n \in \mathcal{X}$, $y_0, \dots, y_n \in \mathcal{Y}$, $n \geq 0$ and $z_0 = (y_0, x_0), \dots, z_n = (y_n, x_n)$, where $\mathbf{y} = \{y'_n\}_{n \geq 1}$ is any sequence in \mathcal{Y} satisfying $y'_k = y_k$ for $n \geq k \geq 1$.² $\Phi_\eta(\xi, z)$ is the function defined by

$$\Phi_\eta(\xi, z) = \Phi_{\eta,y}(\xi),$$

where $x \in \mathcal{X}$, $y \in \mathcal{Y}$ and $z = (y, x)$ ($\Phi_{\eta,y}(\xi)$ is specified in (30)).³ $\Phi_\eta^n(\xi, z)$ is the function defined by

$$\begin{aligned} \Phi_\eta^n(\xi, z) &= \int \cdots \int \Phi_\eta \left(F_\eta^n(\xi, z_{1:n}), z_{n+1} \right) \\ &\quad \cdot T_\eta(z_n, dz_{n+1}) \cdots T_\eta(z, dz_1) \end{aligned} \quad (94)$$

for $n \geq 0$. $\bar{A}_\eta^n(\xi)$, $A_\eta^{k,n}(\xi, z)$, $B_\eta^n(\xi, z)$ are the functions defined by

$$\begin{aligned} \bar{A}_\eta^n(\xi) &= \int \cdots \int \int \left(\Phi_\eta \left(F_\eta^n(\xi, z_{1:n}), z_{n+1} \right) \right. \\ &\quad \left. - \Phi_\eta \left(F_\eta^{n-1}(\xi, z_{2:n}), z_{n+1} \right) \right) \\ &\quad \cdot T_\eta(z_n, dz_{n+1}) \cdots T_\eta(z_0, dz_1) \tau_\eta(dz_0), \\ A_\eta^{k,n}(\xi, z) &= \int \cdots \int \int \left(\Phi_\eta \left(F_\eta^{n-k+1}(\xi, z_{k:n}), z_{n+1} \right) \right. \\ &\quad \left. - \Phi_\eta \left(F_\eta^{n-k}(\xi, z_{k+1:n}), z_{n+1} \right) \right) \\ &\quad \cdot T_\eta(z_n, dz_{n+1}) \cdots T_\eta(z_k, dz_{k+1}) (T_\eta^k - \tau_\eta)(z, dz_k), \\ B_\eta^n(\xi, z) &= \int \Phi_\eta(\xi, z') (T_\eta^{n+1} - \tau_\eta)(z, dz') \end{aligned}$$

for $n \geq k \geq 1$.

Under the notations introduced above, we have

$$\log q_\theta^n(y_{1:n} | \lambda) = \sum_{k=0}^{n-1} \Phi_\theta \left(F_\theta^k(\lambda, z_{1:k}), z_{k+1} \right) \quad (95)$$

²Symbols $y_{1:0}$, $z_{1:0}$ denote empty sequences (i.e., sequences without any element). $u_\eta^n(x_{0:n}, y_{1:n})$, $F_{\eta,y}^{0:n}(\xi)$ are specified in (13),(38). $F_{\eta,y}^{0:n}(\xi)$ depends only on y'_1, \dots, y'_n and is independent of other elements of \mathbf{y} .

³Functions $u_\eta^n(z_{0:n})$, $F_\eta^n(\xi, z_{1:n})$, $\Phi_\eta(\xi, z)$ are just another notations for $u_\eta^n(x_{0:n}, y_{1:n})$, $F_{\eta,y}^{0:n}(\xi)$, $\Phi_{\eta,y}(\xi)$. However, notations $u_\eta^n(z_{0:n})$, $F_\eta^n(\xi, z_{1:n})$, $\Phi_\eta(\xi, z)$ are more suitable (than the original one) for measure-theoretic arguments which the analysis carried out in this section is based on.

for $\theta \in \Theta$, $\lambda \in \mathcal{P}(\mathcal{X})$, $x_1, \dots, x_n \in \mathcal{X}$, $y_1, \dots, y_n \in \mathcal{Y}$, $n \geq 1$ and $z_1 = (y_1, x_1), \dots, z_n = (y_n, x_n)$. We also have

$$\begin{aligned} \Phi_\eta^n(\xi, z') - \Phi_\eta^n(\xi, z'') &= \sum_{k=1}^n \left(A_\eta^{k,n}(\xi, z') - A_\eta^{k,n}(\xi, z'') \right) \\ &\quad + B_\eta^n(\xi, z') - B_\eta^n(\xi, z''), \end{aligned} \quad (96)$$

$$\begin{aligned} \Phi_\eta^{n+1}(\xi, z) - \Phi_\eta^n(\xi, z) &= \sum_{k=1}^{n+1} A_\eta^{k,n+1}(\xi, z) - \sum_{k=1}^n A_\eta^{k,n}(\xi, z) \\ &\quad + \bar{A}_\eta^{n+1}(\xi) + B_\eta^{n+1}(\xi, z) - B_\eta^n(\xi, z) \end{aligned} \quad (97)$$

for $\eta \in V_\alpha(\Theta)$, $\xi \in \mathcal{M}_c(\mathcal{X})$, $z, z', z'' \in \mathcal{Z}$, $n \geq 1$.

Lemma 7.1. Let Assumptions 2.1 – 2.3, 7.1 and 7.2 hold. Then, there exist a function ϕ_η mapping $\eta \in \mathbb{C}^d$ to \mathbb{C} and real numbers $\delta_9, \gamma_4 \in (0, 1)$, $C_7 \in [1, \infty)$ such that

$$\left| \Phi_\eta^n(\xi, z) - \phi_\eta \right| \leq C_7 n \gamma_4^n \quad (98)$$

for all $\eta \in V_{\delta_9}(\Theta)$, $\xi \in V_{\delta_9}(\mathcal{P}(\mathcal{X}))$, $z \in \mathcal{Z}$, $n \geq 1$.

Proof. Throughout the proof, the following notations is used. γ_4, δ_9 are the real numbers defined by $\gamma_4 = \max\{\beta^{1/2}, \gamma_3^{1/2}\}$, $\delta_9 = \min\{\delta_7, \delta_8, (1 - \gamma_4)/L\}$ ($\beta, \delta_7, \delta_8, \gamma_3, L$ are specified in Assumption 7.2 and Lemmas 6.5, 6.6). η is any element in $V_{\delta_9}(\Theta)$, while θ is any element of Θ satisfying $\|\eta - \theta\| < \delta_9$. ξ, ξ', ξ'' are any elements of $V_{\delta_9}(\mathcal{P}(\mathcal{X}))$, while z, z', z'' are any elements in \mathcal{Z} . B is any element of $\mathcal{B}(\mathcal{Z})$. n, k are any integers satisfying $n \geq k \geq 1$.

Owing to Assumptions 7.1 and 7.2, we have

$$\begin{aligned} |T_\eta|(z, B) &\leq T_\theta(z, B) + |T_\eta - T_\theta|(z, B) \\ &\leq 1 + L \|\eta - \theta\| \\ &< 1 + L \delta_9 \leq \frac{1}{\gamma_4} \end{aligned} \quad (99)$$

as $L \delta_9 \leq 1 - \gamma_4 \leq 1/\gamma_4 - 1$. Consequently, Assumption 7.1 yields

$$|\tau_\eta|(B) \leq |T_\eta - \tau_\eta|(z, B) + |T_\eta|(z, B) \leq L + \frac{1}{\gamma_4}. \quad (100)$$

Let $\tilde{C}_1 = 4C_4C_6$ where C_4, C_6 are specified in Lemmas 6.4, 6.6. Then, due to Lemmas 6.1, 6.4, 6.6, we have

$$\begin{aligned} &\left| \Phi_\eta \left(F_\eta^{n-k+1}(\xi, z_{k:n}), z_{n+1} \right) - \Phi_\eta \left(F_\eta^{n-k}(\xi, z_{k+1:n}), z_{n+1} \right) \right| \\ &\leq C_4 \tilde{\psi}(z_{n+1}) \left\| F_\eta^{n-k+1}(\xi, z_{k:n}) - F_\eta^{n-k}(\xi, z_{k+1:n}) \right\| \\ &= C_4 \tilde{\psi}(z_{n+1}) \left\| F_\eta^{n-k} \left(F_\eta^1(\xi, z_k), z_{k+1:n} \right) - F_\eta^{n-k}(\xi, z_{k+1:n}) \right\| \\ &\leq C_4 C_6 \gamma_3^{n-k} \tilde{\psi}(z_{n+1}) \left\| F_\eta^1(\xi, z_k) - \xi \right\| \\ &\leq \tilde{C}_1 \gamma_4^{2(n-k)} \tilde{\psi}(z_{n+1}) \end{aligned} \quad (101)$$

for $z_1, \dots, z_{n+1} \in \mathcal{Z}$.⁴ Similarly, owing to Lemmas 6.4, 6.6, we have

$$\begin{aligned} & \left| \Phi_\eta \left(F_\eta^n(\xi', z_{1:n}), z_{n+1} \right) - \Phi_\eta \left(F_\eta^n(\xi'', z_{1:n}), z_{n+1} \right) \right| \\ & \leq C_4 \tilde{\psi}(z_{n+1}) \left\| F_\eta^n(\xi', z_{1:n}) - F_\eta^n(\xi'', z_{1:n}) \right\| \\ & \leq C_4 C_6 \gamma_3^n \tilde{\psi}(z_{n+1}) \|\xi' - \xi''\| \\ & \leq \tilde{C}_1 \gamma_4^{2n} \tilde{\psi}(z_{n+1}). \end{aligned} \quad (102)$$

Let $\tilde{C}_2 = 2\tilde{C}_1 L^2 / \gamma_4^3$. Then, using Assumption 7.2 and (99), (102), we conclude

$$\begin{aligned} \left| \Phi_\eta^n(\xi', z) - \Phi_\eta^n(\xi'', z) \right| & \leq \tilde{C}_1 \gamma_4^{2n} \int \cdots \int \tilde{\psi}(z_{n+1}) \\ & \quad \cdot |T_\eta|(z_n, dz_{n+1}) \cdots |T_\eta|(z, dz_1) \\ & \leq \tilde{C}_1 L \gamma_4^n \leq \tilde{C}_2 \gamma_4^n. \end{aligned} \quad (103)$$

Similarly, relying on Assumption 7.2 and (99), (101), we deduce

$$\begin{aligned} \left| A_\eta^{k,n}(\xi, z) \right| & \leq \tilde{C}_1 \gamma_4^{2(n-k)} \int \cdots \int \tilde{\psi}(z_{n+1}) \\ & \quad \cdot |T_\eta|(z_n, dz_{n+1}) \cdots |T_\eta|(z_k, dz_{k+1}) \\ & \quad \cdot |T_\eta^k - \tau_\eta|(z, dz_k) \\ & \leq \tilde{C}_1 L^2 \beta^k \gamma_4^{n-k} \leq \tilde{C}_2 \gamma_4^n. \end{aligned} \quad (104)$$

Moreover, using Assumption 7.2 and (100), (101), we get

$$\begin{aligned} \left| \bar{A}_\eta^n(\xi) \right| & \leq \tilde{C}_1 \gamma_4^{2(n-1)} \int \cdots \int \tilde{\psi}(z_{n+1}) \\ & \quad \cdot |T_\eta|(z_n, dz_{n+1}) \cdots |T_\eta|(z_0, dz_1) |\tau_\eta|(dz_0) \\ & \leq \tilde{C}_1 L \left(L + \frac{1}{\gamma_4} \right) \gamma_4^{n-2} \leq \tilde{C}_2 \gamma_4^n. \end{aligned} \quad (105)$$

Let $\tilde{C}_3 = C_4 L^2$, $\tilde{C}_4 = 4(\tilde{C}_2 + \tilde{C}_3)$. Then, owing to Assumption 7.2 and Lemma 6.4, we have

$$\begin{aligned} \left| B_\eta^n(\xi, z) \right| & \leq C_4 \int \int \tilde{\psi}(z'') |T_\eta|(z', dz'') |T_\eta^n - \tau_\eta|(z, dz') \\ & \leq C_4 L^2 \beta^n \leq \tilde{C}_3 \gamma_4^n. \end{aligned} \quad (106)$$

Consequently, (97), (104), (105) yield

$$\begin{aligned} & \left| \Phi_\eta^{n+1}(\xi, z) - \Phi_\eta^n(\xi, z) \right| \\ & \leq \sum_{k=1}^{n+1} \left| A_\eta^{k,n+1}(\xi, z) \right| + \sum_{k=1}^n \left| A_\eta^{k,n}(\xi, z) \right| \\ & \quad + \left| \bar{A}_\eta^{n+1}(\xi) \right| + \left| B_\eta^{n+1}(\xi, z) \right| + \left| B_\eta^n(\xi, z) \right| \\ & \leq 2\tilde{C}_2(n+1)\gamma_4^n + 2\tilde{C}_3\gamma_4^n \leq \tilde{C}_4 n \gamma_4^n. \end{aligned} \quad (107)$$

⁴To get the first two relations in (101), use Lemmas 6.1, 6.4, and notice that inclusions $\eta \in V_{\delta_4}(\Theta)$, $F_\eta^{n-k+1}(\xi', z_{k:n}) \in V_{\delta_4}(\mathcal{P}(\mathcal{X}))$, $F_\eta^{n-k}(\xi'', z_{k+1:n}) \in V_{\delta_4}(\mathcal{P}(\mathcal{X}))$ follow from Lemma 6.6 and $\eta \in V_{\delta_9}(\Theta) \subseteq V_{\delta_5}(\Theta)$, $\xi', \xi'' \in V_{\delta_9}(\mathcal{P}(\mathcal{X})) \subseteq V_{\delta_6}(\mathcal{P}(\mathcal{X}))$. To get the third relation in (101), use Lemma 6.6 and notice that $F_\eta^1(\xi, z_k) \in V_{\delta_6}(\mathcal{P}(\mathcal{X}))$ follows from Lemma 6.4 and $\eta \in V_{\delta_9}(\Theta) \subseteq V_{\delta_7}(\Theta)$, $\xi \in V_{\delta_9}(\mathcal{P}(\mathcal{X})) \subseteq V_{\delta_8}(\mathcal{P}(\mathcal{X}))$. To get the last relation in (101), use inequality $\|F_\eta^1(\xi, z_k) - \xi\| \leq \|F_\eta^1(\xi, z_k)\| + \|\xi\| \leq 2 + \delta_6 + \delta_9 \leq 4$.

Hence, we have

$$\begin{aligned} & \sum_{n=1}^{\infty} \left| \Phi_\eta^{n+1}(\xi, z) - \Phi_\eta^n(\xi, z) \right| \\ & \leq \tilde{C}_4 \sum_{n=1}^{\infty} n \gamma_4^n \leq \frac{\tilde{C}_4}{(1-\gamma_4)^2} < \infty. \end{aligned} \quad (108)$$

Now, combining (96), (104), (106), we get

$$\begin{aligned} \left| \Phi_\eta^n(\xi, z') - \Phi_\eta^n(\xi, z'') \right| & \leq \sum_{k=1}^n \left| A_\eta^{k,n}(\xi, z') \right| + \sum_{k=1}^n \left| A_\eta^{k,n}(\xi, z'') \right| \\ & \quad + \left| B_\eta^n(\xi, z') \right| + \left| B_\eta^n(\xi, z'') \right| \\ & \leq 2\tilde{C}_2 n \gamma_4^n + 2\tilde{C}_3 \gamma_4^n. \end{aligned}$$

Then, (103) implies

$$\begin{aligned} \left| \Phi_\eta^n(\xi', z') - \Phi_\eta^n(\xi'', z'') \right| & \leq \left| \Phi_\eta^n(\xi', z') - \Phi_\eta^n(\xi'', z') \right| \\ & \quad + \left| \Phi_\eta^n(\xi'', z') - \Phi_\eta^n(\xi'', z'') \right| \\ & \leq \tilde{C}_2(2n+1)\gamma_4^n + 2\tilde{C}_3\gamma_4^n \leq \tilde{C}_4 n \gamma_4^n. \end{aligned} \quad (109)$$

Let $C_7 = \tilde{C}_4 / (1-\gamma_4)^2$. Moreover, let

$$\phi_\eta(\xi, z) = \Phi_\eta^0(\xi, z) + \sum_{n=0}^{\infty} \left(\Phi_\eta^{n+1}(\xi, z) - \Phi_\eta^n(\xi, z) \right).$$

Then, due to (108), $\phi_\eta(\xi, z)$ is well-defined. Now, (107) implies

$$\begin{aligned} \left| \Phi_\eta^n(\xi, z) - \phi_\eta(\xi, z) \right| & \leq \sum_{k=n}^{\infty} \left| \Phi_\eta^{k+1}(\xi, z) - \Phi_\eta^k(\xi, z) \right| \\ & \leq \tilde{C}_4 \sum_{k=n}^{\infty} k \gamma_4^k \leq C_7 n \gamma_4^n. \end{aligned} \quad (110)$$

Consequently, (109) yields

$$\begin{aligned} \left| \phi_\eta(\xi', z') - \phi_\eta(\xi'', z'') \right| & \leq \left| \Phi_\eta^n(\xi', z') - \Phi_\eta^n(\xi'', z'') \right| \\ & \quad + \left| \Phi_\eta^n(\xi', z') - \phi_\eta(\xi', z') \right| \\ & \quad + \left| \Phi_\eta^n(\xi'', z'') - \phi_\eta(\xi'', z'') \right| \\ & \leq 3C_7 n \gamma_4^n. \end{aligned}$$

Therefore, $\phi_\eta(\xi', z') = \phi_\eta(\xi'', z'')$ for any $\xi', \xi'' \in V_{\delta_9}(\mathcal{P}(\mathcal{X}))$, $z', z'' \in \mathcal{Z}$. Hence, there exists a function ϕ_η which maps $\eta \in \mathbb{C}^d$ to \mathbb{C} and satisfies $\phi_\eta = \phi_\eta(\xi, z)$ for all $\eta \in V_{\delta_9}(\Theta)$, $\xi \in V_{\delta_9}(\mathcal{P}(\mathcal{X}))$, $z \in \mathcal{Z}$. Then, using (110), we conclude that (98) holds for $\eta \in V_{\delta_9}(\Theta)$, $\xi \in V_{\delta_9}(\mathcal{P}(\mathcal{X}))$, $z \in \mathcal{Z}$. \square

Lemma 7.2. (i) Let Assumptions 2.1 – 2.3 and 2.5 hold. Then, integral

$$\int \cdots \int \Phi_\eta \left(F_\eta^n(\lambda, z_{1:n}), z_{n+1} \right) S(z_n, dz_{n+1}) \cdots S(z, dz_1) \quad (111)$$

is analytic in η for all $\eta \in V_{\delta_5}(\Theta)$, $\lambda \in \mathcal{P}(\mathcal{X})$, $z \in \mathcal{Z}$, $n \geq 1$ (δ_5 is specified in Lemmas 6.5, 6.6).

(ii) Let Assumptions 2.1 – 2.4 hold. Then, integral

$$\int \cdots \int \Phi_\eta \left(F_\eta^n(\lambda, z_{1:n}), z_{n+1} \right) S_\eta(z_n, dz_{n+1}) \cdots S_\eta(z, dz_1) \quad (112)$$

is analytic in η for all $\eta \in V_{\delta_5}(\Theta)$, $\lambda \in \mathcal{P}(\mathcal{X})$, $z \in \mathcal{Z}$, $n \geq 1$.

Proof. Throughout the proof, the following notations is used. $\tilde{\phi}(z)$ is the function defined by $\tilde{\phi}(z) = \phi(y)$ for $x \in \mathcal{X}$, $y \in \mathcal{Y}$ and $z = (y, x)$. η is any element of $V_{\delta_5}(\Theta)$, while λ is any element in $\mathcal{P}(\mathcal{X})$. $\{x_n\}_{n \geq 0}$, $\{y_n\}_{n \geq 0}$ are any sequences in \mathcal{X} , \mathcal{Y} (respectively), while $\{z_n\}_{n \geq 0}$ is the sequence defined by $z_n = (y_n, x_n)$ for $n \geq 0$ (notice that $\{z_n\}_{n \geq 0}$ can be any sequence in \mathcal{Z}). $n \geq 1$ is any integer.

Using Lemmas 6.1, 6.6, we conclude

$$\begin{aligned} \Phi_\eta \left(F_\eta^n(\lambda, z_{1:n}), z_{n+1} \right) &= \Phi_{\eta, y_{n+1}} \left(F_{\eta, y}^{0:n}(\lambda) \right) \\ &= \log \left\langle R_{\eta, y}^{n:n+1} \left(F_{\eta, y}^{0:n}(\lambda) \right) \right\rangle \\ &= \log \left(\frac{\left\langle R_{\eta, y}^{0:n+1}(\lambda) \right\rangle}{\left\langle R_{\eta, y}^{0:n}(\lambda) \right\rangle} \right), \end{aligned}$$

where $\mathbf{y} = \{y'_k\}_{k \geq 1}$ is any sequence in \mathcal{Y} satisfying $y'_k = y_k$ for $1 \leq k \leq n+1$. Combining this with Lemmas 5.3, 6.3, 6.6, we deduce that $\Phi_\eta \left(F_\eta^n(\lambda, z_{1:n}), z_{n+1} \right)$, $u_\eta^n(z_{0:n})$ are analytic in η for each $\eta \in V_{\delta_5}(\Theta)$. Moreover, due to Lemmas 5.3, 6.4, 6.6, we have

$$\left| \Phi_\eta \left(F_\eta^n(\lambda, z_{1:n}), z_{n+1} \right) \right| \leq C_4 \tilde{\psi}(z_{n+1}), \quad (113)$$

$$\left| u_\eta^n(z_{0:n}) \right| \leq K_n \prod_{k=1}^n \tilde{\phi}(z_k) \quad (114)$$

($\tilde{\psi}(z)$ is specified in (12)).

Owing to Assumption 2.5, we have

$$\begin{aligned} &\int \cdots \int \tilde{\psi}(z_{n+1}) S(z_n, dz_{n+1}) \cdots S(z_0, dz_1) \\ &= \int \int \cdots \int (1 + \psi(y_{n+1})) Q(x_{n+1}, dy_{n+1}) \\ &\quad \cdot P(x_n, dx_{n+1}) \cdots P(x_0, dx_1) \\ &\leq K + 1 < \infty. \end{aligned}$$

Consequently, Lemma A1.1 (see Appendix 1) and (113) imply that integral (111) is analytic in η for each $\eta \in V_{\delta_5}(\Theta)$.

Relying on (16), it is easy to show

$$\begin{aligned} &\int \cdots \int \Phi_\eta \left(F_\eta^n(\lambda, z_{1:n}), z_{n+1} \right) S_\eta(z_n, dz_{n+1}) \cdots S_\eta(z_0, dz_1) \\ &= \int \cdots \int \Phi_\eta \left(F_\eta^n(\lambda, z_{1:n}), z_{n+1} \right) u_\eta^{n+1}(z_{0:n+1}) \\ &\quad \cdot (v \times \mu)(dz_{n+1}) \cdots (v \times \mu)(dz_1). \end{aligned}$$

Moreover, due to Assumptions 2.2, 2.4, we have

$$\begin{aligned} &\int \cdots \int \tilde{\psi}(z_{n+1}) \left(\prod_{k=1}^{n+1} \tilde{\phi}(z_k) \right) \\ &\quad \cdot (v \times \mu)(dz_{n+1}) \cdots (v \times \mu)(dz_1) \\ &= \|\mu\|^{n+1} \left(\int (1 + \psi(y_{n+1})) \phi(y_{n+1}) v(dy_{n+1}) \right) \\ &\quad \cdot \left(\prod_{k=1}^n \int \phi(y_k) v(dy_k) \right) \\ &< \infty. \end{aligned}$$

Consequently, Lemma A1.1 (see Appendix 1) and (113), (114) imply that integral (112) is analytic in η for $\eta \in V_{\delta_5}(\Theta)$. \square

Proof of Theorem 2.1. Let $T_\eta(z, dz')$ be the kernel defined by $T_\eta(z, B) = S(z, B)$ for $\eta \in \mathbb{C}^d$, $z \in \mathcal{Z}$, $B \in \mathcal{B}(\mathcal{Z})$ ($S(z, dz')$ is specified in (14)). Moreover, let $T_\eta^n(z, dz')$, $\Phi_\eta^n(\lambda, z)$ have the same meaning as in (94). Then, owing to Lemma 7.2, $\Phi_\eta^n(\lambda, z)$ is analytic in η for each $\eta \in V_{\delta_5}(\Theta)$, $\lambda \in \mathcal{P}(\mathcal{X})$, $z \in \mathcal{Z}$, $n \geq 1$. Moreover, due to Lemma 5.1, kernel $T_\eta(z, dz')$ (defined here) satisfies Assumptions 7.1, 7.2. Combining this with Lemma 7.1, we deduce that there exist a function ϕ_η mapping $\eta \in \mathbb{C}^d$ to \mathbb{C} and real numbers $\delta_\theta \in (0, \delta_5]$, $\gamma_4 \in (0, 1)$, $C_7 \in [1, \infty)$ such that (98) holds for $\eta \in V_{\delta_\theta}(\Theta)$, $\lambda \in \mathcal{P}(\mathcal{X})$, $z \in \mathcal{Z}$, $n \geq 1$. Since the limit of uniformly convergent analytic functions is also analytic (see e.g., [28, Theorem 2.4.1]), ϕ_η is analytic in η for each $\eta \in V_{\delta_\theta}(\Theta)$.

In what follows in the proof, θ, λ, z are any elements of Θ , $\mathcal{P}(\mathcal{X})$, \mathcal{Z} (respectively), while $n \geq 1$ is any integer. It is straightforward to verify

$$\Phi_\theta^n(\lambda, z) = E \left(\Phi_\theta \left(F_\theta^n(\lambda, Z_{1:n}), Z_{n+1} \right) \mid Z_0 = z \right),$$

where $Z_n = (Y_n, X_n)$. Therefore, (95) yields

$$E \left(\log q_\theta^n(Y_{1:n} | \lambda) \right) = \sum_{k=0}^{n-1} E \left(\Phi_\theta^k(\lambda, Z_0) \right).$$

Then, Lemma 7.1 implies

$$\begin{aligned} \left| E \left(\frac{1}{n} \log q_\theta(Y_{1:n} | \lambda) \right) - \phi_\theta \right| &\leq \frac{1}{n} \sum_{k=0}^{n-1} E \left| \Phi_\theta^k(\lambda, Z_0) - \phi_\theta \right| \\ &\leq \frac{C_7}{n} \sum_{k=0}^{n-1} \gamma_4^k \leq \frac{C_7}{n(1 - \gamma_4)}. \end{aligned}$$

Consequently, there exists a function $l : \Theta \rightarrow \mathbb{R}$ with the properties specified in the statement of the theorem. \square

Proof of Theorem 2.2. Let $T_\eta(z, dz')$ be the kernel defined by $T_\eta(z, B) = S_\eta(z, B)$ for $\eta \in \mathbb{C}^d$, $z \in \mathcal{Z}$, $B \in \mathcal{B}(\mathcal{Z})$ ($S_\eta(z, dz')$ is specified in (15)). Moreover, let $T_\eta^n(z, dz')$, $\Phi_\eta^n(\lambda, z)$ have the same meaning as in (94). Then, due to Lemma 7.2, $\Phi_\eta^n(\lambda, z)$ is analytic in η for each $\eta \in V_{\delta_5}(\Theta)$, $\lambda \in \mathcal{P}(\mathcal{X})$, $z \in \mathcal{Z}$, $n \geq 1$. Moreover, Lemma 5.4 implies that Assumptions 7.1, 7.2 hold for kernel $T_\eta(z, dz')$ (defined here). Combining this with Lemma 7.1, we conclude that there

exist a function ϕ_η mapping $\eta \in \mathbb{C}^d$ to \mathbb{C} and real numbers $\delta_9 \in (0, \delta_5]$, $\gamma_4 \in (0, 1)$, $C_7 \in [1, \infty)$ such that (98) holds for $\eta \in V_{\delta_9}(\Theta)$, $\lambda \in \mathcal{P}(\mathcal{X})$, $z \in \mathcal{Z}$, $n \geq 1$. As the limit of uniformly convergent analytic functions is also analytic (see e.g., [28, Theorem 2.4.1]), ϕ_η is analytic in η for each $\eta \in V_{\delta_9}(\Theta)$.

In the rest of the proof, θ , λ , z are any elements of Θ , $\mathcal{P}(\mathcal{X})$, \mathcal{Z} (respectively), while $n \geq 1$ is any integer. It is easy to show

$$\Phi_\theta^n(\lambda, z) = E \left(\Phi_\theta \left(F_\theta^n(\lambda, Z_{1:n}^{\theta, \lambda}, Z_{n+1}^{\theta, \lambda}) \middle| Z_0^{\theta, \lambda} = z \right) \right),$$

where $Z_n^{\theta, \lambda} = (Y_{1:n}^{\theta, \lambda}, X_n^{\theta, \lambda})$. Then, (95) yields

$$E \left(\log q_\theta^n(Y_{1:n}^{\theta, \lambda} | \lambda) \right) = \sum_{k=0}^{n-1} E \left(\Phi_\theta^k(\lambda, Z_0^{\theta, \lambda}) \right).$$

Therefore, Lemma 7.1 implies

$$\begin{aligned} \left| E \left(\frac{1}{n} \log q_\theta(Y_{1:n}^{\theta, \lambda} | \lambda) \right) - \phi_\theta \right| &\leq \frac{1}{n} \sum_{k=0}^{n-1} E \left| \Phi_\theta^k(\lambda, Z_0^{\theta, \lambda}) - \phi_\theta \right| \\ &\leq \frac{C_7}{n} \sum_{k=0}^{n-1} \gamma_4^k \leq \frac{C_7}{n(1-\gamma_4)}. \end{aligned}$$

Consequently, there exists a function $h : \Theta \rightarrow \mathbb{R}$ with the properties specified in the statement of the theorem. \square

VIII. PROOF OF COROLLARIES 3.1 – 4.2

Proof of Corollaries 3.1 and 3.2. Let $\tilde{\Theta}$ be any non-empty bounded open set satisfying $\text{cl}\tilde{\Theta} \subset \Theta$. As $\text{cl}\tilde{\Theta}$, \mathcal{X} are compact sets, Assumption 3.2 and Lemma A1.2 (see Appendix 1) imply that there exist functions $\{\hat{a}_\eta^i(x)\}_{1 \leq i \leq N_x}$, $\{\hat{b}_\eta^j(x)\}_{1 \leq j \leq N_y}$ with the following properties:

- (i) $\{\hat{a}_\eta^i(x)\}_{1 \leq i \leq N_x}$, $\{\hat{b}_\eta^j(x)\}_{1 \leq j \leq N_y}$ map $\eta \in \mathbb{C}^d$, $x \in \mathbb{C}^{d_x}$ to \mathbb{C} .
- (ii) $\hat{a}_\theta^i(x) = a_\theta^i(x)$, $\hat{b}_\theta^j(x) = b_\theta^j(x)$ for $\theta \in \tilde{\Theta}$, $x \in \mathcal{X}$, $1 \leq i \leq N_x$, $1 \leq j \leq N_y$.
- (iii) There exists a real number $\alpha_1 \in (0, 1)$ such that $\hat{a}_\eta^i(x)$, $\hat{b}_\eta^j(x)$ are analytic in (η, x) for $\eta \in V_{\alpha_1}(\tilde{\Theta})$, $x \in V_{\alpha_1}(\mathcal{X})$, $1 \leq i \leq N_x$, $1 \leq j \leq N_y$.

Owing to Assumption 3.2, $\{\hat{a}_\theta^i(x)\}_{1 \leq i \leq N_x}$, $\{\hat{b}_\theta^j(x)\}_{1 \leq j \leq N_y}$ are positive and uniformly bounded away from zero for $\theta \in \text{cl}\tilde{\Theta}$, $x \in \mathcal{X}$. Then, due to (iii), there exist real numbers $\alpha \in (0, \alpha_1)$, $\beta \in (0, 1)$ such that

$$\text{Re} \left\{ \hat{a}_\eta^i(x) \right\} \geq \beta, \quad |\hat{a}_\eta^i(x)| \leq \frac{1}{\beta}, \quad (115)$$

$$\text{Re} \left\{ \hat{b}_\eta^j(x) \right\} \geq \beta, \quad |\hat{b}_\eta^j(x)| \leq \frac{1}{\beta} \quad (116)$$

for $\eta \in V_\alpha(\tilde{\Theta})$, $x \in V_\alpha(\mathcal{X})$, $1 \leq i \leq N_x$, $1 \leq j \leq N_y$.

Let $\hat{p}_\eta(x'|x)$, $\hat{q}_\eta(y|x)$ be the functions defined by

$$\begin{aligned} \hat{p}_\eta(x'|x) &= \sum_{i=1}^{N_x} \hat{a}_\eta^i(x) v_i(x'), \\ \hat{q}_\eta(y|x) &= \sum_{j=1}^{N_y} \hat{b}_\eta^j(x) w_j(y) \end{aligned}$$

for $\eta \in \mathbb{C}^d$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$, while

$$\begin{aligned} r_\theta(y, x'|x) &= q_\theta(y|x') p_\theta(x'|x), \\ \hat{r}_\eta(y, x'|x) &= \hat{q}_\eta(y|x') \hat{p}_\eta(x'|x) \end{aligned}$$

for the same η, x, x', y and $\theta \in \Theta$. Then, owing to (ii), (iii), $\hat{r}_\eta(y, x'|x)$ is analytic in η for each $\eta \in V_\alpha(\tilde{\Theta})$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$. For similar reasons, $\hat{r}_\theta(y, x'|x) = r_\theta(y, x'|x)$ for the same x, x', y and $\theta \in \tilde{\Theta}$. Moreover, Assumption 3.3 and (115) imply

$$|\hat{p}_\eta(x'|x)| \geq \sum_{i=1}^{N_x} \text{Re} \left\{ \hat{a}_\eta^i(x) \right\} v_i(x') \geq \beta \varepsilon N_x, \quad (117)$$

$$|\hat{p}_\eta(x'|x)| \leq \sum_{i=1}^{N_x} |\hat{a}_\eta^i(x)| v_i(x') \leq \frac{N_x}{\beta \varepsilon} \quad (118)$$

for $\eta \in V_\alpha(\tilde{\Theta})$, $x, x' \in \mathcal{X}$. Similarly, (116) yields

$$|\hat{q}_\eta(y|x)| \geq \sum_{j=1}^{N_y} \text{Re} \left\{ \hat{b}_\eta^j(x) \right\} w_j(y) \geq \beta \sum_{j=1}^{N_y} w_j(y), \quad (119)$$

$$|\hat{q}_\eta(y|x)| \leq \sum_{j=1}^{N_y} |\hat{b}_\eta^j(x)| w_j(y) \leq \frac{1}{\beta} \sum_{j=1}^{N_y} w_j(y) \quad (120)$$

for the same η, x and $y \in \mathcal{Y}$.

Let $\tilde{C}_1 = \beta^{-2} \varepsilon^{-1} N_x$, $\tilde{C}_2 = \tilde{C}_1 N_x$, $\gamma = \beta^4 \varepsilon^2$. Moreover, let $\phi(y)$, $\psi(y)$ be the functions defined by

$$\begin{aligned} \phi(y) &= \tilde{C}_1 \sum_{j=1}^{N_y} w_j(y), \\ \psi(y) &= \tilde{C}_2 \left(1 + \sum_{j=1}^{N_y} |\log w_j(y)| \right) \end{aligned}$$

for $y \in \mathcal{Y}$. Then, combining (117) – (120), we get

$$\gamma \phi(y) \leq |\hat{r}_\eta(y, x'|x)| \leq \frac{\phi(y)}{\gamma} \quad (121)$$

for $\eta \in V_\alpha(\tilde{\Theta})$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$. We also get

$$\log \phi(y) \leq \log(\tilde{C}_1 N_x) + \max_{1 \leq j \leq N_y} \log w_j(y)$$

$$\leq \tilde{C}_1 N_x \left(1 + \sum_{j=1}^{N_y} |\log w_j(y)| \right),$$

$$\log \phi(y) \geq \log(\tilde{C}_1 N_x) + \min_{1 \leq j \leq N_y} \log w_j(y)$$

$$\geq -\tilde{C}_1 N_x \left(1 + \sum_{j=1}^{N_y} |\log w_j(y)| \right).$$

Therefore, we have

$$|\log \phi(y)| \leq \psi(y). \quad (122)$$

Since $\int \phi(y) \nu(dy) = \tilde{C}_1 N_y < \infty$, (121), (122) imply that Assumptions 2.1 – 2.3 follow from Assumptions 4.1 – 4.3 when Θ is restricted to $\tilde{\Theta}$ (i.e., when Θ is replaced with $\tilde{\Theta}$).

Owing to Assumption 3.5, we have

$$\begin{aligned} \int \psi(y) Q(x, dy) &= \tilde{C}_2 + \sum_{j=1}^{N_y} |\log w_j(y)| Q(x, dy) \\ &\leq \tilde{C}_2 + KN_y < \infty. \end{aligned}$$

Hence, Assumption 2.5 results from Assumption 3.5. Moreover, due to Assumption 3.4, we have

$$\begin{aligned} \int \psi(y) \phi(y) \nu(dy) &= \tilde{C}_1 \tilde{C}_2 \sum_{j,k=1}^{N_y} \int |\log w_j(y)| w_k(y) \nu(dy) \\ &+ \tilde{C}_1 \tilde{C}_2 N_y < \infty. \end{aligned}$$

Thus, Assumption 2.4 results from Assumption 3.4.

Using Theorems 2.1, 2.2, we conclude that there exist functions $\tilde{l}, \tilde{h}: \tilde{\Theta} \rightarrow \mathbb{R}$ such that $\tilde{l}(\theta), \tilde{h}(\theta)$ are real-analytic in θ and satisfy $\lim_{n \rightarrow \infty} l_n(\theta, \lambda) = \tilde{l}(\theta)$, $\lim_{n \rightarrow \infty} h_n(\theta, \lambda) = \tilde{h}(\theta)$ for each $\theta \in \tilde{\Theta}$, $\lambda \in \mathcal{P}(\mathcal{X})$ ($l_n(\theta, \lambda), h_n(\theta, \lambda)$ have the same meaning as in (1)). Consequently, Corollaries 4.1, 4.2 hold. We use here the representation $\Theta = \bigcup_{n=1}^{\infty} \tilde{\Theta}_n$, where $\{\tilde{\Theta}_n\}_{n \geq 1}$ is a sequence of non-empty open balls satisfying $\text{cl}\tilde{\Theta}_n \subset \Theta$ for $n \geq 1$. \square

Proof of Corollaries 4.1 and 4.2. Let $\tilde{\Theta}$ be a non-empty bounded open set satisfying $\text{cl}\tilde{\Theta} \subset \Theta$. As $\text{cl}\tilde{\Theta}, \mathcal{X}, \mathcal{Y}$ are compact and $A_\theta(x), B_\theta^{-1}(x), C_\theta(x), D_\theta^{-1}(x)$ are continuous in (θ, x) , it follows from Assumption 4.3 that there exists a real number $r \in [1, \infty)$ such that

$$\|B_\theta^{-1}(x)(x' - A_\theta(x))\| \leq r, \quad (123)$$

$$\|D_\theta^{-1}(x)(y - C_\theta(x))\| \leq r \quad (124)$$

for $\theta \in \text{cl}\tilde{\Theta}$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$.

Let $\tilde{\mathcal{X}} = \{x \in \mathbb{R}^{d_x} : \|x\| \leq r\}$, $\tilde{\mathcal{Y}} = \{y \in \mathbb{R}^{d_y} : \|y\| \leq r\}$. Since $\text{cl}\tilde{\Theta}, \mathcal{X}, \tilde{\mathcal{X}}, \tilde{\mathcal{Y}}$ are compact sets, Assumptions 4.2, 4.3 and Lemma A1.2 (see Appendix 1) imply that there exist functions $\hat{A}_\eta(x), \hat{B}_\eta(x), \hat{C}_\eta(x), \hat{D}_\eta(x)$ and $\hat{v}(x), \hat{w}(y)$ with the following properties:

(i) $\hat{A}_\eta(x), \hat{B}_\eta(x), \hat{C}_\eta(x), \hat{D}_\eta(x)$ map $\eta \in \mathbb{C}^d$, $x \in \mathbb{C}^{d_x}$ to $\mathbb{C}^{d_x}, \mathbb{C}^{d_x \times d_x}, \mathbb{C}^{d_y}, \mathbb{C}^{d_y \times d_y}$ (respectively), while $\hat{v}(x), \hat{w}(y)$ map $x \in \mathbb{C}^{d_x}, y \in \mathbb{C}^{d_y}$ to \mathbb{C} .

(ii) $\hat{A}_\theta(x) = A_\theta(x), \hat{B}_\theta(x) = B_\theta(x), \hat{C}_\theta(x) = C_\theta(x), \hat{D}_\theta(x) = D_\theta(x)$ for $\theta \in \tilde{\Theta}$, $x \in \mathcal{X}$, and $\hat{v}(x) = v(x), \hat{w}(y) = w(y)$ for $x \in \tilde{\mathcal{X}}, y \in \tilde{\mathcal{Y}}$.

(iii) There exists a real number $\alpha_1 \in (0, 1)$ such that $\hat{A}_\eta(x), \hat{B}_\eta(x), \hat{C}_\eta(x), \hat{D}_\eta(x)$ are analytic in (η, x) for $\eta \in V_{\alpha_1}(\tilde{\Theta})$, $x \in V_{\alpha_1}(\mathcal{X})$.

(iv) There exists a real number $\alpha_2 \in (0, 1)$ such that $\hat{v}(x), \hat{w}(y)$ are analytic in x, y (respectively) for $x \in V_{\alpha_2}(\tilde{\mathcal{X}}), y \in V_{\alpha_2}(\tilde{\mathcal{Y}})$.

Since $|\det \hat{B}_\theta(x)|, |\det \hat{D}_\theta(x)|$ are uniformly bounded away from zero for $\theta \in \text{cl}\tilde{\Theta}$, $x \in \mathcal{X}$, Assumption 4.3 and (iii) imply that there exists a real number $\alpha_3 \in (0, \alpha_1)$ such that $\det \hat{B}_\eta(x) \neq 0, \det \hat{D}_\eta(x) \neq 0$ for $\eta \in V_{\alpha_3}(\tilde{\Theta})$,

$x \in V_{\alpha_3}(\mathcal{X})$. Therefore,

$$\begin{aligned} \hat{B}_\eta^{-1}(x)(x' - \hat{A}_\eta(x)), \\ \hat{D}_\eta^{-1}(x)(y - \hat{C}_\eta(x)) \end{aligned}$$

are well-defined and analytic in (η, x, x', y) for $\eta \in V_{\alpha_3}(\tilde{\Theta})$, $x \in V_{\alpha_3}(\mathcal{X})$, $x' \in \mathbb{C}^{d_x}$, $y \in \mathbb{C}^{d_y}$. As $\text{cl}\tilde{\Theta}, \mathcal{X}, \mathcal{Y}$ are compact sets, it follows from (123), (124) that there exists a real number $\alpha_4 \in (0, \alpha_3)$ such that

$$\begin{aligned} \left\| \text{Re} \left\{ \hat{B}_\eta^{-1}(x)(x' - \hat{A}_\eta(x)) \right\} \right\| &< r + \frac{\alpha_2}{2}, \\ \left\| \text{Im} \left\{ \hat{B}_\eta^{-1}(x)(x' - \hat{A}_\eta(x)) \right\} \right\| &< \frac{\alpha_2}{2}, \\ \left\| \text{Re} \left\{ \hat{D}_\eta^{-1}(x)(y - \hat{C}_\eta(x)) \right\} \right\| &< r + \frac{\alpha_2}{2}, \\ \left\| \text{Im} \left\{ \hat{D}_\eta^{-1}(x)(y - \hat{C}_\eta(x)) \right\} \right\| &< \frac{\alpha_2}{2} \end{aligned}$$

for $\eta \in V_{\alpha_4}(\tilde{\Theta})$, $x, x' \in V_{\alpha_4}(\mathcal{X})$, $y \in V_{\alpha_4}(\mathcal{Y})$. Hence, we have

$$\begin{aligned} \hat{B}_\eta^{-1}(x)(x' - \hat{A}_\eta(x)) &\in V_{\alpha_2}(\tilde{\mathcal{X}}), \\ \hat{D}_\eta^{-1}(x)(y - \hat{C}_\eta(x)) &\in V_{\alpha_2}(\tilde{\mathcal{Y}}) \end{aligned}$$

for the same η, x, x', y . Consequently, (iv) yields that

$$\hat{v} \left(\hat{B}_\eta^{-1}(x)(x' - \hat{A}_\eta(x)) \right), \quad (125)$$

$$\hat{w} \left(\hat{D}_\eta^{-1}(x)(y - \hat{C}_\eta(x)) \right) \quad (126)$$

are analytic in (η, x, x', y) for $\eta \in V_{\alpha_4}(\tilde{\Theta})$, $x, x' \in V_{\alpha_4}(\mathcal{X})$, $y \in V_{\alpha_4}(\mathcal{Y})$. Since functions (125), (126) are positive and uniformly bounded away from zero for $\eta \in \text{cl}\tilde{\Theta}$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$, Assumption 4.2 implies that there exist real numbers $\alpha \in (0, \alpha_4)$, $\beta \in (0, 1)$ such that

$$\text{Re} \left\{ \hat{v} \left(\hat{B}_\eta^{-1}(x)(x' - \hat{A}_\eta(x)) \right) \right\} \geq \beta, \quad (127)$$

$$\left| \hat{v} \left(\hat{B}_\eta^{-1}(x)(x' - \hat{A}_\eta(x)) \right) \right| \leq \frac{1}{\beta}, \quad (128)$$

$$\text{Re} \left\{ \hat{w} \left(\hat{D}_\eta^{-1}(x)(y - \hat{C}_\eta(x)) \right) \right\} \geq \beta, \quad (129)$$

$$\left| \hat{w} \left(\hat{D}_\eta^{-1}(x)(y - \hat{C}_\eta(x)) \right) \right| \leq \frac{1}{\beta} \quad (130)$$

for $\eta \in V_\alpha(\tilde{\Theta})$, $x, x' \in V_\alpha(\mathcal{X})$, $y \in V_\alpha(\mathcal{Y})$.

Owing to (127), (128), we have

$$\begin{aligned} \left| \int_{\mathcal{X}} \hat{v} \left(\hat{B}_\eta^{-1}(x)(x' - \hat{A}_\eta(x)) \right) dx' \right| \\ \geq \int_{\mathcal{X}} \text{Re} \left\{ \hat{v} \left(\hat{B}_\eta^{-1}(x)(x' - \hat{A}_\eta(x)) \right) \right\} dx' \geq \beta m(\mathcal{X}) > 0, \end{aligned} \quad (131)$$

$$\begin{aligned} \left| \int_{\mathcal{X}} \hat{v} \left(\hat{B}_\eta^{-1}(x)(x' - \hat{A}_\eta(x)) \right) dx' \right| \\ \leq \int_{\mathcal{X}} \left| \hat{v} \left(\hat{B}_\eta^{-1}(x)(x' - \hat{A}_\eta(x)) \right) \right| dx' \leq \frac{m(\mathcal{X})}{\beta} \end{aligned} \quad (132)$$

for $\eta \in V_\alpha(\tilde{\Theta})$, $x \in V_\alpha(\mathcal{X})$, where $m(\mathcal{X})$ is the Lebesgue measure of \mathcal{X} . Similarly, due to (129), (130), we have

$$\begin{aligned} & \left| \int_{\mathcal{Y}} \hat{w} \left(\hat{D}_\eta^{-1}(x) \left(y - \hat{C}_\eta(x) \right) \right) dy \right| \\ & \geq \int_{\mathcal{Y}} \operatorname{Re} \left\{ \hat{w} \left(\hat{D}_\eta^{-1}(x) \left(y - \hat{C}_\eta(x) \right) \right) \right\} dy \geq \beta m(\mathcal{Y}) > 0, \end{aligned} \quad (133)$$

$$\begin{aligned} & \left| \int_{\mathcal{Y}} \hat{w} \left(\hat{D}_\eta^{-1}(x) \left(y - \hat{C}_\eta(x) \right) \right) dy \right| \\ & \leq \int_{\mathcal{Y}} \left| \hat{w} \left(\hat{D}_\eta^{-1}(x) \left(y - \hat{C}_\eta(x) \right) \right) \right| dy \leq \frac{m(\mathcal{Y})}{\beta} \end{aligned} \quad (134)$$

for the same η, x , where $m(\mathcal{Y})$ is the Lebesgue measure of \mathcal{Y} . Further to this, Lemma A1.1 (see Appendix 1) and (128), (130) imply that

$$\int_{\mathcal{X}} \hat{v} \left(\hat{B}_\eta^{-1}(x) \left(x' - \hat{A}_\eta(x) \right) \right) dx', \quad (135)$$

$$\int_{\mathcal{Y}} \hat{w} \left(\hat{D}_\eta^{-1}(x) \left(y - \hat{C}_\eta(x) \right) \right) dy \quad (136)$$

are analytic in (η, x) for $\eta \in V_\alpha(\tilde{\Theta})$, $x \in V_\alpha(\mathcal{X})$.

In the rest of the proof, the following notations is used. $\hat{p}_\eta(x'|x)$, $\hat{q}_\eta(y|x)$ are the functions defined by $\hat{p}_{\eta'}(x'|x) = 0$, $\hat{q}_{\eta'}(y|x) = 0$ and

$$\begin{aligned} \hat{p}_\eta(x'|x) &= \frac{\hat{v} \left(\hat{B}_\eta^{-1}(x) \left(x' - \hat{A}_\eta(x) \right) \right)}{\int_{\mathcal{X}} \hat{v} \left(\hat{B}_\eta^{-1}(x) \left(x'' - \hat{A}_\eta(x) \right) \right) dx''}, \\ \hat{q}_\eta(y|x) &= \frac{\hat{w} \left(\hat{D}_\eta^{-1}(x) \left(y - \hat{C}_\eta(x) \right) \right)}{\int_{\mathcal{Y}} \hat{w} \left(\hat{D}_\eta^{-1}(x) \left(y' - \hat{C}_\eta(x) \right) \right) dy'}. \end{aligned}$$

for $\eta \in V_\alpha(\tilde{\Theta})$, $\eta' \in \mathbb{C}^d \setminus V_\alpha(\tilde{\Theta})$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$, while $r_\theta(y, x'|x)$, $\hat{r}_\eta(y, x'|x)$ are the functions defined by

$$\begin{aligned} r_\theta(y, x'|x) &= q_\theta(y|x') p_\theta(x'|x), \\ \hat{r}_\eta(y, x'|x) &= \hat{q}_\eta(y|x') \hat{p}_\eta(x'|x) \end{aligned}$$

for the same x, x', y and $\theta \in \Theta$, $\eta \in \mathbb{C}^d$.

As functions (125), (126) and integrals (135), (136) are analytic in (η, x, x', y) for $\eta \in V_\alpha(\tilde{\Theta})$, $x, x' \in V_\alpha(\mathcal{X})$, $y \in V_\alpha(\mathcal{Y})$, it follows from (131), (133) that $\hat{r}_\eta(y, x'|x)$ is well-defined and analytic in η for the same η, x, x', y . Similarly, (131) – (134) imply that there exists a real number $\gamma \in (0, 1)$ such that $\gamma \leq |\hat{r}_\eta(y, x'|x)| \leq 1/\gamma$ for $\eta \in V_\alpha(\tilde{\Theta})$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$. Further to this, (ii) yields $\hat{r}_\theta(y, x'|x) = r_\theta(y, x'|x)$ for the same x, x', y and $\theta \in \tilde{\Theta}$. Consequently, Assumptions 2.1 – 2.5 follow from Assumptions 4.1 – 4.4 when Θ is restricted to $\tilde{\Theta}$ (i.e., when Θ is replaced with $\tilde{\Theta}$). Then, using Theorems 2.1, 2.2, we conclude that there exist functions $\tilde{l}, \tilde{h} : \tilde{\Theta} \rightarrow \mathbb{R}$ such that $\tilde{l}(\theta)$, $\tilde{h}(\theta)$ are real-analytic in θ and satisfy $\lim_{n \rightarrow \infty} l_n(\theta, \lambda) = \tilde{l}(\theta)$, $\lim_{n \rightarrow \infty} h_n(\theta, \lambda) = \tilde{h}(\theta)$ for $\theta \in \tilde{\Theta}$, $\lambda \in \mathcal{P}(\mathcal{X})$ ($l_n(\theta, \lambda)$, $h_n(\theta, \lambda)$ have the same meaning as in (1)). Consequently, Corollaries 4.1, 4.2 hold (we use here representation $\Theta = \bigcup_{n=1}^{\infty} \tilde{\Theta}_n$, where $\{\tilde{\Theta}_n\}_{n \geq 1}$ is a sequence of non-empty open balls satisfying $\operatorname{cl} \tilde{\Theta}_n \subset \Theta$ for $n \geq 1$). \square

APPENDIX 1

This section contains some auxiliary results which are relevant for the proof of Lemmas 5.2, 5.3, 6.3, 7.2 and Corollaries 3.1 – 4.2. Here, we rely on the following notations. $d_w \geq 1$ and $d_z \geq 1$ are integers, while A is a bounded convex set in \mathbb{C}^{d_w} . $F(w, z)$ is a function mapping $w \in \mathbb{C}^{d_w}$, $z \in \mathbb{R}^{d_z}$ to \mathbb{C} , while $\lambda(dz)$ is a measure on \mathbb{R}^{d_z} . $f(w)$ is the function defined by

$$f(w) = \int F(w, z) \lambda(dz)$$

for $w \in \mathbb{C}^{d_w}$.

Lemma A1.1. *Assume the following:*

(i) *There exists a real number $\delta \in (0, 1)$ such that $F(w, z)$ is analytic in w for each $w \in V_\delta(A)$, $z \in \mathbb{R}^{d_z}$.*

(ii) *There exists a function $\phi : \mathbb{R}^{d_z} \rightarrow [1, \infty)$ such that $|F(w, z)| \leq \phi(z)$ for all $w \in V_\delta(A)$, $z \in \mathbb{R}^{d_z}$.*

Then, we have

$$|F(w', z) - F(w'', z)| \leq \frac{d_w \phi(z) \|w' - w''\|}{\delta}$$

for all $w', w'' \in V_\delta(A)$, $z \in \mathbb{R}^{d_z}$. Moreover, if $\int \phi(z) \lambda(dz) < \infty$, then $f(w)$ is well-defined and analytic for all $w \in V_\delta(A)$.

Proof. Owing to Cauchy's inequality (see e.g., [28, Proposition 2.1.3]) and (i), (ii), we have

$$\|\nabla_x F(w, z)\| \leq \frac{d_w \phi(z)}{\delta} \quad (137)$$

for $w \in V_\delta(A)$, $z \in \mathbb{R}^{d_z}$. Consequently, we get

$$\begin{aligned} & |F(w', z) - F(w'', z)| \\ &= \left| \int_0^1 (\nabla_w F(tw' + (1-t)w'', z))^T (w' - w'') dt \right| \\ &\leq \int_0^1 \|\nabla_w F(tw' + (1-t)w'', z)\| \|w' - w''\| dt \\ &\leq \frac{d_w \phi(z) \|w' - w''\|}{\delta} \end{aligned}$$

for $w', w'' \in V_\delta(A)$, $z \in \mathbb{R}^{d_z}$, as $tw' + (1-t)w'' \in V_\delta(A)$ results from $t \in [0, 1]$ and the convexity of $V_\delta(A)$. Moreover, if $\int \phi(z) \lambda(dz) < \infty$, the dominated convergence theorem and (137) imply that $f(w)$ is well-defined and differentiable for $w \in V_\delta(A)$. Consequently, $f(w)$ is analytic for $w \in V_\delta(A)$. \square

In the rest of this appendix, we use the following notations. B is a compact set in \mathbb{R}^{d_w} , while $g(w)$ is a function mapping $w \in \mathbb{R}^{d_w}$ to \mathbb{R} (d_w is specified at the beginning in the appendix).

Lemma A1.2. *Assume that there exists an open set C in \mathbb{R}^{d_w} such that $B \subset C$ and $g(w)$ is real-analytic on C . Then, there exists a function $\hat{g}(w)$ with the following properties:*

(i) $\hat{g}(w)$ maps $w \in \mathbb{C}^{d_w}$ to \mathbb{C} .

(ii) $\hat{g}(w) = g(w)$ for all $w \in B$.

(iii) *There exists a real number $\delta \in (0, 1)$ such that $\hat{g}(w)$ is analytic on $V_\delta(B)$.*

Proof. First, we assume that B is connected (latter, this assumption is dropped). As $g(w)$ is real-analytic on C , $g(w)$ has an analytic continuation in an open vicinity of any point in C . Hence, there exist functions $\hat{g}(w, v)$, $\delta(v)$ with the following properties:

(iv) $\hat{g}(w, v)$, $\delta(v)$ map $w \in \mathbb{C}^{d_w}$, $v \in C$ to \mathbb{C} , $(0, 1)$ (respectively).

(v) $\hat{g}(w, v) = g(w)$ for $w \in V_{\delta(v)}(v) \cap \mathbb{R}^{d_w}$, $v \in C$.

(vi) $\hat{g}(w, v)$ is analytic in w for $w \in V_{\delta(v)}(v)$, $v \in C$.

Since B is compact, there exist an integer $M \geq 1$ and points $\{v_i\}_{1 \leq i \leq M}$ such that $v_i \in B$ for $1 \leq i \leq M$ and $B \subset \bigcup_{i=1}^M V_{\delta(v_i)}(v_i)$. Let $\hat{g}_i(w) = \hat{g}(w, v_i)$, $V_i = V_{\delta(v_i)}(v_i)$ for $w \in \mathbb{C}^{d_w}$, $1 \leq i \leq M$. As B is connected, for each $1 \leq i \leq M$, there exists $1 \leq j \leq M$, $j \neq i$ such that $V_i \cap V_j \cap \mathbb{R}^{d_w} \neq \emptyset$. Moreover, if $V_i \cap V_j \cap \mathbb{R}^{d_w} \neq \emptyset$, then $V_i \cap V_j \cap \mathbb{R}^{d_w}$ is a non-empty open set and $\hat{g}_i(w) = \hat{g}_j(w) = g(w)$ for $w \in V_i \cap V_j \cap \mathbb{R}^{d_w}$. Then, by the uniqueness of analytic continuation (see e.g., [15, Corollary 1.2.6]), for each $1 \leq i \leq M$, there exist $1 \leq j \leq M$, $j \neq i$ and a function $\hat{g}_{ij}(w)$ with the following properties:

(vii) $\hat{g}_{ij}(w)$ maps $w \in \mathbb{C}^{d_w}$ to \mathbb{C} .

(viii) $\hat{g}_{ij}(w)$ is analytic on $V_i \cup V_j$.

(ix) $\hat{g}_{ij}(w) = \hat{g}_i(w)$ for $w \in V_i$ and $\hat{g}_{ij}(w) = \hat{g}_j(w)$ for $w \in V_j$.

Following these arguments, we conclude that there exists a function $\hat{g}(w)$ with the following properties:

(x) $\hat{g}(w)$ maps $w \in \mathbb{C}^{d_w}$ to \mathbb{C} .

(xi) $\hat{g}(w)$ is analytic on $\bigcup_{i=1}^M V_i$.

(xii) $\hat{g}(w) = \hat{g}_i(w)$ for $w \in V_i$, $1 \leq i \leq M$.

Now, we drop the assumption that B is connected (i.e., B is any compact set in \mathbb{R}^{d_w}). Since B is compact, there exist an integer $N \geq 1$ and open sets $\{W_i\}_{1 \leq i \leq N}$ in \mathbb{R}^{d_w} such that $W_i \subseteq C$, $B \cap W_i \neq \emptyset$, $W_i \cap W_j = \emptyset$ for $1 \leq i, j \leq N$, $i \neq j$ and $B \subset \bigcup_{i=1}^N W_i$. Let $B_i = B \cap W_i$ for $1 \leq i \leq N$. Hence, $\{B_i\}_{1 \leq i \leq N}$ are connected components of B , and thus, $\{B_i\}_{1 \leq i \leq N}$ are compact and disjoint. Then, according to what has already been shown, there exist open sets $\{U_i\}_{1 \leq i \leq N}$ in \mathbb{C}^{d_w} and functions $\{\hat{g}_i(w)\}_{1 \leq i \leq N}$ with the following properties:

(xiii) $B_i \subset U_i$, $U_i \cap U_j = \emptyset$ for $1 \leq i, j \leq N$, $i \neq j$.

(xiv) $\hat{g}_i(w)$ maps $w \in \mathbb{C}^{d_w}$ to \mathbb{C} for $1 \leq i \leq N$.

(xv) $\hat{g}_i(w) = g(w)$ for $w \in B_i$, $1 \leq i \leq N$.

(xvi) $\hat{g}_i(w)$ is analytic on U_i for $1 \leq i \leq N$.

Let $\hat{g}(w)$ be the function defined by $\hat{g}(w) = \hat{g}_i(w)$ for $w \in U_i$, $1 \leq i \leq N$ and $\hat{g}(w) = 0$ for $w \notin \bigcup_{i=1}^N U_i$. Due to (xiii), $\hat{g}(w)$ is well-defined. As B is compact and $B \subset \bigcup_{i=1}^N U_i$ (owing to (xiii)), there exists a real number $\delta \in (0, 1)$ such that $B \subset V_\delta(B) \subset \bigcup_{i=1}^N U_i$. Then, (xv), (xvi) imply that $\hat{g}(w)$ is analytic on $V_\delta(B)$ and satisfies $\hat{g}(w) = g(w)$ for $w \in B$. \square

APPENDIX 2

In this section, we show how Theorem 2.2 can be applied to finite-state hidden Markov models. We also provide a link between Theorem 2.2 and the results of [12]. Here, we assume that \mathcal{X} has a finite number of elements. We also assume $\mathcal{X} = \{1, \dots, N\}$ and $\mu(x) = 1$ for each $x \in \mathcal{X}$ (in this case, $p_\theta(x'|x)$ is the conditional probability of $X_{n+1}^{\theta, \lambda} = x'$

given $X_n^{\theta, \lambda} = x$). Further to this, we introduce the following assumptions.

Assumption A2.1. $p_\theta(x'|x)$ and $q_\theta(y|x)$ are real-analytic in θ for each $\theta \in \Theta$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$. Moreover, $p_\theta(x'|x)$ and $q_\theta(y|x)$ have complex-valued continuations $\hat{p}_\eta(x'|x)$ and $\hat{q}_\eta(y|x)$ with the following properties:

(i) $\hat{p}_\eta(x'|x)$ and $\hat{q}_\eta(y|x)$ map $\eta \in \mathbb{C}^d$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$ to \mathbb{C} .

(ii) $\hat{p}_\theta(x'|x) = p_\theta(x'|x)$ and $\hat{q}_\theta(y|x) = q_\theta(y|x)$ for all $\theta \in \Theta$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$.

(iii) There exists a real number $\delta \in (0, 1)$ such that $\hat{p}_\eta(x'|x)$ and $\hat{q}_\eta(y|x)$ are analytic in η for each $\eta \in V_\delta(\Theta)$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$.

(iv) There exists a real number $\varepsilon \in (0, 1)$ such that $\varepsilon \leq |\hat{p}_\eta(x'|x)| \leq 1/\varepsilon$ for all $\eta \in V_\delta(\Theta)$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$.

Assumption A2.2. There exists a real number $\alpha \in (0, 1)$ and a vector $\hat{\theta} \in \Theta$ with the following properties:

(i) $q_{\hat{\theta}}(y|x) \neq 0$, $q_\theta(y|x)/q_{\hat{\theta}}(y|x) \geq \alpha$ and $|\hat{q}_\eta(y|x)/q_{\hat{\theta}}(y|x)| \leq 1/\alpha$ for all $\theta \in \Theta$, $\eta \in V_\delta(\Theta)$, $x \in \mathcal{X}$, $y \in \mathcal{Y}$.

(ii) $\int |\log q_{\hat{\theta}}(y|x')| q_{\hat{\theta}}(y|x) \nu(dy) < \infty$ for all $x, x' \in \mathcal{X}$.

Assumption A2.3. There exists a real number $\beta \in (0, 1)$, a vector $\hat{x} \in \mathcal{X}$ and functions $\tilde{\phi}, \tilde{\psi} : \mathcal{Y} \rightarrow (0, \infty)$ with the following properties:

(i) $\hat{q}_\eta(y|\hat{x}) \neq 0$, $|\hat{q}_\eta(y|x)/\hat{q}_\eta(y|\hat{x})| \leq 1/\beta$ for all $\eta \in V_\delta(\Theta)$, $x \in \mathcal{X}$, $y \in \mathcal{Y}$.

(ii) $|\hat{q}_\eta(y|\hat{x})| \leq \tilde{\phi}(y)$ and $|\log |\hat{q}_\eta(y|\hat{x})|| \leq \tilde{\psi}(y)$ for all $\eta \in V_\delta(\Theta)$, $y \in \mathcal{Y}$.

(iii) $\int \tilde{\phi}(y) \nu(dy) < \infty$ and $\int \tilde{\psi}(y) \tilde{\phi}(y) \nu(dy) < \infty$.

Assumptions A2.1 – A2.3 are a particular case of Assumptions 2.1 – 2.4 (see Corollary A2.1 and its proof). At the same time, Assumptions A2.1 – A2.3 include, as a special case, all conditions which the results of [12] are based on.⁵ Further to this, Assumptions A2.1 – A2.3 considerably simplify the conditions adopted in [12].

Corollary A2.1. Let Assumption A2.1 and one of Assumptions A2.2, A2.3 hold. Then, all conclusions of Theorem 2.2 are true.

Proof. It is sufficient to show that Assumptions 2.1 – 2.4 follow from Assumption A2.1 and one of Assumptions A2.2, A2.3.

(i) In this part of the proof, we demonstrate that Assumption 2.1 holds under Assumption A2.1. Let $\lambda_\theta(dx|y)$ be the measure on \mathcal{X} defined by

$$\lambda_\theta(B|y) = \sum_{x \in \mathcal{X}} q_\theta(y|x) I_B(x) \mu(x)$$

for $\theta \in \Theta$, $y \in \mathcal{Y}$, $B \subseteq \mathcal{X}$. Then, Assumption A2.1 implies

$$\begin{aligned} \sum_{x' \in B} r_\theta(y, x'|x) I_B(x') \mu(x') &\geq \varepsilon \sum_{x' \in B} q_\theta(y|x') I_B(x') \mu(x') \\ &= \lambda_\theta(B|y), \end{aligned}$$

⁵Assumptions A2.1 and A2.2 follow (respectively) from [12, Conditions (a), (c.i)] and [12, Conditions (b), (c.iii), Equation (11)], while Assumption A2.3 results from one of [12, Conditions (c.iii), (d.i), Equation (7)] and [12, Conditions (c.ii), (c.iii), (d.i), Equation (8)].

$$\begin{aligned} \sum_{x' \in B} r_\theta(y, x'|x) I_B(x') \mu(x') &\leq \frac{1}{\varepsilon} \sum_{x' \in B} q_\theta(y|x') I_B(x') \mu(x') \\ &= \frac{\lambda_\theta(B|y)}{\varepsilon} \end{aligned}$$

for the same θ, y, B and $x \in \mathcal{X}$. Hence, Assumption 2.1 holds.

(ii) In the next part of the proof, we show that Assumptions 2.2 – 2.4 follow from Assumptions A2.1, A2.2. Let $\tilde{C}_1 = \alpha^{-1} \varepsilon^{-1}$, $\gamma = \alpha^2 \varepsilon^2$, while $\hat{r}_\eta(y, x'|x)$ and $\varphi_\eta(y)$ are the functions defined by

$$\begin{aligned} \hat{r}_\eta(y, x'|x) &= \hat{q}_\eta(y|x') \hat{p}_\eta(x'|x), \\ \varphi_\eta(y) &= \tilde{C}_1 \sum_{x \in \mathcal{X}} q_{\hat{\theta}}(y|x) \end{aligned}$$

for $\eta \in \mathbb{C}^d$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$. Since $\varphi_\eta(y)$ is constant in η , it follows from Assumption A2.1 that $\hat{r}_\eta(y, x'|x)$ and $\varphi_\eta(y)$ are analytic in η for each $\eta \in V_\delta(\Theta)$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$. Moreover, Assumptions A2.1, A2.2 yield $\varphi_\eta(y) \neq 0$ and

$$|\hat{r}_\eta(y, x'|x)| \leq \frac{|\hat{q}_\eta(y|x')|}{\varepsilon} \leq \frac{q_{\hat{\theta}}(y|x')}{\alpha \varepsilon} = |\varphi_\eta(y)| \quad (138)$$

for the same η, x, x', y . Assumptions A2.1, A2.2 also imply

$$\begin{aligned} \sum_{x' \in \mathcal{X}} r_\theta(y, x'|x) \mu(x') &\geq \varepsilon \sum_{x' \in \mathcal{X}} q_\theta(y|x') \geq \alpha \varepsilon \sum_{x' \in \mathcal{X}} q_{\hat{\theta}}(y|x') \\ &= \gamma |\varphi_\theta(y)| \quad (139) \end{aligned}$$

for $\theta \in \Theta$, $x \in \mathcal{X}$, $y \in \mathcal{Y}$.

Let $\tilde{C}_2 = \tilde{C}_1 N$, while $\phi(y)$ and $\psi(y)$ are the functions defined by

$$\begin{aligned} \phi(y) &= \tilde{C}_1 \sum_{x \in \mathcal{X}} q_{\hat{\theta}}(y|x), \\ \psi(y) &= \tilde{C}_2 \left(1 + \sum_{x \in \mathcal{X}} |\log q_{\hat{\theta}}(y|x)| \right) \end{aligned}$$

for $y \in \mathcal{Y}$. Then, due to Part (ii) of Assumption A2.2, we have $\int \phi(y) \nu(dy) = \tilde{C}_1 N < \infty$ and

$$\begin{aligned} \int \psi(y) \phi(y) \nu(dy) &= \tilde{C}_1 \tilde{C}_2 \sum_{x, x' \in \mathcal{X}} \int |\log q_{\hat{\theta}}(y|x')| q_{\hat{\theta}}(y|x) \nu(dy) \\ &\quad + \tilde{C}_1 \tilde{C}_2 N < \infty. \quad (140) \end{aligned}$$

We also have

$$\begin{aligned} \log |\varphi_\eta(y)| &\leq \log(\tilde{C}_1 N) + \max_{x \in \mathcal{X}} \log q_{\hat{\theta}}(y|x) \\ &\leq \tilde{C}_1 N \left(1 + \sum_{x \in \mathcal{X}} |\log q_{\hat{\theta}}(y|x)| \right), \\ \log |\varphi_\eta(y)| &\geq \log(\tilde{C}_1 N) + \min_{x \in \mathcal{X}} \log q_{\hat{\theta}}(y|x) \\ &\geq -\tilde{C}_1 N \left(1 + \sum_{x \in \mathcal{X}} |\log q_{\hat{\theta}}(y|x)| \right) \end{aligned}$$

for $\eta \in V_\delta(\Theta)$, $y \in \mathcal{Y}$. Consequently, we get

$$|\varphi_\eta(y)| \leq \phi(y), \quad |\log |\varphi_\eta(y)|| \leq \psi(y) \quad (141)$$

for the same η, y . Then, using (138) – (141), we conclude that Assumptions 2.2 – 2.4 hold.

(iii) In this part of the proof, we show that Assumptions 2.2 – 2.4 follow from Assumptions A2.1, A2.3. Let $\tilde{C} = \beta^{-1} \varepsilon^{-1}$, $\gamma = \beta \varepsilon^2$, while $\hat{r}_\eta(y, x'|x)$ and $\varphi_\eta(y)$ are the functions defined by

$$\begin{aligned} \hat{r}_\eta(y, x'|x) &= \hat{q}_\eta(y|x') \hat{p}_\eta(x'|x), \\ \varphi_\eta(y) &= \tilde{C} \hat{q}_\eta(y|\hat{x}) \end{aligned}$$

for $\eta \in \mathbb{C}^d$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$. Then, due to Assumption A2.1, $\hat{r}_\eta(y, x'|x)$ and $\varphi_\eta(y)$ are analytic in η for each $\eta \in V_\delta(\Theta)$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$. Moreover, Assumptions A2.1, A2.3 yield $\varphi_\eta(y) \neq 0$ and

$$|\hat{r}_\eta(y, x'|x)| \leq \frac{|\hat{q}_\eta(y|x')|}{\varepsilon} \leq \frac{|\hat{q}_\eta(y|\hat{x})|}{\beta \varepsilon} = |\varphi_\eta(y)| \quad (142)$$

for η, x, x', y . Assumptions A2.1, A2.3 also imply

$$\begin{aligned} \sum_{x' \in \mathcal{X}} r_\theta(y, x'|x) \mu(x') &\geq q_\theta(y|\hat{x}) p_\theta(\hat{x}|x) \geq \varepsilon q_\theta(y|\hat{x}) \\ &= \gamma |\varphi_\theta(y)| \quad (143) \end{aligned}$$

for $\theta \in \Theta$, $x \in \mathcal{X}$, $y \in \mathcal{Y}$.

Let $\phi(y)$ and $\psi(y)$ be the functions defined by

$$\phi(y) = \tilde{C} \tilde{\phi}(y), \quad \psi(y) = \tilde{C} (1 + \tilde{\psi}(y))$$

for $y \in \mathcal{Y}$. Then, Assumption A2.3 yields

$$|\varphi_\eta(y)| = \tilde{C} |\hat{q}_\eta(y|\hat{x})| \leq \tilde{C} \tilde{\phi}(y) = \phi(y) \quad (144)$$

for $\eta \in V_\delta(\Theta)$, $y \in \mathcal{Y}$. Assumption A2.3 also implies

$$\begin{aligned} |\log |\varphi_\eta(y)|| &\leq \log \tilde{C} + |\log |\hat{q}_\eta(y|\hat{x})|| \leq \tilde{C} (1 + \tilde{\psi}(y)) \\ &= \psi(y) \quad (145) \end{aligned}$$

for the same η, y . Then, using Part (iii) of Assumption A2.3, and (142) – (145), we conclude that Assumptions 2.2 – 2.4 hold. \square

In rest of the section, we explain how Theorem 2.2 can further be extended in the context of finite-state hidden Markov models. Here, we rely on the following notations. \mathcal{P}^N is the set of N -dimensional probability vectors, while e is the N -dimensional vector whose all elements are one. For $\theta \in \Theta$, $y \in \mathcal{Y}$, $R_\theta(y)$ is the $N \times N$ matrix whose (x', x) -entry is $r_\theta(y, x'|x)$, where $r_\theta(y, x'|x)$ has the same meaning as in Section II. $G_\theta(\lambda, y)$ and $h_\theta(\lambda, y)$ are the functions defined by

$$G_\theta(\lambda, y) = \frac{R_\theta(y) \lambda}{e^T R_\theta(y) \lambda}, \quad h_\theta(\lambda, y) = \log \left(e^T R_\theta(y) \lambda \right)$$

for $\theta \in \Theta$, $\lambda \in \mathcal{P}^N$, $y \in \mathcal{Y}$. Regarding functions $r_\theta(y, x'|x)$, $G_\theta(\lambda, y)$ and $h_\theta(\lambda, y)$, we assume the following.

Assumption A2.4. *There exist a real number $\varepsilon \in (0, 1)$ and a function $s_\theta(y, x)$ mapping $\theta \in \Theta$, $x \in \mathcal{X}$, $y \in \mathcal{Y}$ to $[0, \infty)$ such that*

$$\varepsilon s_\theta(y, x') \leq r_\theta(y, x'|x) \leq \frac{s_\theta(y, x')}{\varepsilon}$$

for all $\theta \in \Theta$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$.

Assumption A2.5. *$G_\theta(\lambda, y)$ and $h_\theta(\lambda, y)$ are real-analytic in (θ, λ) for all $\theta \in \Theta$, $\lambda \in \mathcal{P}^N$, $y \in \mathcal{Y}$. Moreover, $G_\theta(\lambda, y)$*

and $h_\theta(\lambda, y)$ have complex-valued continuations $\hat{G}_\eta(\zeta, y)$ and $\hat{h}_\eta(\zeta, y)$ with the following properties:

(i) $\hat{G}_\eta(\zeta, y)$ and $\hat{h}_\eta(\zeta, y)$ map $\eta \in \mathbb{C}^d$, $\zeta \in \mathbb{C}^N$, $y \in \mathcal{Y}$ to \mathbb{C}^N and \mathbb{C} (respectively).

(ii) $\hat{G}_\theta(\lambda, y) = G_\theta(\lambda, y)$ and $\hat{h}_\theta(\lambda, y) = h_\theta(\lambda, y)$ for all $\theta \in \Theta$, $\lambda \in \mathcal{P}^N$, $y \in \mathcal{Y}$.

(iii) There exists a real number $\delta \in (0, 1)$ such that $\hat{G}_\eta(\zeta, y)$ and $\hat{h}_\eta(\zeta, y)$ are analytic in (η, ζ) for each $\eta \in V_\delta(\Theta)$, $\zeta \in V_\delta(\mathcal{P}^N)$, $y \in \mathcal{Y}$.

(iv) There exist a real number $K \in [1, \infty)$ and a function $\tilde{\psi} : \mathcal{Y} \rightarrow [1, \infty)$ such that $\int \exp(\tilde{\psi}(y)) \tilde{\psi}(y) \nu(dy) < \infty$ and

$$\|\hat{G}_\eta(\zeta, y)\| \leq K, \quad |\hat{h}_\eta(\zeta, y)| \leq \tilde{\psi}(y)$$

for all $\eta \in V_\delta(\Theta)$, $\zeta \in V_\delta(\mathcal{P}^N)$, $y \in \mathcal{Y}$.

Assumption A2.4 corresponds to the stability of the hidden Markov model $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$ and its optimal filter, while Assumption A2.5 is related to the parameterization of the model $\{(X_n^{\theta, \lambda}, Y_n^{\theta, \lambda})\}_{n \geq 0}$. Assumptions A2.4 and A2.5 are the same as the (corresponding) assumptions adopted in [23]. Further to this, Assumptions A2.4 and A2.5 include, as a particular case, all conditions which the results of [12] are based on.

Theorem A2.1. *Let Assumptions A2.4 and A2.5 hold. Then, all conclusions of Theorem 2.2 are true.*

Proof. Let e_i be the i -th standard unit vector in \mathbb{R}^N , where $1 \leq i \leq N$. Moreover, let $\hat{r}_\eta(y, x'|x)$ be the function defined by

$$\hat{r}_\eta(y, x'|x) = e_{x'}^T \hat{G}_\eta(e_x, y) \exp(\hat{h}_\eta(e_x, y))$$

for $\eta \in \mathbb{C}^d$, $x, x' \in \mathcal{X}$, $y \in \mathcal{Y}$, while $\varphi_\eta(y)$, $\phi(y)$ and $\psi(y)$ are the functions defined by

$$\varphi_\eta(y) = \phi(y) = K \exp(\tilde{\psi}(y)), \quad \psi(y) = 2K \tilde{\psi}(y)$$

for the same η, y . Then, it is straightforward to demonstrate that Assumptions 2.1, 2.2 and 2.4 hold.

Let $\hat{s}_\eta(x)$ be the function defined by

$$\hat{s}_\eta(x) = \sum_{x' \in \mathcal{X}} \int \hat{r}_\eta(y, x'|x) \nu(dy)$$

for $\eta \in \mathbb{C}^d$, $x \in \mathcal{X}$, while $\tilde{r}_\eta(y, x'|x)$ is the function be defined by

$$\tilde{r}_\eta(y, x'|x) = \begin{cases} \hat{r}_\eta(y, x'|x) / \hat{s}_\eta(x), & \text{if } \hat{s}_\eta(x) \neq 0 \\ 0, & \text{otherwise} \end{cases}$$

for the same η, x and $x' \in \mathcal{X}$, $y \in \mathcal{Y}$. Moreover, let $T_\eta(z, B)$ be the kernel defined by

$$T_\eta(z, B) = \sum_{x' \in \mathcal{X}} \int I_B(y', x') \tilde{r}_\eta(y', x'|x) \nu(dy')$$

for $\eta \in \mathbb{C}^d$, $x \in \mathcal{X}$, $y \in \mathcal{Y}$, a Borel-set $B \subseteq \mathcal{Y} \times \mathcal{X}$ and $z = (y, x)$. Since Assumptions 2.1, 2.2 and 2.4 hold, Lemma 5.4 implies that Assumptions 7.1, 7.2 hold, too.

Let $\Phi_\eta(\zeta, z)$ be the function defined by

$$\Phi_\eta(\zeta, z) = \hat{h}_\eta(\zeta, y)$$

for $\eta \in \mathbb{C}^d$, $\zeta \in \mathbb{C}^N$, $x \in \mathcal{X}$, $y \in \mathcal{Y}$ and $z = (y, x)$, while $\{F_{\eta, y}^{m, n}(\zeta)\}_{n \geq m \geq 0}$ are the functions recursively defined by $F_{\eta, y}^{m, m}(\zeta) = \zeta$ and

$$F_{\eta, y}^{m, n+1}(\zeta) = \hat{G}_\eta(F_{\eta, y}^{m, n}(\zeta), y_{n+1})$$

for the same η, ζ and a sequence $y = \{y_n\}_{n \geq 1}$ in \mathcal{Y} . Then, owing to Lemma A1.1, the conclusions of Lemma 6.4 hold. Moreover, due to [23, Lemma 3], the conclusions of Lemma 6.6 also hold provided that elements of \mathbb{C}^N are interpreted as complex measures on \mathcal{X} . Combining Assumptions 7.1, 7.2 and the conclusions of Lemmas 6.4, 6.6, we get the conclusions of Lemmas 7.1, 7.2. Then, as a direct consequence of the conclusions of Lemmas 7.1, 7.2, we get the conclusions of Theorem 2.2. \square

REFERENCES

- [1] L. E. Baum and T. Petrie, "Statistical inference for probabilistic functions of finite state Markov chains," *Ann. Math. Statist.*, vol. 37, no. 6, pp. 1554–1563, 1966.
- [2] O. Cappe, E. Moulines, and T. Ryden, *Inference in Hidden Markov Models*. New York, NY, USA: Springer-Verlag, 2005.
- [3] D. Crisan and B. Rozovskii, *The Oxford Handbook of Nonlinear Filtering*. London, U.K.: Oxford Univ. Press, 2011.
- [4] P. Del Moral and A. Guionnet, "On the stability of interacting processes with applications to filtering and genetic algorithms," *Ann. l'Inst. Henri Poincaré, Probab. Stat.*, vol. 37, no. 2, pp. 155–194, 2001.
- [5] R. Douc, E. Moulines, and T. Rydén, "Asymptotic properties of the maximum likelihood estimator in autoregressive models with Markov regime," *Ann. Statist.*, vol. 32, no. 5, pp. 2254–2304, 2004.
- [6] R. Douc, E. Moulines, and D. S. Stoffer, *Nonlinear Time Series: Theory, Methods and Applications With R Examples*. Boca Raton, FL, USA: CRC Press, 2014.
- [7] T. M. Cover and J. A. Thomas, *Elements of Information Theory*, 2nd ed. Hoboken, NJ, USA: Wiley, 2006.
- [8] Y. Ephraim and N. Merhav, "Hidden Markov processes," *IEEE Trans. Inf. Theory*, vol. 48, no. 6, pp. 1518–1569, Aug. 2002.
- [9] R. M. Gray, *Entropy and Information Theory*, 2nd ed. New York, NY, USA: Springer-Verlag, 2011.
- [10] G. Han and B. Marcus, "Analyticity of entropy rate of hidden Markov chains," *IEEE Trans. Inf. Theory*, vol. 52, no. 12, pp. 5251–5266, Nov. 2006.
- [11] G. Han and B. Marcus, "Derivatives of entropy rate in special families of hidden Markov chains," *IEEE Trans. Inf. Theory*, vol. 53, no. 7, pp. 2642–2652, Jul. 2007.
- [12] G. Han and B. Marcus, "Analyticity of entropy rate of hidden Markov chains with continuous alphabet," *IEEE Trans. Inf. Theory*, vol. 61, no. 6, pp. 3013–3028, Apr. 2015.
- [13] G. Han, "A randomized algorithm for the capacity of finite-state channels," *IEEE Trans. Inf. Theory*, vol. 61, no. 7, pp. 3651–3669, May 2015.
- [14] T. Holliday, A. Goldsmith, and P. Glynn, "Capacity of finite state channels based on Lyapunov exponents of random matrices," *IEEE Trans. Inf. Theory*, vol. 52, no. 8, pp. 3509–3532, Aug. 2006.
- [15] S. G. Krantz and H. R. Parks, *A Primer of Real Analytic Functions*. Basel, Switzerland: Birkhäuser, 2002.
- [16] F. Le Gland and L. Mevel, "Exponential forgetting and geometric ergodicity in hidden Markov models," *Math. Control, Signals Syst.*, vol. 13, no. 1, pp. 63–93, 2000.
- [17] F. Le Gland and N. Oudjane, "Stability and uniform approximation of nonlinear filters using the Hilbert metric and application to particle filters," *Ann. Appl. Probab.*, vol. 14, no. 1, pp. 144–187, 2004.
- [18] S. P. Meyn and R. L. Tweedie, *Markov Chains and Stochastic Stability*, 2nd ed. Cambridge, U.K.: Cambridge Univ. Press, 2009.
- [19] E. Ordentlich and T. Weissman, "On the optimality of symbol-by-symbol filtering and denoising," *IEEE Trans. Inf. Theory*, vol. 52, no. 1, pp. 19–40, Jan. 2006.
- [20] Y. Peres, "Domains of analytic continuation for the top Lyapunov exponent," *Ann. l'Inst. Henri Poincaré, Probab. Stat.*, vol. 28, no. 1, pp. 131–148, 1992.

- [21] G. Poyiadjis, A. Doucet, and S. S. Singh, "Particle approximations of the score and observed information matrix in state space models with application to parameter estimation," *Biometrika*, vol. 98, no. 1, pp. 65–80, 2011.
- [22] A. Schönhuth, "On analytic properties of entropy rate," *IEEE Trans. Inf. Theory*, vol. 55, no. 5, pp. 2119–2127, Apr. 2009.
- [23] V. B. Tadić, "Analyticity, convergence, and convergence rate of recursive maximum-likelihood estimation in hidden Markov models," *IEEE Trans. Inf. Theory*, vol. 56, no. 12, pp. 6406–6432, Dec. 2010.
- [24] V. B. Tadić and A. Doucet, "Exponential forgetting and geometric ergodicity for optimal filtering in general state-space models," *Stochastic Process. Appl.*, vol. 115, no. 8, pp. 1408–1436, 2005.
- [25] V. B. Tadić and A. Doucet, "Asymptotic bias of stochastic gradient search," *Ann. Appl. Probab.*, vol. 27, no. 6, pp. 3255–3304, 2017.
- [26] V. Z. B. Tadić and A. Doucet, "Asymptotic properties of recursive maximum likelihood estimation in non-linear state-space models," 2018, *arXiv:1806.09571*. [Online]. Available: <https://arxiv.org/abs/1806.09571>
- [27] M. Taniguchi and Y. Kakizawa, *Asymptotic Theory of Statistical Inference for Time Series*. New York, NY, USA: Springer, 2000.
- [28] J. L. Taylor, *Several Complex Variables with Connections to Algebraic Geometry and Lie Groups*. Providence, RI, USA: AMS, 2002.

Vladislav Z. B. Tadić obtained B.Sc., M.Sc. and Ph.D. all in electrical engineering from the University of Belgrade in 1989, 1995 and 1997, respectively. During the period 1989–2000, he held various research and teaching positions at the Mihajlo Pupin Institute, Belgrade and the Department of Electrical Engineering, University of Belgrade. From 2000 until 2003, he was a research fellow at the Department of Electrical and Electronic Engineering, University of Melbourne. During the period 2003–2007, he was a lecturer at the Department of Automatic Control and Systems Engineering, University of Sheffield. In 2007, he joined the Department of Mathematics, University of Bristol. His research interests include stochastic approximation, dynamic programming, stochastic optimization and Monte-Carlo methods.

Arnaud Doucet obtained his Ph.D. from University Paris-XI (Orsay) in 1997. He has held ever since faculty positions at the University of Melbourne, Cambridge University, the University of British Columbia and the Institute of Statistical Mathematics in Tokyo. He joined the Department of Statistics of the University of Oxford in 2011 where he is currently a Chair Professor. His research interests include computational statistics, Monte-Carlo methods and statistical machine learning.