

1. Consider a sample  $(X_1, \dots, X_n) = (x_1, \dots, x_n)$  of independent observations from a canonical exponential family with density

$$f(x; \theta) = b(x)e^{\theta^\top t(x) - c(\theta)}.$$

The *reciprocal score*  $R(\theta)$  is defined by

$$\mathcal{R}(\theta) = -\frac{\partial}{\partial \theta} L(\theta)^{-1}$$

i.e. the derivative of the reciprocal of the likelihood function rather than its logarithm.

- (a) Show that

$$R(\theta) = L(\theta)^{-1}\{t(X) - \tau(\theta)\} = L(\theta)^{-1}S(\theta)$$

where  $\tau(\theta) = \mathbf{E}_\theta\{t(X)\}$ .

- (b) Show that using Newton's method on the reciprocal score equation  $R(\theta) = 0$  leads to the iteration

$$\theta \leftarrow \theta + \{v(\theta) + S(\theta)S(\theta)^\top\}^{-1}S(\theta).$$

- (c) Compare this method to the method of scoring.

2. Let  $X = (X_1, \dots, X_n)$  be a sample from the *Weibull* distribution with individual densities

$$f(x; \theta) = \theta x^{\theta-1} e^{-x^\theta} \text{ for } x > 0$$

where  $\theta > 0$  is unknown.

- Find the score statistic and the likelihood equation;
- Show that if the likelihood equation has a solution, it must be the MLE;
- Describe the Newton–Raphson method for solving the likelihood equation;
- Describe the method of scoring for solving the likelihood equation;
- Can you think of other methods for solving the likelihood equation?

3. Consider a sample  $X = (X_1, \dots, X_n)$  from a normal distribution  $\mathcal{N}(\mu, \mu^2)$ , where  $\mu > 0$  is unknown. This corresponds to the coefficient of variation  $\sqrt{\mathbf{V}(X)}/\mathbf{E}(X)$  being known and equal to 1.

- (a) Find the score function for  $\mu$ ;
- (b) Show that the likelihood equation has a unique root  $\hat{\mu}$  within the parameter space unless  $X_i$  are all equal to zero;
- (c) Show that the observed information at  $\hat{\mu}$  is

$$j(\hat{\mu}) = \frac{n}{\hat{\mu}^2} + \frac{\sum_i X_i^2}{\hat{\mu}^4},$$

and use this to argue that the root  $\hat{\mu}$  is indeed the MLE of  $\mu$ ;

- (d) Show that the Fisher information is equal to  $(3n)/\mu^2$ .

4. Let  $X = (X_1, \dots, X_n)$  be a sample from the Gamma distribution with parameters  $\alpha > 0$  and  $\beta > 0$  both unknown, i.e. the distribution with individual densities

$$f(x; \alpha, \beta) = \frac{\beta^\alpha x^{\alpha-1}}{\Gamma(\alpha)} e^{-\beta x}, \quad x > 0.$$

- (a) Show that the score statistic for  $\theta = (\alpha, \beta)$  is equal to

$$S(\alpha, \beta) = \begin{pmatrix} \sum_i \log X_i + n \log \beta - n\psi(\alpha) \\ n\alpha/\beta - \sum_i X_i \end{pmatrix}$$

where  $\psi(\alpha) = D \log \Gamma(\alpha)$  is the *digamma* function;

- (b) Show that the method of scoring for  $\theta$  leads to the iteration

$$\begin{pmatrix} \alpha \\ \beta \end{pmatrix} \leftarrow \begin{pmatrix} \alpha \\ \beta \end{pmatrix} + \frac{1}{n\{\alpha\psi'(\alpha) - 1\}} \begin{pmatrix} \alpha & \beta \\ \beta & \beta^2\psi'(\alpha) \end{pmatrix} S(\alpha, \beta),$$

where  $\psi'(\alpha)$  is the *trigamma* function.

- (c) Consider a simpler iteration for solving the likelihood equation by first eliminating  $\beta$  from the equation.