

# Conditional Independence and Graphical Models

**MSc Further Statistical Methods, Lecture 2**  
**Hilary Term 2005**

Steffen Lauritzen, University of Oxford; January 21, 2005

## Three-way tables


Admissions to Berkeley by department

Department	Sex	Whether admitted	
		Yes	No
I	Male	512	313
	Female	89	19
II	Male	353	207
	Female	17	8
III	Male	120	205
	Female	202	391
IV	Male	138	279
	Female	131	244
V	Male	53	138
	Female	94	299
VI	Male	22	351
	Female	24	317

Here are three variables  $A$ : Admitted?,  $S$ : Sex, and  $D$ : Department.

## Conditional independence

For three variables it is of interest to see whether independence holds for fixed value of one of them, e.g. *is the admission independent of sex for every department separately?* We denote this as  $A \perp\!\!\!\perp S \mid D$  and graphically as



Algebraically, this corresponds to the relations

$$p_{ijk} = p_{i+|k} p_{+j|k} p_{++k} = \frac{p_{i+k} p_{+jk}}{p_{++k}}.$$

## Marginal and conditional independence

Note that there the two conditions


$$A \perp\!\!\!\perp S, \quad A \perp\!\!\!\perp S \mid D$$

are very different and will typically not both hold unless we either have  $A \perp\!\!\!\perp (D, S)$  or  $(A, S) \perp\!\!\!\perp D$ , i.e. if one of the variables are completely independent of both others.

This fact is a simple form of what is known as *Yule–Simpson paradox*. It can be much worse than this:

*A positive conditional association can turn into a negative marginal association and vice-versa.*

## Graphical models



For several variables, complex systems of conditional independence can be described by undirected graphs.

Then a set of variables  $A$  is conditionally independent of set  $B$ , given the values of a set of variables  $C$  if  $C$  separates  $A$  from  $B$ .

For example in picture above

$$1 \perp\!\!\!\perp \{4, 7\} \mid \{2, 3\}, \quad \{1, 2\} \perp\!\!\!\perp 7 \mid \{4, 5, 6\}.$$

Algebraically the picture represents the fact that the joint probability of all variables factorizes into terms that only depends on *cliques* of the graph. In the pictures:

$$p_{ijklmno} = a_{ij}b_{ik}c_{jm}c_{jl}c_{kmn}c_{lo}c_{mno}.$$