

Ordinal variables and measures of association

MSc Further Statistical Methods, Lecture 3
Hilary Term 2005

Steffen Lauritzen, University of Oxford; January 28, 2005

Ordinal variables

Centre	Status	Treatment	Response		
			Poor	Moderate	Excellent
1	1	Active	3	20	5
		Placebo	11	14	8
	2	Active	3	14	12
		Placebo	6	13	5
2	1	Active	12	12	0
		Placebo	11	10	0
	2	active	3	9	4
		Placebo	6	9	3

Multicentre analgesic trial. Here are four variables C :
Centre, S : Status, T : Treatment, and R : Response.

Wilcoxon test-statistic compares distribution of *ranks* between two distributions. Ranks are well-defined for ordinal data.

Several categories

Drug regimen	Response		
	None	Partial	Complete
1	2	0	0
2	1	1	0
3	3	0	0
4	2	2	0
5	1	1	4

Two variables D : Drug regimen, R : response.

Kruskal-Wallis test statistic measure deviations from independence in direction of at *least one distribution stochastically larger* than the others.

Kruskal-Wallis test specializes to Wilcoxon test for binary variables

Two ordinal variables

Income	Job satisfaction			
	Very diss.	Little diss.	Mod. sat.	Very sat.
< 15,000	1	3	10	6
15,000–25,000	2	3	10	7
25,000–40,000	1	6	14	12
> 40,000	0	1	9	11

Two ordinal variables: J : Job satisfaction, I : Income.
Jonckheere-Terpstra test measures deviations from independence in direction of *all distributions being stochastically ordered*.

The Jonckheere–Terpstra test specializes to the Wilcoxon test if one of the two ordinal variables are binary.